

2008 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS

x	$h(x)$	$h'(x)$	$h''(x)$	$h'''(x)$	$h^{(4)}(x)$
1	11	30	42	99	18
2	80	128	$\frac{488}{3}$	$\frac{448}{3}$	$\frac{584}{9}$
3	317	$\frac{753}{2}$	$\frac{1383}{4}$	$\frac{3483}{16}$	$\frac{1125}{16}$

3. Let h be a function having derivatives of all orders for $x > 0$. Selected values of h and its first four derivatives are indicated in the table above. The function h and these four derivatives are increasing on the interval $1 \leq x \leq 3$.
- (a) Write the first-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$. Is this approximation greater than or less than $h(1.9)$? Explain your reasoning.
- (b) Write the third-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$.
- (c) Use the Lagrange error bound to show that the third-degree Taylor polynomial for h about $x = 2$ approximates $h(1.9)$ with error less than 3×10^{-4} .
-

WRITE ALL WORK IN THE PINK EXAM BOOKLET.

END OF PART A OF SECTION II


2008 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS

**CALCULUS BC
SECTION II, Part B**

Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.



Graph of v

4. A particle moves along the x -axis so that its velocity at time t , for $0 \leq t \leq 6$, is given by a differentiable function v whose graph is shown above. The velocity is 0 at $t = 0$, $t = 3$, and $t = 5$, and the graph has horizontal tangents at $t = 1$ and $t = 4$. The areas of the regions bounded by the t -axis and the graph of v on the intervals $[0, 3]$, $[3, 5]$, and $[5, 6]$ are 8, 3, and 2, respectively. At time $t = 0$, the particle is at $x = -2$.
- (a) For $0 \leq t \leq 6$, find both the time and the position of the particle when the particle is farthest to the left. Justify your answer.
- (b) For how many values of t , where $0 \leq t \leq 6$, is the particle at $x = -8$? Explain your reasoning.
- (c) On the interval $2 < t < 3$, is the speed of the particle increasing or decreasing? Give a reason for your answer.
- (d) During what time intervals, if any, is the acceleration of the particle negative? Justify your answer.
-

WRITE ALL WORK IN THE PINK EXAM BOOKLET.

AP[®] CALCULUS BC
2008 SCORING GUIDELINES

Question 3

x	$h(x)$	$h'(x)$	$h''(x)$	$h'''(x)$	$h^{(4)}(x)$
1	11	30	42	99	18
2	80	128	$\frac{488}{3}$	$\frac{448}{3}$	$\frac{584}{9}$
3	317	$\frac{753}{2}$	$\frac{1383}{4}$	$\frac{3483}{16}$	$\frac{1125}{16}$

Let h be a function having derivatives of all orders for $x > 0$. Selected values of h and its first four derivatives are indicated in the table above. The function h and these four derivatives are increasing on the interval $1 \leq x \leq 3$.

- (a) Write the first-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$. Is this approximation greater than or less than $h(1.9)$? Explain your reasoning.
- (b) Write the third-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$.
- (c) Use the Lagrange error bound to show that the third-degree Taylor polynomial for h about $x = 2$ approximates $h(1.9)$ with error less than 3×10^{-4} .

(a) $P_1(x) = 80 + 128(x - 2)$, so $h(1.9) \approx P_1(1.9) = 67.2$

$P_1(1.9) < h(1.9)$ since h' is increasing on the interval $1 \leq x \leq 3$.

$$4 : \begin{cases} 2 : P_1(x) \\ 1 : P_1(1.9) \\ 1 : P_1(1.9) < h(1.9) \text{ with reason} \end{cases}$$

(b) $P_3(x) = 80 + 128(x - 2) + \frac{488}{6}(x - 2)^2 + \frac{448}{18}(x - 2)^3$

$h(1.9) \approx P_3(1.9) = 67.988$

$$3 : \begin{cases} 2 : P_3(x) \\ 1 : P_3(1.9) \end{cases}$$

(c) The fourth derivative of h is increasing on the interval

$1 \leq x \leq 3$, so $\max_{1.9 \leq x \leq 2} |h^{(4)}(x)| = \frac{584}{9}$.

Therefore, $|h(1.9) - P_3(1.9)| \leq \frac{584}{9} \frac{|1.9 - 2|^4}{4!}$
 $= 2.7037 \times 10^{-4}$
 $< 3 \times 10^{-4}$

$$2 : \begin{cases} 1 : \text{form of Lagrange error estimate} \\ 1 : \text{reasoning} \end{cases}$$