

1998 Calculus BC Free-Response Questions

CALCULUS BC

Section II

Time—1 hour and 30 minutes

Number of problems—6

Percent of total grade—50

A GRAPHING CALCULATOR IS REQUIRED FOR SOME PROBLEMS OR PARTS OF PROBLEMS ON THIS SECTION OF THE EXAMINATION.

REMEMBER TO SHOW YOUR SETUPS AS DESCRIBED IN THE GENERAL INSTRUCTIONS.

-
1. Let R be the region in the first quadrant bounded by the graph of $y = 8 - x^{\frac{3}{2}}$, the x -axis, and the y -axis.
- (a) Find the area of the region R .
 - (b) Find the volume of the solid generated when R is revolved about the x -axis.
 - (c) The vertical line $x = k$ divides the region R into two regions such that when these two regions are revolved about the x -axis, they generate solids with equal volumes. Find the value of k .
-

GO ON TO THE NEXT PAGE 

1998 Calculus BC Free-Response Questions


2. Let f be the function given by $f(x) = 2xe^{2x}$.
- (a) Find $\lim_{x \rightarrow -\infty} f(x)$ and $\lim_{x \rightarrow \infty} f(x)$.
 - (b) Find the absolute minimum value of f . Justify that your answer is an absolute minimum.
 - (c) What is the range of f ?
 - (d) Consider the family of functions defined by $y = bxe^{bx}$, where b is a nonzero constant. Show that the absolute minimum value of bxe^{bx} is the same for all nonzero values of b .
-

GO ON TO THE NEXT PAGE 

1998 Calculus BC Scoring Guidelines

1. Let R be the region in the first quadrant bounded by the graph of $y = 8 - x^{3/2}$, the x -axis, and the y -axis.
- (a) Find the area of the region R .
- (b) Find the volume of the solid generated when R is revolved about the x -axis.
- (c) The vertical line $x = k$ divides the region R into two regions such that when these two regions are revolved about the x -axis, they generate solids with equal volumes. Find the value of k .

(a)



$$A = \int_0^4 (8 - x^{3/2}) dx$$

$$= 8x - \frac{2}{5}x^{5/2} \Big|_0^4 = 32 - \frac{64}{5} = \frac{96}{5} = 19.2$$

(b) $V = \pi \int_0^4 (8 - x^{3/2})^2 dx$

$$= \frac{576\pi}{5} = 115.2\pi \approx 361.911$$

(c) $\pi \int_0^k (8 - x^{3/2})^2 dx = \frac{115.2\pi}{2}$

[or $\pi \int_0^k (8 - x^{3/2})^2 dx = \pi \int_k^4 (8 - x^{3/2})^2 dx$]

$$\int_0^k (8 - x^{3/2})^2 dx = 57.6$$

$$\int_0^k (64 - 16x^{3/2} + x^3) dx = 57.6$$

$$64k - \frac{32}{5}k^{5/2} + \frac{k^4}{4} = 57.6$$

$$k \approx 0.995 \text{ or } 0.994$$

3 { 2: integral
1: integrand
1: limits
1: answer

3 { 2: integral
1: integrand
1: limits and constant
1: answer

3 { 1: integral with k in limits
1: equates volumes
1: answer

Note: 0/1 for answer in each part if no setup points earned