

2016 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS**CHEMISTRY****Section II****7 Questions****Time—1 hour and 45 minutes****YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.**

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

Write your response in the space provided following each question. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.


1. A student investigates the enthalpy of solution, ΔH_{soln} , for two alkali metal halides, LiCl and NaCl. In addition to the salts, the student has access to a calorimeter, a balance with a precision of ± 0.1 g, and a thermometer with a precision of $\pm 0.1^\circ\text{C}$.
 - (a) To measure ΔH_{soln} for LiCl, the student adds 100.0 g of water initially at 15.0°C to a calorimeter and adds 10.0 g of LiCl(s), stirring to dissolve. After the LiCl dissolves completely, the maximum temperature reached by the solution is 35.6°C .
 - (i) Calculate the magnitude of the heat absorbed by the solution during the dissolution process, assuming that the specific heat capacity of the solution is $4.18 \text{ J/(g}\cdot^\circ\text{C)}$. Include units with your answer.
 - (ii) Determine the value of ΔH_{soln} for LiCl in kJ/mol_{rxn} .

To explain why ΔH_{soln} for NaCl is different than that for LiCl, the student investigates factors that affect ΔH_{soln} and finds that ionic radius and lattice enthalpy (which can be defined as the ΔH associated with the separation of a solid crystal into gaseous ions) contribute to the process. The student consults references and collects the data shown in the table below.

Ion	Ionic Radius (pm)
Li ⁺	76
Na ⁺	102

2016 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

- (b) Write the complete electron configuration for the Na^+ ion in the ground state.
- (c) Using principles of atomic structure, explain why the Na^+ ion is larger than the Li^+ ion.
- (d) Which salt, LiCl or NaCl , has the greater lattice enthalpy? Justify your answer.
- (e) Below is a representation of a portion of a crystal of LiCl . Identify the ions in the representation by writing the appropriate formulas (Li^+ or Cl^-) in the boxes below.



- (f) The lattice enthalpy of LiCl is positive, indicating that it takes energy to break the ions apart in LiCl . However, the dissolution of LiCl in water is an exothermic process. Identify all particle-particle interactions that contribute significantly to the dissolution process being exothermic. For each interaction, include the particles that interact and the specific type of intermolecular force between those particles.

AP[®] CHEMISTRY

2016 SCORING GUIDELINES

Question 1

A student investigates the enthalpy of solution, ΔH_{soln} , for two alkali metal halides, LiCl and NaCl. In addition to the salts, the student has access to a calorimeter, a balance with a precision of ± 0.1 g, and a thermometer with a precision of $\pm 0.1^\circ\text{C}$.

- (a) To measure ΔH_{soln} for LiCl, the student adds 100.0 g of water initially at 15.0°C to a calorimeter and adds 10.0 g of LiCl(s), stirring to dissolve. After the LiCl dissolves completely, the maximum temperature reached by the solution is 35.6°C .

- (i) Calculate the magnitude of the heat absorbed by the solution during the dissolution process, assuming that the specific heat capacity of the solution is $4.18 \text{ J}/(\text{g}\cdot^\circ\text{C})$. Include units with your answer.

$q = mc\Delta T = (110.0 \text{ g})(4.18 \text{ J}/(\text{g}\cdot^\circ\text{C}))(35.6^\circ\text{C} - 15.0^\circ\text{C})$ $= 9,470 \text{ J} = 9.47 \text{ kJ}$	<p>1 point is earned for the correct setup.</p> <p>1 point is earned for the correct answer with units.</p>
---	---

- (ii) Determine the value of ΔH_{soln} for LiCl in $\text{kJ}/\text{mol}_{rxn}$.

$10.0 \text{ g LiCl} \times \frac{1 \text{ mol LiCl}}{42.39 \text{ g LiCl}} = 0.236 \text{ mol LiCl}$ $\frac{-9.47 \text{ kJ}}{0.236 \text{ mol LiCl}} = -40.1 \text{ kJ}/\text{mol}_{rxn}$	<p>1 point is earned for the number of moles of LiCl.</p> <p>1 point is earned for the correct ΔH_{soln} <u>and</u> the correct sign.</p>
---	--

To explain why ΔH_{soln} for NaCl is different than that for LiCl, the student investigates factors that affect ΔH_{soln} and finds that ionic radius and lattice enthalpy (which can be defined as the ΔH associated with the separation of a solid crystal into gaseous ions) contribute to the process. The student consults references and collects the data shown in the table below.

Ion	Ionic Radius (pm)
Li^+	76
Na^+	102

- (b) Write the complete electron configuration for the Na^+ ion in the ground state.

$1s^2 2s^2 2p^6$	1 point is earned for the complete correct configuration.
------------------	---

AP[®] CHEMISTRY
2016 SCORING GUIDELINES

Question 1 (continued)


- (c) Using principles of atomic structure, explain why the Na^+ ion is larger than the Li^+ ion.

The valence electrons in the Na^+ ion are in a higher principal energy level than the valence electrons in the Li^+ ion. Electrons in higher principal energy levels are, on average, farther from the nucleus.	1 point is earned for a correct explanation based on occupied principal energy levels.
---	--

- (d) Which salt, LiCl or NaCl , has the greater lattice enthalpy? Justify your answer.

LiCl . Because the Li^+ ion is smaller than the Na^+ ion, the Coulombic attractions between ions in LiCl are stronger than in NaCl . This results in a greater lattice enthalpy.	1 point is earned for the correct choice and justification.
---	---

- (e) Below is a representation of a portion of a crystal of LiCl . Identify the ions in the representation by writing the appropriate formulas (Li^+ or Cl^-) in the boxes below.



See diagram above.	1 point is earned for both identifications.
--------------------	---

- (f) The lattice enthalpy of LiCl is positive, indicating that it takes energy to break the ions apart. in LiCl . However, the dissolution of LiCl in water is an exothermic process. Identify all particle-particle interactions that contribute significantly to the exothermic dissolution process being exothermic. For each interaction, include the particles that interact and the specific type of intermolecular force between those particles.

There are interactions between Li^+ ions and polar water molecules and between Cl^- ions and polar water molecules. These are ion-dipole interactions.	1 point is earned for identifying the particles that interact. 1 point is earned for correctly identifying the type of interaction.
---	--