

2007 AP[®] CALCULUS AB FREE-RESPONSE QUESTIONS

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	6	4	2	5
2	9	2	3	1
3	10	-4	4	2
4	-1	3	6	7

3. The functions f and g are differentiable for all real numbers, and g is strictly increasing. The table above gives values of the functions and their first derivatives at selected values of x . The function h is given by $h(x) = f(g(x)) - 6$.
- (a) Explain why there must be a value r for $1 < r < 3$ such that $h(r) = -5$.
- (b) Explain why there must be a value c for $1 < c < 3$ such that $h'(c) = -5$.
- (c) Let w be the function given by $w(x) = \int_1^{g(x)} f(t) dt$. Find the value of $w'(3)$.
- (d) If g^{-1} is the inverse function of g , write an equation for the line tangent to the graph of $y = g^{-1}(x)$ at $x = 2$.
-

WRITE ALL WORK IN THE PINK EXAM BOOKLET.

END OF PART A OF SECTION II

2007 AP[®] CALCULUS AB FREE-RESPONSE QUESTIONS**CALCULUS AB
SECTION II, Part B****Time—45 minutes****Number of problems—3****No calculator is allowed for these problems.**

-
4. A particle moves along the x -axis with position at time t given by $x(t) = e^{-t} \sin t$ for $0 \leq t \leq 2\pi$.
- (a) Find the time t at which the particle is farthest to the left. Justify your answer.
- (b) Find the value of the constant A for which $x(t)$ satisfies the equation $Ax''(t) + x'(t) + x(t) = 0$ for $0 < t < 2\pi$.
-

t (minutes)	0	2	5	7	11	12
$r'(t)$ (feet per minute)	5.7	4.0	2.0	1.2	0.6	0.5

5. The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function r of time t , where t is measured in minutes. For $0 < t < 12$, the graph of r is concave down. The table above gives selected values of the rate of change, $r'(t)$, of the radius of the balloon over the time interval $0 \leq t \leq 12$. The radius of the balloon is 30 feet when $t = 5$. (Note: The volume of a sphere of radius r is given by $V = \frac{4}{3}\pi r^3$.)
- (a) Estimate the radius of the balloon when $t = 5.4$ using the tangent line approximation at $t = 5$. Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when $t = 5$. Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate $\int_0^{12} r'(t) dt$. Using correct units, explain the meaning of $\int_0^{12} r'(t) dt$ in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than $\int_0^{12} r'(t) dt$? Give a reason for your answer.
-

WRITE ALL WORK IN THE PINK EXAM BOOKLET.

AP[®] CALCULUS AB
2007 SCORING GUIDELINES

Question 3

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	6	4	2	5
2	9	2	3	1
3	10	-4	4	2
4	-1	3	6	7

The functions f and g are differentiable for all real numbers, and g is strictly increasing. The table above gives values of the functions and their first derivatives at selected values of x . The function h is given by $h(x) = f(g(x)) - 6$.

- (a) Explain why there must be a value r for $1 < r < 3$ such that $h(r) = -5$.
- (b) Explain why there must be a value c for $1 < c < 3$ such that $h'(c) = -5$.
- (c) Let w be the function given by $w(x) = \int_1^{g(x)} f(t) dt$. Find the value of $w'(3)$.
- (d) If g^{-1} is the inverse function of g , write an equation for the line tangent to the graph of $y = g^{-1}(x)$ at $x = 2$.

- (a) $h(1) = f(g(1)) - 6 = f(2) - 6 = 9 - 6 = 3$
 $h(3) = f(g(3)) - 6 = f(4) - 6 = -1 - 6 = -7$
 Since $h(3) < -5 < h(1)$ and h is continuous, by the Intermediate Value Theorem, there exists a value r , $1 < r < 3$, such that $h(r) = -5$.

- (b) $\frac{h(3) - h(1)}{3 - 1} = \frac{-7 - 3}{3 - 1} = -5$
 Since h is continuous and differentiable, by the Mean Value Theorem, there exists a value c , $1 < c < 3$, such that $h'(c) = -5$.

- (c) $w'(3) = f(g(3)) \cdot g'(3) = f(4) \cdot 2 = -2$

- (d) $g(1) = 2$, so $g^{-1}(2) = 1$.
 $(g^{-1})'(2) = \frac{1}{g'(g^{-1}(2))} = \frac{1}{g'(1)} = \frac{1}{5}$
 An equation of the tangent line is $y - 1 = \frac{1}{5}(x - 2)$.

$$2 : \begin{cases} 1 : h(1) \text{ and } h(3) \\ 1 : \text{conclusion, using IVT} \end{cases}$$

$$2 : \begin{cases} 1 : \frac{h(3) - h(1)}{3 - 1} \\ 1 : \text{conclusion, using MVT} \end{cases}$$

$$2 : \begin{cases} 1 : \text{apply chain rule} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : g^{-1}(2) \\ 1 : (g^{-1})'(2) \\ 1 : \text{tangent line equation} \end{cases}$$