

2000 AP[®] STATISTICS FREE-RESPONSE QUESTIONS

STATISTICS

SECTION II

Part B

Question 6

Spend about 25 minutes on this part of the exam.

Percent of Section II grade—25

Show all your work. Indicate clearly the methods you use, because you will be graded on the correctness of your methods as well as on the accuracy of your results and explanation.

6. A random sample of 400 married couples was selected from a large population of married couples.
- Heights of married men are approximately normally distributed with mean 70 inches and standard deviation 3 inches.
 - Heights of married women are approximately normally distributed with mean 65 inches and standard deviation 2.5 inches.
 - There were 20 couples in which the wife was taller than her husband, and there were 380 couples in which the wife was shorter than her husband.
- (a) Find a 95 percent confidence interval for the proportion of married couples in the population for which the wife is taller than her husband. Interpret your interval in the context of this question.

2000 AP[®] STATISTICS FREE-RESPONSE QUESTIONS

- (b) Suppose that a married man is selected at random and a married woman is selected at random. Find the approximate probability that the woman will be taller than the man.

- (c) Based on your answers to (a) and (b), are the heights of wives and their husbands independent? Explain your reasoning.

2000 AP[®] STATISTICS FREE-RESPONSE QUESTIONS




Table entry for z is the probability lying below z .

Table A (Continued)

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

AP[®] Statistics 2000 — Scoring Guidelines

Question 6

Solution

- a. Large sample confidence interval for a population proportion. Assumptions: large sample. Here,

$$n\hat{p} = 20 \geq 5 \quad (\text{or } 10) \quad n(1 - \hat{p}) = 380 \geq 5 \quad (\text{or } 10)$$

or

$$\hat{p} \pm 3\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \quad \text{is in the interval } (0,1)$$

$$\hat{p} \pm 1.96\sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

$$.05 \pm 1.96\sqrt{\frac{(.05)(.95)}{400}}$$

$$.05 \pm .02146$$

$$(.02854, .07146)$$

Calculator solution: (.02864, .07136), but still need to name the interval used and check assumptions.

Interpretation: Based on this sample, we can be 95% confident that the proportion of married couples for which the wife is taller than her husband is between .028 and .071.