

2016 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS




Figure 1. Detectability of eDNA fragments of varying lengths




Figure 2. Map of the waterways that connect a nearby river system to Lake Michigan

6. Living and dead organisms continuously shed DNA fragments, known as eDNA, into the environment. To detect eDNA fragments in the environment, the polymerase chain reaction (PCR) can be used to amplify specific eDNA fragments. eDNA fragments of different lengths persist in the environment for varying amounts of time before becoming undetectable (Figure 1).

To investigate whether silver carp, an invasive fish, have moved from a nearby river system into Lake Michigan, researchers tested water samples for the presence of eDNA specific to silver carp (Figure 2).

- Justify** the use of eDNA sampling as an appropriate technique for detecting the presence of silver carp in an environment where many different species of fish are found. **Propose** ONE advantage of identifying long eDNA fragments as opposed to short fragments for detecting silver carp.
- The researchers tested a large number of water samples from Lake Michigan and found eDNA specific to silver carp in a single sample in the lake, as indicated in Figure 2. The researchers concluded that the single positive sample was a false positive and that no silver carp had entered Lake Michigan. **Provide reasoning** other than human error to support the researchers' claim.

AP[®] BIOLOGY
2016 SCORING GUIDELINES

Question 6




Figure 1. Detectability of eDNA fragments of varying lengths




Figure 2. Map of the waterways that connect a nearby river system to Lake Michigan

Living and dead organisms continuously shed DNA fragments, known as eDNA, into the environment. To detect eDNA fragments in the environment, the polymerase chain reaction (PCR) can be used to amplify specific eDNA fragments. eDNA fragments of different lengths persist in the environment for varying amounts of time before becoming undetectable (Figure 1).

To investigate whether silver carp, an invasive fish, have moved from a nearby river system into Lake Michigan, researchers tested water samples for the presence of eDNA specific to silver carp (Figure 2).

- (a) **Justify** the use of eDNA sampling as an appropriate technique for detecting the presence of silver carp in an environment where many different species of fish are found. **Propose** ONE advantage of identifying long eDNA fragments as opposed to short fragments for detecting silver carp. **(2 points)**

Justify (1 point)

- eDNA allows detection of the fish without visual identification/catching the fish.

Proposed advantage (1 point)

- Longer fragments indicate more recent presence of fish.
- Longer fragments are more likely to contain a sequence that is specific to silver carp.
- Longer sequences/more base pairs may increase accuracy/specificity/confidence that the eDNA is from a silver carp and not a related species.

- (b) The researchers tested a large number of water samples from Lake Michigan and found eDNA specific to silver carp in a single sample in the lake, as indicated in Figure 2. The researchers concluded that the single positive sample was a false positive and that no silver carp had entered Lake Michigan.

Provide reasoning other than human error to support the researchers' claim. **(1 point)**

Reasoning (1 point)

- eDNA entered the lake by means other than the fish (e.g., river flow, boats, waste from predators).