CALCULUS BC
SECTION II, Part B
Time—60 minutes
Number of problems—4

No calculator is allowed for these problems.



Graph of f

3. The function f is defined on the closed interval $[-5, 4]$. The graph of f consists of three line segments and is shown in the figure above. Let g be the function defined by $g(x) = \int_{-3}^{x} f(t)\,dt$.

(a) Find $g(3)$.

(b) On what open intervals contained in $-5 < x < 4$ is the graph of g both increasing and concave down? Give a reason for your answer.

(c) The function h is defined by $h(x) = \dfrac{g(x)}{5x}$. Find $h'(3)$.

(d) The function p is defined by $p(x) = f\left(x^2 - x\right)$. Find the slope of the line tangent to the graph of p at the point where $x = -1$.

t (minutes)	0	2	5	8	12
$v_A(t)$ (meters/minute)	0	100	40	–120	–150

4. Train A runs back and forth on an east-west section of railroad track. Train A's velocity, measured in meters per minute, is given by a differentiable function $v_A(t)$, where time t is measured in minutes. Selected values for $v_A(t)$ are given in the table above.

   (a) Find the average acceleration of train A over the interval $2 \le t \le 8$.

   (b) Do the data in the table support the conclusion that train A's velocity is -100 meters per minute at some time t with $5 < t < 8$? Give a reason for your answer.

   (c) At time $t = 2$, train A's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train A, in meters from the Origin Station, at time $t = 12$. Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time $t = 12$.

   (d) A second train, train B, travels north from the Origin Station. At time t the velocity of train B is given by $v_B(t) = -5t^2 + 60t + 25$, and at time $t = 2$ the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train A and train B is changing at time $t = 2$.

Question 3

The function f is defined on the closed interval $[-5, 4]$. The graph of f consists of three line segments and is shown in the figure above.



Graph of f

Let g be the function defined by $g(x) = \int_{-3}^{x} f(t)\, dt$.

(a) Find $g(3)$.

(b) On what open intervals contained in $-5 < x < 4$ is the graph of g both increasing and concave down? Give a reason for your answer.

(c) The function h is defined by $h(x) = \dfrac{g(x)}{5x}$. Find $h'(3)$.

(d) The function p is defined by $p(x) = f\left(x^2 - x\right)$. Find the slope of the line tangent to the graph of p at the point where $x = -1$.

(a) $g(3) = \int_{-3}^{3} f(t)\, dt = 6 + 4 - 1 = 9$

1 : answer

(b) $g'(x) = f(x)$

The graph of g is increasing and concave down on the intervals $-5 < x < -3$ and $0 < x < 2$ because $g' = f$ is positive and decreasing on these intervals.

$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

(c) $h'(x) = \dfrac{5xg'(x) - g(x)5}{(5x)^2} = \dfrac{5xg'(x) - 5g(x)}{25x^2}$

$$h'(3) = \frac{(5)(3)g'(3) - 5g(3)}{25 \cdot 3^2} = \frac{15(-2) - 5(9)}{225} = \frac{-75}{225} = -\frac{1}{3}$$

$3 : \begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$

(d) $p'(x) = f'\left(x^2 - x\right)(2x - 1)$

$p'(-1) = f'(2)(-3) = (-2)(-3) = 6$

$3 : \begin{cases} 2 : p'(x) \\ 1 : \text{answer} \end{cases}$