CALCULUS BC
SECTION II, Part B
Time—60 minutes
Number of problems—4

No calculator is allowed for these problems.

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	–220	150

3. Johanna jogs along a straight path. For $0 \le t \le 40$, Johanna's velocity is given by a differentiable function v. Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

   (a) Use the data in the table to estimate the value of $v'(16)$.

   (b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)|\, dt$ in the context of the problem. Approximate the value of $\int_0^{40} |v(t)|\, dt$ using a right Riemann sum with the four subintervals indicated in the table.

   (c) Bob is riding his bicycle along the same path. For $0 \le t \le 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute. Find Bob's acceleration at time $t = 5$.

   (d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \le t \le 10$.

4. Consider the differential equation $\frac{dy}{dx} = 2x - y$.

   (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.



   (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y. Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.

   (c) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(2) = 3$. Does f have a relative minimum, a relative maximum, or neither at $x = 2$? Justify your answer.

   (d) Find the values of the constants m and b for which $y = mx + b$ is a solution to the differential equation.

GO ON TO THE NEXT PAGE.

AP® CALCULUS AB/CALCULUS BC
2015 SCORING GUIDELINES

Question 3

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	–220	150

Johanna jogs along a straight path. For $0 \le t \le 40$, Johanna's velocity is given by a differentiable function v. Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

(a) Use the data in the table to estimate the value of $v'(16)$.

(b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)|\, dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)|\, dt$ using a right Riemann sum with the four subintervals indicated in the table.

(c) Bob is riding his bicycle along the same path. For $0 \le t \le 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute.

Find Bob's acceleration at time $t = 5$.

(d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \le t \le 10$.

(a) $v'(16) \approx \dfrac{240 - 200}{20 - 12} = 5$ meters/min^2

1 : approximation

(b) $\int_0^{40} |v(t)|\, dt$ is the total distance Johanna jogs, in meters, over the time interval $0 \le t \le 40$ minutes.

$$\begin{aligned}\int_0^{40} |v(t)|\, dt &\approx 12 \cdot |v(12)| + 8 \cdot |v(20)| + 4 \cdot |v(24)| + 16 \cdot |v(40)| \\ &= 12 \cdot 200 + 8 \cdot 240 + 4 \cdot 220 + 16 \cdot 150 \\ &= 2400 + 1920 + 880 + 2400 \\ &= 7600 \text{ meters}\end{aligned}$$

$$3 : \begin{cases} 1 : \text{explanation} \\ 1 : \text{right Riemann sum} \\ 1 : \text{approximation} \end{cases}$$

(c) Bob's acceleration is $B'(t) = 3t^2 - 12t$.

$B'(5) = 3(25) - 12(5) = 15$ meters/min^2

$$2 : \begin{cases} 1 : \text{uses } B'(t) \\ 1 : \text{answer} \end{cases}$$

(d)
$$\begin{aligned}\text{Avg vel} &= \frac{1}{10}\int_0^{10} \left(t^3 - 6t^2 + 300\right) dt \\ &= \frac{1}{10}\left[\frac{t^4}{4} - 2t^3 + 300t\right]_0^{10} \\ &= \frac{1}{10}\left[\frac{10000}{4} - 2000 + 3000\right] = 350 \text{ meters/min}\end{aligned}$$

$$3 : \begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$