


2003 AP[®] CALCULUS AB FREE-RESPONSE QUESTIONS

CALCULUS AB
SECTION II, Part B

Time—45 minutes


Number of problems—3

No calculator is allowed for these problems.



4. Let f be a function defined on the closed interval $-3 \leq x \leq 4$ with $f(0) = 3$. The graph of f' , the derivative of f , consists of one line segment and a semicircle, as shown above.
- (a) On what intervals, if any, is f increasing? Justify your answer.
 - (b) Find the x -coordinate of each point of inflection of the graph of f on the open interval $-3 < x < 4$. Justify your answer.
 - (c) Find an equation for the line tangent to the graph of f at the point $(0, 3)$.
 - (d) Find $f(-3)$ and $f(4)$. Show the work that leads to your answers.
-

2003 AP[®] CALCULUS AB FREE-RESPONSE QUESTIONS



5. A coffeepot has the shape of a cylinder with radius 5 inches, as shown in the figure above. Let h be the depth of the coffee in the pot, measured in inches, where h is a function of time t , measured in seconds. The volume V of coffee in the pot is changing at the rate of $-5\pi\sqrt{h}$ cubic inches per second. (The volume V of a cylinder with radius r and height h is $V = \pi r^2 h$.)

(a) Show that $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$.

(b) Given that $h = 17$ at time $t = 0$, solve the differential equation $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$ for h as a function of t .

- (c) At what time t is the coffeepot empty?

6. Let f be the function defined by

$$f(x) = \begin{cases} \sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ 5-x & \text{for } 3 < x \leq 5. \end{cases}$$

- (a) Is f continuous at $x = 3$? Explain why or why not.
 (b) Find the average value of $f(x)$ on the closed interval $0 \leq x \leq 5$.
 (c) Suppose the function g is defined by

$$g(x) = \begin{cases} k\sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ mx + 2 & \text{for } 3 < x \leq 5, \end{cases}$$

where k and m are constants. If g is differentiable at $x = 3$, what are the values of k and m ?


END OF EXAMINATION

Copyright © 2003 by College Entrance Examination Board. All rights reserved.
 Available to AP professionals at apcentral.collegeboard.com and to
 students and parents at www.collegeboard.com/apstudents.

AP[®] CALCULUS AB
2003 SCORING GUIDELINES

Question 4

Let f be a function defined on the closed interval $-3 \leq x \leq 4$ with $f(0) = 3$. The graph of f' , the derivative of f , consists of one line segment and a semicircle, as shown above.



- (a) On what intervals, if any, is f increasing? Justify your answer.
 (b) Find the x -coordinate of each point of inflection of the graph of f on the open interval $-3 < x < 4$. Justify your answer.
 (c) Find an equation for the line tangent to the graph of f at the point $(0, 3)$.
 (d) Find $f(-3)$ and $f(4)$. Show the work that leads to your answers.

- (a) The function f is increasing on $[-3, -2]$ since $f' > 0$ for $-3 \leq x < -2$.

2 : $\begin{cases} 1 : \text{interval} \\ 1 : \text{reason} \end{cases}$

- (b) $x = 0$ and $x = 2$
 f' changes from decreasing to increasing at $x = 0$ and from increasing to decreasing at $x = 2$

2 : $\begin{cases} 1 : x = 0 \text{ and } x = 2 \text{ only} \\ 1 : \text{justification} \end{cases}$

- (c) $f'(0) = -2$
 Tangent line is $y = -2x + 3$.

1 : equation

- (d) $f(0) - f(-3) = \int_{-3}^0 f'(t) dt$
 $= \frac{1}{2}(1)(1) - \frac{1}{2}(2)(2) = -\frac{3}{2}$

$$f(-3) = f(0) + \frac{3}{2} = \frac{9}{2}$$

$$f(4) - f(0) = \int_0^4 f'(t) dt$$

$$= -\left(8 - \frac{1}{2}(2)^2\pi\right) = -8 + 2\pi$$

$$f(4) = f(0) - 8 + 2\pi = -5 + 2\pi$$

$\begin{cases} 1 : \pm \left(\frac{1}{2} - 2\right) \\ \text{(difference of areas of triangles)} \\ 1 : \text{answer for } f(-3) \text{ using FTC} \\ 4 : \begin{cases} 1 : \pm \left(8 - \frac{1}{2}(2)^2\pi\right) \\ \text{(area of rectangle} \\ \text{— area of semicircle)} \\ 1 : \text{answer for } f(4) \text{ using FTC} \end{cases} \end{cases}$