



2017 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

7. A student wants to determine the concentration of H_2O_2 in a solution of $\text{H}_2\text{O}_2(aq)$. The student can use one of two titrants, either dichromate ion, $\text{Cr}_2\text{O}_7^{2-}(aq)$, or cobalt(II) ion, $\text{Co}^{2+}(aq)$. The balanced chemical equations for the two titration reactions are shown below.



The half-reactions and the E° values for the systems related to the titrations above are given in the following table.

Half-Reaction	E° (V) at 298 K
$\text{Co}^{3+}(aq) + e^- \rightarrow \text{Co}^{2+}(aq)$	1.84
$\text{H}_2\text{O}_2(aq) + 2 \text{H}^+(aq) + 2 e^- \rightarrow 2 \text{H}_2\text{O}(l)$	1.77
$\text{Cr}_2\text{O}_7^{2-}(aq) + 14 \text{H}^+(aq) + 6 e^- \rightarrow 2 \text{Cr}^{3+}(aq) + 7 \text{H}_2\text{O}(l)$	1.33
$\text{O}_2(g) + 2 \text{H}^+(aq) + 2 e^- \rightarrow \text{H}_2\text{O}_2(aq)$	0.70

- (a) Use the information in the table to calculate the following.
- E° for the reaction between $\text{Cr}_2\text{O}_7^{2-}(aq)$ and $\text{H}_2\text{O}_2(aq)$ at 298 K
 - E° for the reaction between $\text{Co}^{2+}(aq)$ and $\text{H}_2\text{O}_2(aq)$ at 298 K
- (b) Based on the calculated values of E° , the student must choose the titrant for which the titration reaction is thermodynamically favorable at 298 K.
- Which titrant should the student choose? Explain your reasoning.
 - Calculate the value of ΔG° , in kJ/mol_{rxn} , for the reaction between the chosen titrant and $\text{H}_2\text{O}_2(aq)$.



STOP

END OF EXAM

AP[®] CHEMISTRY
2017 SCORING GUIDELINES

Question 7

A student wants to determine the concentration of H_2O_2 in a solution of $\text{H}_2\text{O}_2(aq)$. The student can use one of two titrants, either dichromate ion, $\text{Cr}_2\text{O}_7^{2-}(aq)$, or cobalt(II) ion, $\text{Co}^{2+}(aq)$. The balanced chemical equations for the two titration reactions are shown below.



The half-reactions and the E° values for the systems related to the titrations above are given in the following table.

Half-Reaction	E° (V) at 298 K
$\text{Co}^{3+}(aq) + e^- \rightarrow \text{Co}^{2+}(aq)$	1.84
$\text{H}_2\text{O}_2(aq) + 2 \text{H}^+(aq) + 2 e^- \rightarrow 2 \text{H}_2\text{O}(l)$	1.77
$\text{Cr}_2\text{O}_7^{2-}(aq) + 14 \text{H}^+(aq) + 6 e^- \rightarrow 2 \text{Cr}^{3+}(aq) + 7 \text{H}_2\text{O}(l)$	1.33
$\text{O}_2(g) + 2 \text{H}^+(aq) + 2 e^- \rightarrow \text{H}_2\text{O}_2(aq)$	0.70

(a) Use the information in the table to calculate the following.

(i) E° for the reaction between $\text{Cr}_2\text{O}_7^{2-}(aq)$ and $\text{H}_2\text{O}_2(aq)$ at 298 K

$E^\circ = 1.33 - 0.70 = 0.63 \text{ V}$	1 point is earned for correctly combining E° values.
--	---

(ii) E° for the reaction between $\text{Co}^{2+}(aq)$ and $\text{H}_2\text{O}_2(aq)$ at 298 K

$E^\circ = -1.84 + 1.77 = -0.07 \text{ V}$	1 point is earned for correctly combining E° values.
--	---

AP[®] CHEMISTRY
2017 SCORING GUIDELINES

Question 7 (continued)

- (b) Based on the calculated values of E° , the student must choose the titrant for which the titration reaction is thermodynamically favorable at 298 K.

(i) Which titrant should the student choose? Explain your reasoning.

<p>The student should use the dichromate ion for the titration because, for the reaction, the value of E° is positive, which means that the reaction is thermodynamically favorable.</p> <p>OR</p> <p>$\Delta G^\circ = -nFE^\circ$ and n, F, and E° are all positive numbers, therefore $\Delta G^\circ < 0$, which means that the reaction is thermodynamically favorable.</p>	<p>1 point is earned for choosing the correct titrant and for understanding that a positive E° or a negative ΔG° is required.</p>
---	--

(ii) Calculate the value of ΔG° , in $\text{kJ/mol}_{\text{rxn}}$, for the reaction between the chosen titrant and $\text{H}_2\text{O}_2(aq)$.

$\Delta G^\circ = -nFE^\circ = -6(96,485 \frac{\text{C}}{\text{mol}})(0.63 \frac{\text{J}}{\text{C}})(\frac{1 \text{ kJ}}{1000 \text{ J}}) = -360 \text{ kJ/mol}_{\text{rxn}}$	<p>1 point is earned for calculating the value of ΔG°.</p>
--	--