


**2007 AP<sup>®</sup> CALCULUS BC FREE-RESPONSE QUESTIONS**



3. The graphs of the polar curves  $r = 2$  and  $r = 3 + 2\cos \theta$  are shown in the figure above. The curves intersect when  $\theta = \frac{2\pi}{3}$  and  $\theta = \frac{4\pi}{3}$ .
- (a) Let  $R$  be the region that is inside the graph of  $r = 2$  and also inside the graph of  $r = 3 + 2\cos \theta$ , as shaded in the figure above. Find the area of  $R$ .
- (b) A particle moving with nonzero velocity along the polar curve given by  $r = 3 + 2\cos \theta$  has position  $(x(t), y(t))$  at time  $t$ , with  $\theta = 0$  when  $t = 0$ . This particle moves along the curve so that  $\frac{dr}{dt} = \frac{dr}{d\theta}$ . Find the value of  $\frac{dr}{dt}$  at  $\theta = \frac{\pi}{3}$  and interpret your answer in terms of the motion of the particle.
- (c) For the particle described in part (b),  $\frac{dy}{dt} = \frac{dy}{d\theta}$ . Find the value of  $\frac{dy}{dt}$  at  $\theta = \frac{\pi}{3}$  and interpret your answer in terms of the motion of the particle.
- 

**WRITE ALL WORK IN THE PINK EXAM BOOKLET.**

**END OF PART A OF SECTION II**

**2007 AP<sup>®</sup> CALCULUS BC FREE-RESPONSE QUESTIONS****CALCULUS BC  
SECTION II, Part B****Time—45 minutes****Number of problems—3****No calculator is allowed for these problems.**

- 
4. Let  $f$  be the function defined for  $x > 0$ , with  $f(e) = 2$  and  $f'$ , the first derivative of  $f$ , given by  $f'(x) = x^2 \ln x$ .
- (a) Write an equation for the line tangent to the graph of  $f$  at the point  $(e, 2)$ .
- (b) Is the graph of  $f$  concave up or concave down on the interval  $1 < x < 3$ ? Give a reason for your answer.
- (c) Use antidifferentiation to find  $f(x)$ .
- 

$t$ (minutes)	0	2	5	7	11	12
$r'(t)$ (feet per minute)	5.7	4.0	2.0	1.2	0.6	0.5

5. The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function  $r$  of time  $t$ , where  $t$  is measured in minutes. For  $0 < t < 12$ , the graph of  $r$  is concave down. The table above gives selected values of the rate of change,  $r'(t)$ , of the radius of the balloon over the time interval  $0 \leq t \leq 12$ . The radius of the balloon is 30 feet when  $t = 5$ . (Note: The volume of a sphere of radius  $r$  is given by  $V = \frac{4}{3}\pi r^3$ .)
- (a) Estimate the radius of the balloon when  $t = 5.4$  using the tangent line approximation at  $t = 5$ . Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when  $t = 5$ . Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate  $\int_0^{12} r'(t) dt$ . Using correct units, explain the meaning of  $\int_0^{12} r'(t) dt$  in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than  $\int_0^{12} r'(t) dt$ ? Give a reason for your answer.
- 

**WRITE ALL WORK IN THE PINK EXAM BOOKLET.**