

2. To investigate how increases in environmental temperatures affect the metabolism of certain organisms, researchers incubated liver cells from toads at different temperatures and measured two markers of metabolic activity (Table 1): the rate of oxygen consumption and the rate of ATP synthesis.

(a) **Describe** the role of water in the hydrolysis of ATP.

TABLE 1. RATE OF OXYGEN CONSUMPTION AND ATP SYNTHESIS AT DIFFERENT TEMPERATURES

Metabolic Marker	20°C	25°C	30°C
Rate of Oxygen Consumption (nmol / min / mg of mitochondrial protein $\pm 2 SE_{\bar{x}}$)	12.8 ± 2.2	16.5 ± 2.0	22.1 ± 0.7
Rate of ATP Synthesis (nmol / min / mg of mitochondrial protein $\pm 2 SE_{\bar{x}}$)	12.6 ± 1.6	16.8 ± 2.0	21.07 ± 0.8

(b)

- Using the template in the space provided for your response, **construct** a bar graph that represents the data shown in Table 1. Your graph should be appropriately plotted and labeled.
- Based on the data provided, **determine** the temperature in °C at which the rate of oxygen consumption is different from the rate of oxygen consumption at 25°C.

(c)

- Based on the data in Table 1, **describe** the effect of temperature on the rate of ATP synthesis in liver cells from toads.
- Based on the data in Table 1, **calculate** the average amount of oxygen consumed, in nmol, for 10 mg of mitochondrial protein after 10 minutes at 25°C.

(d)

- Oligomycin is a compound that can block the channel protein function of ATP synthase. **Predict** the effects of using oligomycin on the proton gradient across the inner mitochondrial membrane.
- Justify** your prediction.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

If there are multiple parts to this question, write the part letter with your response.

Question 2: Interpreting and Evaluating Experimental Results
with Graphing

9 points

To investigate how increases in environmental temperatures affect the metabolism of certain organisms, researchers incubated liver cells from toads at different temperatures and measured two markers of metabolic activity (Table 1): the rate of oxygen consumption and the rate of ATP synthesis.

TABLE 1. RATE OF OXYGEN CONSUMPTION AND ATP SYNTHESIS AT DIFFERENT TEMPERATURES

Metabolic Marker	20°C	25°C	30°C
Rate of Oxygen Consumption (nmol/min/mg of mitochondrial protein $\pm 2SE_{\bar{x}}$)	12.8 \pm 2.2	16.5 \pm 2.0	22.1 \pm 0.7
Rate of ATP Synthesis (nmol/min/mg of mitochondrial protein $\pm 2SE_{\bar{x}}$)	12.6 \pm 1.6	16.8 \pm 2.0	21.07 \pm 0.8

- (a)


Describe the role of water in the hydrolysis of ATP.
Accept one of the following:
 - Water is added in the process of cleaving/splitting (a phosphate from) ATP.
 - Water breaks down/splits ATP.

1 point
- (b)

Using the template in the space provided for your response, **construct** a bar graph that represents the data shown in Table 1. Your graph should be appropriately plotted and labeled.

1 point

Sample Responses:



- Data are represented in a bar graph.
- Using the template in the space provided for your response, **construct** a bar graph that represents the data shown in Table 1. Your graph should be appropriately plotted and labeled.

1 point
- Graph is appropriately labeled.

	Using the template in the space provided for your response, construct a <u>bar</u> graph that represents the data shown in <u>Table 1</u> . Your graph should be appropriately <u>plotted</u> and <u>labeled</u> .	1 point
	<ul style="list-style-type: none"> Data points and error bars are correctly plotted. 	
	Based on the data provided, determine the temperature in °C at which the rate of oxygen consumption is different from the rate of oxygen consumption at 25°C.	1 point
	<ul style="list-style-type: none"> 30 	
Total for part (b)		4 points
(c)	Based on the data in <u>Table 1</u> , describe the effect of temperature on the rate of ATP synthesis in liver cells from toads. Accept one of the following: <ul style="list-style-type: none"> As the temperature increases, the rate of ATP synthesis also increases. There is a positive relationship (between temperature and ATP synthesis). Temperature and ATP synthesis are directly correlated. 	1 point
	Based on the data in <u>Table 1</u> , calculate the average amount of oxygen consumed, in nmol, for 10 mg of mitochondrial protein after 10 minutes at 25°C. 1,650 [16.5 nmol/min/mg × 10 mg × 10 min]	1 point
Total for part (c)		2 points
(d)	Oligomycin is a compound that can block the channel protein function of ATP synthase. Predict the effects of using oligomycin on the proton gradient across the inner mitochondrial membrane. Accept one of the following: <ul style="list-style-type: none"> (The proton gradient) will <u>increase/become steeper</u> (and may eventually plateau). The difference in the <u>concentration of protons/pH</u> (across the inner mitochondrial membrane) will increase. There will be <u>an increase in the concentration of protons/a decrease in pH</u> in the intermembrane space relative to that found within the mitochondrial matrix. 	1 point
	Justify your prediction. Accept one of the following: <ul style="list-style-type: none"> (Without protons being able to flow back into the matrix through ATP synthase), more protons will accumulate <u>in the intermembrane space/between the two mitochondrial membranes</u>. (Without protons being able to flow back into the matrix through ATP synthase), there will be a lower pH <u>in the intermembrane space/between the two mitochondrial membranes</u>. Protons will not be able to flow across the membrane (through ATP synthase), but the electron transport chain will still pump protons into the intermembrane space. 	1 point
Total for part (d)		2 points
Total for question 2		9 points