

Begin your response to **QUESTION 1** on this page.


PHYSICS C: ELECTRICITY AND MAGNETISM

SECTION II

Time—45 minutes

3 Questions

Directions: Answer all three questions. The suggested time is about 15 minutes for answering each of the questions, which are worth 15 points each. The parts within a question may not have equal weight. Show all your work in this booklet in the spaces provided after each part.



Note: Figure not drawn to scale.


1. Students perform an experiment to determine the value of vacuum permittivity ϵ_0 . Sphere A is nonconducting with charge $+q$ and is attached to an insulating rod. Sphere B is nonconducting with charge $+Q$, and has mass M . Sphere B rests on an insulating platform of negligible mass that is attached to a vertical ideal spring with spring constant k_s . Sphere B and the spring are initially at rest.

Sphere A is then brought near Sphere B without touching. When the centers of the spheres are separated by a vertical distance H , the spring has been compressed a distance y , as shown in the figure. The students measure y for different values of H .

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (a) On the following dot that represents Sphere B in the figure on the previous page, draw and label the forces (not components) that are exerted on Sphere B. Each force must be represented by a distinct arrow starting on, and pointing away from, the dot.




- (b) Derive the relationship between y and H to show that $y = \frac{1}{4\pi\epsilon_0} \frac{Qq}{k_s H^2} + \frac{Mg}{k_s}$.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (c) The students plot collected data of y as a function of $\frac{1}{H^2}$, as shown in the graph.



- i. Draw the best-fit line for the data.
- ii. Using the best-fit line, calculate an experimental value for the vacuum permittivity ϵ_0 when $Q = q = 2.00 \times 10^{-6} \text{ C}$ and $k_s = 25 \text{ N/m}$.
- iii. Using the best-fit line, calculate an experimental value for the mass of Sphere B.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (d) The students modify the experiment by replacing nonconducting Sphere B with conducting Sphere C that has the same charge $+Q$ and mass M . Sphere A is brought near Sphere C without touching, compressing the spring. Sphere C comes to rest.


i. In the original experiment, when the centers of spheres A and B are a vertical distance H_1 apart, the spring is compressed a distance y_1 . In the modified experiment, when the centers of spheres A and C are a vertical distance H_1 apart, the spring is compressed a distance y_2 .

Is y_2 greater than, less than, or equal to y_1 ?

_____ $y_2 > y_1$ _____ $y_2 < y_1$ _____ $y_2 = y_1$


Justify your answer.

- ii. Sphere C is then grounded with a wire. On the following figure, draw an arrow indicating the direction that the platform will move immediately after being grounded. If the platform remains stationary, write “does not move.”



GO ON TO THE NEXT PAGE.

Begin your response to **QUESTION 2** on this page.



2. Two parallel conducting rails are separated by distance $d = 0.30$ m. A resistor of resistance $R = 0.20 \, \Omega$ connects the rails. A conducting bar is placed on a sloped section of the rails at height H above the horizontal section of the rails. Frictional forces and the resistances of the bar and rails are negligible.

- At time $t = 0$, the bar is released from rest from position x_0 and slides down the sloped section of the rails, as shown in the Perspective View.
- At time t_1 , the bar reaches position x_1 and smoothly transitions to the horizontal section of the rails and enters a uniform magnetic field of magnitude $B_1 = 0.40$ T that is directed in the $+y$ -direction.
- At time t_2 , the bar reaches position x_2 and enters a region with no magnetic field.
- At time t_3 , the bar reaches position x_3 and enters a uniform magnetic field of magnitude $B_2 = 0.60$ T that is directed in the $+z$ -direction.
- At time t_4 , the bar reaches position x_4 and enters a region with no magnetic field.

The bar is at position x_B (shown in Top View) at time t_B such that $t_1 < t_B < t_2$.

GO ON TO THE NEXT PAGE.