



- D. The chemist weighed out 0.100 g  $\text{P}_4\text{O}_{10}$  of and 100.0 g of  $\text{H}_2\text{O}$  to perform a second trial. In the second trial, some of the solid  $\text{P}_4\text{O}_{10}$  stuck to the weighing paper and was not transferred to the calorimeter. Given that  $\text{P}_4\text{O}_{10}$  is the limiting reactant, would  $\Delta T$  for the second trial be greater than, less than, or equal to the value in the first trial? Justify your answer.


$\text{P}_4(\text{s})$  also reacts readily with  $\text{Cl}_2(\text{g})$  to produce phosphorus trichloride,  $\text{PCl}_3(\text{g})$ , which in turn reacts with  $\text{Cl}_2(\text{g})$  in an equilibrium process to produce  $\text{PCl}_5(\text{g})$ . The reactions are represented by equations 3 and 4.




- E. Calculate the standard enthalpy of formation of  $\text{PCl}_5(\text{g})$  represented by equation 5.



The following particle-level diagram represents the contents of the vessel in an equilibrium mixture at 546 K involving equation 4.




- F. Equation 4 for the reaction that occurs is shown.



- If each particle in the diagram represents a partial pressure of 1.00 atm, what is the value of  $K_p$  for the equilibrium mixture at 546 K?
- Does the value of  $K_p$  increase, decrease, or remain the same when the temperature is increased to 596 K? Justify your answer based on  $\Delta H_2^\circ$ .

4. A scientist is investigating the properties of a mixture of  $\text{CH}_3\text{OH}$  and  $\text{H}_2\text{CO}$ . The scientist generates the following diagram to represent the mixture.



- A. Identify the hybridization of the valence orbitals of the C atom in the  $\text{H}_2\text{CO}$  molecule.
- B. In the diagram provided, draw a SINGLE dashed line (---) to represent a strong hydrogen-bonding attraction between one  $\text{CH}_3\text{OH}$  molecule and one  $\text{H}_2\text{CO}$  molecule in the mixture.
- C. The scientist plans to cool a gaseous mixture of  $\text{CH}_3\text{OH}$  and  $\text{H}_2\text{CO}$  to form a liquid mixture and finds data on the two compounds. The data are summarized in the table.

Substance	Melting Point (K)	Boiling Point (K)	Enthalpy of Vaporization (kJ/mol)
$\text{CH}_3\text{OH}$	176	338	37.6
$\text{H}_2\text{CO}$	181	254	24.2

- i. Propose a temperature to which the mixture should be cooled such that  $\text{CH}_3\text{OH}$  and  $\text{H}_2\text{CO}$  will both be liquids.
- ii. The scientist analyzes the mixture after it is cooled and determines that 8.59 g of  $\text{CH}_3\text{OH}(l)$  is present. Calculate the amount of thermal energy, in kJ, that was removed to condense the 8.59 g of  $\text{CH}_3\text{OH}$  (molar mass 32.04 g/mol) at its boiling point.

## Question 4: Short Answer

4 points

A For the correct answer:


Point 01

 $sp^2$ 

B For a correct diagram:

Point 02

The diagram should show a dashed line between the O atom in one  $\text{H}_2\text{CO}$  molecule and the H atom in the  $-\text{OH}$  group of one  $\text{CH}_3\text{OH}$  molecule. See example response below.



C (i) For a correct proposal:

Point 03

The proposed temperature should be in the range 181 K–254 K.

(ii) For the correct calculated value:

Point 04

$$8.59 \text{ g CH}_3\text{OH} \times \frac{1 \text{ mol CH}_3\text{OH}}{32.04 \text{ g CH}_3\text{OH}} \times \frac{-37.6 \text{ kJ}}{1 \text{ mol CH}_3\text{OH}} = -10.1 \text{ kJ}, \text{ so } 10.1 \text{ kJ are removed.}$$