	Phase 1 Error generation	Phase 2 Precision weighting	Phase 3 Accumulation	Phase 4 Ignition decision	Phase 5 Global broadcast
Approx. time after onset	**0–50 ms**	**50–120 ms**	**120–250 ms**	**250–320 ms**	**320–500 ms**
Neural substrate	Feedforward sensory input; prediction error in early cortex	Neuromodulatory gating of precision (interoceptive and exteroceptive)	Recurrent loops in association cortex	Frontoparietal–thalamic loop compares S_t with θ_t	Cortico–thalamo–cortical broadcast (global workspace)
APGI variable	Prediction errors ε^e, ε^i	Precision Π^e, Π^i_{eff} (gain control)	Accumulated surprise S_t (rises over time)	Ignition if $S_t > \theta_t$	Ignition state $B_t = 1$ (workspace stabilisation)
EEG/MEG marker	C1/P1 and early gamma ~50–100 ms	Pre-stimulus alpha power (attentional gain)	N2 / sustained gamma ~150–250 ms	P3b onset ~250 ms	P3b (250–500 ms; to ~600 ms when θ_t is high)

Latencies are approximate and stimulus-dependent; phases are temporal, not the pipeline stages of Figure 1.

APGI 1.1 / P1F4 / Hypothesized timing schematic
Phase boundaries and substrate assignments are hypotheses; latencies depend on stimulus and state.