

PBIO-141

Sensory and Physiological Ecology of Plants

6: Responses to shading

Pedro J. Aphalo

January–February 2022

M.Sc. in Plant Biology, University of Helsinki

<http://blogs.helsinki.fi/aphalo/>

©2006-2022 by Pedro J. Aphalo

University of Helsinki, Finland.

<http://blogs.helsinki.fi/senpep-blog/>

Sensory and Physiological Ecology of Plants slides by Pedro J. Aphalo are licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.



Typeset in Lucida Sans, Luicda Bright, Lucida Console and Lucida Math. Icons from fonts "WebHostingHub Glyphs" (under SIL-Open Font License) from

<https://www.webhostinghub.com/>; "insect icons" (free from

<http://www.woodcutter.es/>); "linea-basic-10" and "linea-weather-10" (free from <https://github.com/linea-io>), "Mini Pics Uprooted Twig" and "Mini Pics Uprooted Twig" (commercial, from Image Club Graphics, Inc.). Plant icon as .svg by Abdul Wahhab (free from NounProject.com).

Illustrations and text quoted from copyrighted sources is excluded from this license and their use should respect the original licenses.

Outline

A new paradigm

Responses to (impending) shading

A new paradigm

A new paradigm is emerging

An exciting time to study plants!!

- Currently it is a very exciting time to study plants...
- ...especially how their molecular biology, evolution and ecology are linked.
- We are in the middle of a paradigm shift...
- ...for a long time we have disregarded/underestimated the central and complex role of plants' "senses".
- However, we should be open minded and use our imagination to understand plant life as it is...
- ...rather than to project 'human' features and qualities onto plants.

Responses to (impending) shading

Shade avoidance vs. tolerance strategies: ≠ “goals”

These are two extreme strategies with putative roles:

- Shade avoidance → compete for access to strong light
- Shade tolerance → efficiently use weak light
- Acclimated phenotypes are very different!
- Sources of information used in some combination are many:
- Cues: R:FR photon ratio, B:G photon ratio, blue light irradiance, UV irradiance, wind, touch, ethylene (signal?), including their temporal dynamics.

Shade avoidance vs. tolerance 10 min

- How do you expect avoidance and tolerance to work in practice?
- Can you think about clear-cut and less clear-cut cases?
- In which environments tolerance is likely to be better for fitness than shade avoidance?

Leaf size and angle, petiole length, chlorophyll

End-of-day R or FR light (10 min/day), *Petunia axillaris*, (Casal et al. 1987)






Figure 2. Effect of end-of-day R (○, □ or ---) or FR (●, ▨ or ...) irradiations on leaf chlorophyll content of the largest leaf (a), area of the largest leaf (b), leaf area per plant (c), total biomass per plant (d), and net assimilation rate (NAR) (e). Measurements were taken on different plants at each harvest except for Day 0 in (a). Data from Experiment A. Bars indicate \pm SEM.

Stem elongation

- In sparse canopies ‘horizontal’ light flux is important for anticipated sensing of neighbours.
- Growing internodes are a key site of perception.



Stem elongation



Irradiation with red or far-red light from LEDs at the growing internode (whole photoperiod), *Betula pendula*. (from Aphalo and Rikala 2006).

Leaf size, thickness and photosynthesis

High (HL) and low (LL) PAR, *Fuchsia magellanica* (Aphalo et al. 1991).





Figure 4. Responses of CO₂ flux per unit projected leaf area (left) and CO₂ flux per unit leaf dry weight (right), to quantum flux density in plants grown under white light. HL = 350 μmol m⁻² s⁻¹ and LL = 25 μmol m⁻² s⁻¹, equal to 12.6 and 0.9 mol m⁻² d⁻¹. Measured between 134 and 150 d after the start of treatment, each replicate is from a different plant; data shown with the same symbols were measured on consecutive days.

Stems position and length, apical dominance

Sunlight, full and minus blue, *Medicago truncatula* Rai, Yan and Aphalo, unpublished.



Preemptive acclimation: shade




Flow of information in preemptive acclimation to shade by perception of radiation changes. Arrows represent flows of information: **blue** = retrieved from genome (stored during earlier generations), **black** = acquired and/or 'memorized' during an individual's or its progenitor's lifetime, **teal** = regulation of gene expression by phenome or downward causation, **red** = lagged correlation between early changes in spectral irradiance and future low PAR irradiance, **orange** = outcome of information processing: a 'decision', based on an 'implicit forecast of impending shade', leading to developmental adjustments that would increase the probability of higher fitness in the presence of neighbours in comparison with phenotypes lacking preemptive acclimation. **green** = 'Shading mitigated' compared, in probabilistic terms, to no acclimation. Dashed boxes and arrows represent the likely or forecasted future.

Preemptive acclimation: What is the evidence?

- Several plant responses can be only explained from the evolutive/fitness point of view as being a ‘preparation’ to tolerate or escape future stress events or take advantage of future favourable conditions.
- Preemptive shade avoidance as a response to reflected far-red light from neighbouring plants.
- Winter hardening and dehardening, timing of bud burst, timing of flowering, etc.
- *Anticipatory responses* and *future perception* are terms also used in this context.


Timing of shade

Arabidopsis seedlings (Sellaro2012).



Sunflecks, UVR8, CRY1

Arabidopsis seedlings (from Moriconi et al. 2018). In A $\text{mmol m}^{-2} \text{s}^{-1}$ should read $\mu\text{mol m}^{-2} \text{s}^{-1}$.



Acclimation vs. rapid/reversible responses



5 min

(from Casal 2013). What about the balance between light reactions and carbon reactions of photosynthesis?





Figure 3

Plant responses to either sunlight (*left*) or shade light (*right*) perceived by photosensory receptors.

Role of photoreceptors in responses to shade

(from Casal 2013).



References

-  Aphalo, P. J., D. Gibson and A. H. Di Benedetto (1991). "Responses of growth, photosynthesis, and leaf conductance to white light irradiance and end-of-day red and far-red pulses in *Fuchsia magellanica* Lam". In: *New Phytologist* 117, pp. 461–471.
-  Aphalo, P. J. and R. Rikala (2006). "Spacing of silver birch seedlings grown in containers of equal size affects their morphology and its variability.". In: *Tree physiology* 26.9, pp. 1227–1237. DOI: [10.1093/treephys/26.9.1227](https://doi.org/10.1093/treephys/26.9.1227).
-  Aphalo, P. J. and V. O. Sadras (Dec. 2021). "Explaining preemptive acclimation by linking information to plant phenotype". In: *Journal of Experimental Botany*. DOI: [10.1093/jxb/erab537](https://doi.org/10.1093/jxb/erab537).

References ii

-  Casal, J. J., P. J. Aphalo and R. A. Sánchez (1987). "Phytochrome effects on leaf growth and chlorophyll content in Petunia axillaris". In: *Plant, Cell and Environment* 10.6, pp. 509–514. DOI: [10.1111/j.1365-3040.1987.tb01829.x](https://doi.org/10.1111/j.1365-3040.1987.tb01829.x).
-  Casal, J. J. (Apr. 2013). "Photoreceptor Signaling Networks in Plant Responses to Shade". In: *Annual Review of Plant Biology* 64.1, pp. 403–427. DOI: [10.1146/annurev-arplant-050312-120221](https://doi.org/10.1146/annurev-arplant-050312-120221).
-  Moriconi, V., M. Binkert, M. C. C. Rojas, R. Sellaro, R. Ulm and J. J. Casal (2018). "Perception of sunflecks by the UV-B photoreceptor UV RESISTANCE LOCUS 8". In: *Plant physiology* 177, pp. 75–81. DOI: [10.1104/pp.18.00048](https://doi.org/10.1104/pp.18.00048).



Can plants ‘predict’ the arrival of the cold season?
What sources of information do they use?



Copyright P. J. Aphalo



Can plants ‘predict’ the arrival of the spring?
What sources of information do they use?





Can plants emit ‘signals’ for their benefit?
What are the signals used to attract pollinators?





Can plants emit ‘signals’ for their benefit?

What are the signals used to enlist “helpers” for seed dispersal?





Can plants emit signals to deceive other organisms?
How does mimicry work?





Can plants ‘generate’ environmental correlations for their benefit?

How does release of organic volatile compounds create a correlation?





Do symbioses involve communication?

