

R for Photobiology

A handbook

Pedro J. Aphalo

DRAFT

Contents

Contents	i
List of abbreviations and symbols	iii
I Getting ready	1
1 Introduction	3
1.1 Radiation and molecules	3
2 Optics	5
2.1 Task:	5
3 Photochemistry	7
3.1 Task:	7
4 Software	9
4.1 Task:	9
5 Photobiology R packages	11
5.1 Task:	11
II Cookbook	13
6 Unweighted irradiance	15
6.1 Task: (energy) irradiance from spectral irradiance	15
6.2 Task: photon irradiance from spectral irradiance	16
6.3 Task: irradiance from spectral photon irradiance	17
6.4 Task: irradiances for more than one waveband	18
6.5 Task: use simple wavebands	18
6.6 Task: define simple wavebands	20
6.7 Task: photon ratios	21
6.8 Task: energy ratios	22
6.9 Task: calculate average number of photons per unit energy . . .	22
6.10 Task: split energy irradiance into regions	22
6.11 Task: split photon irradiance into regions	23
7 Weighted and effective irradiance	25

7.1 Task:	25
8 Colour	27
8.1 Task:	27
9 Photoreceptors	29
9.1 Task:	29
10 Radiation sources	31
10.1 Task:	31
11 Filters	33
11.1 Task:	33
12 Plotting spectra	35
12.1 Task:	35
13 Calibration	37
13.1 Task:	37
14 Simulation	39
14.1 Task:	39
15 Measurement	41
15.1 Task:	41
16 Optimizing performance	43
16.1 Task:	43

List of abbreviations and symbols

For quantities and units used in photobiology we follow, as much as possible, the recommendations of the Commission Internationale de l'Éclairage as described by Sliney2007.

Symbol	Definition
α	absorptance (%).
Δe	water vapour pressure difference (Pa).
ϵ	emittance (W m^{-2}).
λ	wavelength (nm).
θ	solar zenith angle (degrees).
ν	frequency (Hz or s^{-1}).
ρ	reflectance (%).
σ	Stefan-Boltzmann constant.
τ	transmittance (%).
χ	water vapour content in the air (g m^{-3}).
A	absorbance (absorbance units).
ANCOVA	analysis of covariance.
ANOVA	analysis of variance.
BSWF	BSWF.
c	speed of light in a vacuum.
CCD	charge coupled device, a type of light detector.
CDOM	coloured dissolved organic matter.
CFC	chlorofluorocarbons.
c.i.	confidence interval.
CIE	Commission Internationale de l'Éclairage; or erythema action spectrum standardized by CIE.
CTC	closed-top chamber.
DAD	diode array detector, linear light detector based on photodiodes.
DBP	dibutylphthalate.
DC	direct current.
DIBP	diisobutylphthalate.
DNA(N)	UV action spectrum for 'naked' DNA.
DNA(P)	UV action spectrum for DNA in plants.
DOM	dissolved organic matter.
DU	Dobson units.
e	water vapour partial pressure (Pa).
E	(energy) irradiance (W m^{-2}).
$E(\lambda)$	spectral (energy) irradiance ($\text{W m}^{-2} \text{ nm}^{-1}$).

E_0	fluence rate, also called scalar irradiance (W m^{-2}).
ESR	early stage researcher.
FACE	free air carbon-dioxide enhancement.
FEL	a certain type of 1000 W incandescent lamp.
FLAV	UV action spectrum for accumulation of flavonoids.
FWHM	full-width half-maximum.
GAW	Global Atmosphere Watch.
GEN	generalized plant action spectrum, also abbreviated as GPAS Caldwell1971.
GEN(G)	mathematical formulation of GEN by Green1974 .
GEN(T)	mathematical formulation of GEN by Thimijan1978.
h	Planck's constant.
h'	Planck's constant per mole of photons.
H	exposure, frequently called dose by biologists ($\text{kJ m}^{-2} \text{d}^{-1}$).
H^{BE}	biologically effective (energy) exposure ($\text{kJ m}^{-2} \text{d}^{-1}$).
H_p^{BE}	biologically effective photon exposure ($\text{mol m}^{-2} \text{d}^{-1}$).
HPS	high pressure sodium, a type of discharge lamp.
HSD	honestly significant difference.
k_B	Boltzmann constant.
L	radiance ($\text{W sr}^{-1} \text{m}^{-2}$).
LAI	leaf area index, the ratio of projected leaf area to the ground area.
LED	light emitting diode.
LME	linear mixed effects (type of statistical model).
LSD	least significant difference.
n	number of replicates (number of experimental units per treatment).
N	total number of experimental units in an experiment.
N_A	Avogadro constant (also called Avogadro's number).
NIST	National Institute of Standards and Technology (U.S.A.).
NLME	non-linear mixed effects (statistical model).
OTC	open-top chamber.
PAR	PAR, 400–700 nm. measured as energy or photon irradiance.
PC	polycarbonate, a plastic.
PG	UV action spectrum for plant growth.
PHIN	UV action spectrum for photoinhibition of isolated chloroplasts.
PID	PID (control algorithm).
PMMA	polymethylmethacrylate.
PPFD	PPFD, another name for PAR photon irradiance (Q_{PAR}).
PTFE	polytetrafluoroethylene.
PVC	polyvinylchloride.
q	energy in one photon ('energy of light').
q'	energy in one mole of photons.
Q	photon irradiance ($\text{mol m}^{-2} \text{s}^{-1}$ or $\mu\text{mol m}^{-2} \text{s}^{-1}$).
$Q(\lambda)$	spectral photon irradiance ($\text{mol m}^{-2} \text{s}^{-1} \text{nm}^{-1}$ or $\mu\text{mol m}^{-2} \text{s}^{-1} \text{nm}^{-1}$).
r_0	distance from sun to earth.
RAF	RAF (nondimensional).
RH	relative humidity (%).
s	energy effectiveness (relative units).

$s(\lambda)$	spectral energy effectiveness (relative units).
s^p	quantum effectiveness (relative units).
$s^p(\lambda)$	spectral quantum effectiveness (relative units).
s.d.	standard deviation.
SDK	software development kit.
s.e.	standard error of the mean.
SR	spectroradiometer.
t	time.
T	temperature.
TUV	tropospheric UV.
U	electric potential difference or voltage (e.g. sensor output in V).
UV	ultraviolet radiation ($\lambda = 100\text{--}400\text{ nm}$).
UV-A	ultraviolet-A radiation ($\lambda = 315\text{--}400\text{ nm}$).
UV-B	ultraviolet-B radiation ($\lambda = 280\text{--}315\text{ nm}$).
UV-C	ultraviolet-C radiation ($\lambda = 100\text{--}280\text{ nm}$).
UV ^{BE}	biologically effective UV radiation.
UTC	coordinated universal time, replaces GMT in technical use.
VIS	radiation visible to the human eye ($\approx 400\text{--}700\text{ nm}$).
WMO	World Meteorological Organization.
VPD	water vapour pressure deficit (Pa).
WOUDC	World Ozone and Ultraviolet Radiation Data Centre.

Part I

Getting ready

CHAPTER 1

Introduction

Abstract

In this chapter we explain the physical basis of optics and photo-chemisatry.

1.1 Radiation and molecules

CHAPTER 2

Optics

Abstract

In this chapter we explain how to .

2.1 Task:

CHAPTER 3

Photochemistry

Abstract

In this chapter we explain how to .

3.1 Task:

CHAPTER 4

Software

Abstract

In this chapter we explain how to .

4.1 Task:

CHAPTER 5

Photobiology R packages

Abstract

In this chapter we explain how to .

5.1 Task:

Part II

Cookbook

CHAPTER 6

Unweighted irradiance

Abstract

In this chapter we explain how to calculate unweighted energy and photon irradiances from spectral irradiance.

6.1 Task: (energy) irradiance from spectral irradiance

The task to be completed is to calculate the (energy) irradiance (E) in W m^{-2} from spectral (energy) irradiance ($Q(\lambda)$) in $\text{W m}^{-2} \text{ nm}^{-1}$ and the corresponding wavelengths (λ) in nm.

$$Q_{\lambda_1 < \lambda < \lambda_2} = \int_{\lambda_1}^{\lambda_2} E(\lambda) \, d\lambda \quad (6.1)$$

Let's assume that we want to calculate photosynthetically active radiation (PAR) energy irradiance, for which the most accepted limits are $\lambda_1 = 400\text{nm}$ and $\lambda_2 = 700\text{nm}$. In this example we will use example data for sunlight to calculate $E_{400\text{nm} < \lambda < 700\text{nm}}$:

```
with(sun.data, energy_irradiance(w.length, s.e.irrad,
                                new_waveband(400, 700)))

## range.400.700
##          196.7
```

Function `PAR()` is predefined in package `photobiologyVIS` as a convenience function, so the code above can be replaced by:

```
with(sun.data, energy_irradiance(w.length, s.e.irrad, PAR()))

##      PAR
## 196.7
```

If no waveband is supplied as argument, then the whole range of wavelengths in the spectral data is used for the integration, and the 'name' attribute is generated accordingly:

```
with(sun.data, energy_irradiance(w.length, s.e.irrad))

## range.293.800
##          268.9
```

If a waveband that does not fully overlap with the data is supplied as argument, then spectral irradiance for wavelengths outside the range is assumed to be zero:

```
with(sun.data, energy_irradiance(w.length, s.e.irrad,
                                new_waveband(700,1000)))

## range.700.1000
##          44.1
```

If a waveband that does not overlap with the data is supplied as argument, then spectral irradiance for wavelengths outside the range is assumed to be zero:

```
with(sun.data, energy_irradiance(w.length, s.e.irrad,
                                new_waveband(100,200)))

## range.100.200
##          0
```

6.2 Task: photon irradiance from spectral irradiance

The task to be completed is to calculate the photon irradiance (Q) in $\text{mol m}^{-2} \text{s}^{-1}$ from spectral (energy) irradiance ($E(\lambda)$) in $\text{W m}^{-2} \text{nm}^{-1}$ and the corresponding wavelengths (λ) in nm.

The energy of a quantum of radiation in a vacuum, q , depends on the wavelength, λ , or frequency¹, ν ,

$$q = h \cdot \nu = h \cdot \frac{c}{\lambda} \quad (6.2)$$

with the Planck constant $h = 6.626 \times 10^{-34} \text{ Js}$ and speed of light in vacuum $c = 2.998 \times 10^8 \text{ m s}^{-1}$. When dealing with numbers of photons, the equation (6.2) can be extended by using Avogadro's number $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$. Thus, the energy of one mole of photons, q' , is

$$q' = h' \cdot \nu = h' \cdot \frac{c}{\lambda} \quad (6.3)$$

with $h' = h \cdot N_A = 3.990 \times 10^{-10} \text{ Js mol}^{-1}$. Example 1: red light at 600 nm has about 200 kJ mol^{-1} , therefore, 1 μmol photons has 0.2 J. Example 2: UV-B

¹Wavelength and frequency are related to each other by the speed of light, according to $\nu = c/\lambda$ where c is speed of light in vacuum. Consequently there are two equivalent formulations for equation 6.2.

radiation at 300 nm has about 400 kJ mol⁻¹, therefore, 1 μmol photons has 0.4 J. Equations 6.2 and 6.3 are valid for all kinds of electromagnetic waves.

Combining equations 6.1 and 6.3 we obtain:

$$Q_{\lambda_1 < \lambda < \lambda_2} = \int_{\lambda_1}^{\lambda_2} E(\lambda) \frac{h' \cdot c}{\lambda} d\lambda \quad (6.4)$$

Let's assume that we want to calculate photosynthetically active radiation (PAR) photon irradiance. In this example we will use example data for sunlight.

```
with(sun.data,
      photon_irradiance(w.length, s.e.irrad, PAR()))

##          PAR
## 0.0008938
```

If we want to have Q_{PAR} (PPFD) expressed in the usual units of μmol m⁻² s⁻¹, we need to multiply the result above by 10⁶:

```
with(sun.data,
      photon_irradiance(w.length, s.e.irrad, PAR())) * 1e6

##          PAR
## 893.8
```

PAR() is predefined in package photobiologyVIS as a convenience function, see section 6.1 for an example with arbitrary values for λ₁ and λ₂.

6.3 Task: calculate energy and photon irradiances from spectral photon irradiance

In the case of the calculation of energy irradiance from spectral photon irradiance the calculation is:

$$I_{\lambda_1 < \lambda < \lambda_2} = \int_{\lambda_1}^{\lambda_2} Q_{\lambda} \frac{\lambda}{h' \cdot c} d\lambda \quad (6.5)$$

And the code²:

```
with(sun.data, energy_irradiance(w.length, s.q.irrad, PAR()),
      unit.in = "photon")

##          PAR
## 0.0008938
```

The calculation of photon irradiance from spectral photon irradiance, is a simple integration, analogous to that in equation 6.1, and the code is:

```
with(sun.data, photon_irradiance(w.length, s.q.irrad, PAR()),
      unit.in = "photon")

##          PAR
## 4.158e-09
```

²The dataframe sun.data contains both spectral energy irradiance vales in 'column' s.e.irrad and spectral photon irradiance in 'column' s.q.irrad

6.4 Task: irradiances for more than one waveband

It is possible to calculate the irradiances for several wavebands with a single function call by supplying a list of wavebands as argument:

```
with(sun.data, photon_irradiance(w.length, s.e.irrad, list(Red(),  
  Green(), Blue())) * 1e+06  
  
##      Red.ISO Green.ISO  Blue.ISO  
##      452.2      220.2      149.0  
  
Q.RGB <- with(sun.data, photon_irradiance(w.length, s.e.irrad,  
  list(Red(), Green(), Blue())) * 1e+06  
signif(Q.RGB, 3)  
  
##      Red.ISO Green.ISO  Blue.ISO  
##      452      220      149  
  
Q.RGB[1]  
  
## Red.ISO  
## 452.2  
  
Q.RGB["Green.ISO"]  
  
## Green.ISO  
## 220.2
```

A named list can be used to override the use as names for the output of the waveband names:

```
with(sun.data, photon_irradiance(w.length, s.e.irrad, list(R = Red(),  
  G = Green(), B = Blue())) * 1e+06  
  
##      R      G      B  
## 452.2 220.2 149.0
```

Even when using a single waveband:

```
with(sun.data,  
  photon_irradiance(w.length, s.e.irrad,  
    list(UVB=UVB())) * 1e6  
  
##      UVB  
## 1.527
```

6.5 Task: use simple wavebands

Please, consult the packages' documentation for a list of predefined functions for creating wavebands. Here we will present just a few examples of their use. We usually associate wavebands with colours, however, in many cases there are different definitions in use. For this reason, the functions provided accept an argument that can be used to select the definition to use. In general, the default, is to use the ISO standard whenever it is applicable. The case of the various definitions in use for the UV-B waveband are described on page 19

We can use a predefined function to create a new waveband object, which as any other R object can be assigned to a variable:

```
uvb <- UVB()
uvb

## UVB.ISO
## low (nm) 280
## high (nm) 315
```

As seen above, there is a specialized print function for wavebands. Functions available are min, max, range, center_wl, labels, and color. ■

```
red <- Red()
red

## Red.ISO
## low (nm) 610
## high (nm) 760

min(red)

## [1] 610

max(red)

## [1] 760

range(red)

## [1] 610 760

midpoint(red)

## [1] 685

labels(red)

## $label
## [1] "Red.ISO"

color(red)

## Red.ISO CMF Red.ISO CC
## "#900000" "#FF0000"
```

Here we demonstrate the use of an argument to choose a certain definition:

```
UVB()

## UVB.ISO
## low (nm) 280
## high (nm) 315

UVB("ISO")

## UVB.ISO
## low (nm) 280
## high (nm) 315
```

```

UVB("CIE")

## UVB.CIE
## low (nm) 280
## high (nm) 315

UVB("medical")

## UVB.medical
## low (nm) 290
## high (nm) 320

UVB("none")

## UVB.none
## low (nm) 280
## high (nm) 320

```

Here we demonstrate the importance of complying with standards, and how much the photon irradiance calculated can depend on the definition used.

```

with(sun.data,
      photon_irradiance(w.length, s.e.irrad, UVB("ISO"))) * 1e6

## UVB.ISO
##      1.527

with(sun.data,
      photon_irradiance(w.length, s.e.irrad, UVB("none"))) * 1e6

## UVB.none
##      3.282

```

6.6 Task: define simple wavebands

Here we briefly introduce `new_waveband`, and only in chapter ?? we describe its use in full detail, including the use of spectral weighting functions (SWFs).

Defining a new waveband based on extreme wavelengths expressed in nm.

```

wb1 <- new_waveband(500,600)
wb1

## range.500.600
## low (nm) 500
## high (nm) 600

with(sun.data,
      photon_irradiance(w.length, s.e.irrad, wb1)) * 1e6

## range.500.600
##      314.1

wb2 <- new_waveband(500,600, wb.name="my.colour")
wb2

```

```
## my.colour
## low (nm) 500
## high (nm) 600

with(sun.data,
      photon_irradiance(w.length, s.e.irrad, wb2)) * 1e6

## my.colour
##      314.1
```

6.7 Task: photon ratios

In photobiology sometimes we are interested in calculation the photon ratio between two wavebands. It makes more sense to calculate such ratios if both numerator and denominator wavebands have the same ‘width’ or if the numerator waveband is fully nested in the denominator waveband. However, frequently used ratios like the UV-B to PAR photon ratio do not comply with this. For this reason, our functions do not enforce any such restrictions.

For example a ratio frequently used in plant photobiology is the read to far-red photon ratio (R:FR photon ratio or ζ). If we follow the wavelength ranges in the definition given by **Morgan1981a** using photon irradiance³:

$$\zeta = \frac{Q_{655\text{nm} < \lambda < 665\text{nm}}}{Q_{725\text{nm} < \lambda < 735\text{nm}}} \quad (6.6)$$

To calculate this for our example sunlight spectrum we can use the following code:

```
with(sun.data,
      photon_ratio(w.length, s.e.irrad, Red("Smith"), Far_red("Smith")))

## [1] 1.251
```

or using the predefined convenience function `R_FR_ratio`:

```
with(sun.data,
      R_FR_ratio(w.length, s.e.irrad))

## [1] 1.251
```

Using defaults for waveband definitions:

```
with(sun.data,
      energy_ratio(w.length, s.e.irrad, UVB(), PAR()))

## [1] 0.00299
```

³In the original text photon fluence rate is used but it not clear whether photon irradiance was meant instead.

6.8 Task: energy ratios

An energy ratio, equivalent to ζ can be calculated as follows:

```
with(sun.data,
      energy_ratio(w.length, s.e.irrad, Red("Smith"), Far_red("Smith")))■
## [1] 1.384
```

For this infrequently used ratio, no pre-defined function is provided.

6.9 Task: calculate average number of photons per unit energy

When comparing photo-chemical and photo-biological responses under different light sources it is of interest to calculate the photons per energy in mol W^{-1} . In this case only one waveband definition is used to calculate the quotient:

$$\tilde{q}' = \frac{Q_{\lambda_1 < \lambda < \lambda_2}}{E_{\lambda_1 < \lambda < \lambda_2}} \quad (6.7)$$

```
with(sun.data,
      photons_energy_ratio(w.length, s.e.irrad, PAR()))
## [1] 4.544e-06
```

For obtaining the same quotient in $\mu\text{mol W}^{-1}$ we just need to multiply by 10^6 . We can use such a multiplier to convert E [W m^{-2}] into Q [$\mu\text{mol m}^{-2} \text{s}^{-1}$], or as a divisor to convert Q [$\mu\text{mol m}^{-2} \text{s}^{-1}$] into E [W m^{-2}], *for a given light source and waveband*:

```
with(sun.data, photons_energy_ratio(w.length, s.e.irrad, PAR())) *■
1e+06
## [1] 4.544
```

6.10 Task: calculate the contribution of different regions of a spectrum to energy irradiance

It can be of interest to split the total (energy) irradiance into adjacent regions delimited by arbitrary wavelengths. We can use the function `split_energy_irradiance` to obtain to energy of each of the regions delimited by the values in nm supplied in a numeric vector:

```
with(sun.data, split_energy_irradiance(w.length, s.e.irrad, c(400,■
500, 600, 700)))
## range.400.500 range.500.600 range.600.700
##          69.63          68.53          58.54
```

Here we demonstrate that the sum of the four ‘split’ irradiances add to the total for the range of wavelengths covered:

```
with(sun.data, sum(split_energy_irradiance(w.length, s.e.irrad,
c(400, 500, 600, 700))))

## [1] 196.7

with(sun.data, energy_irradiance(w.length, s.e.irrad, PAR()))

## PAR
## 196.7
```

It also possible to obtain the ‘split’ as a vector of fractions adding up to one,

```
with(sun.data, split_energy_irradiance(w.length, s.e.irrad, c(400,
500, 600, 700), scale = "relative"))

## range.400.500 range.500.600 range.600.700
## 0.3540 0.3484 0.2976
```

or as percentages:

```
with(sun.data, split_energy_irradiance(w.length, s.e.irrad, c(400,
500, 600, 700), scale = "percent"))

## range.400.500 range.500.600 range.600.700
## 35.40 34.84 29.76
```

A vector of two wavelengths is valid input, although not very useful for percentages:

```
with(sun.data, split_energy_irradiance(w.length, s.e.irrad, c(400,
700), scale = "percent"))

## range.400.700
## 100
```

Although for `scale="absolute"`, the default, it can be used as a quick way of calculating an irradiance for a range of wavelengths without having to define a waveband:

```
with(sun.data, split_energy_irradiance(w.length, s.e.irrad, c(400,
700)))

## range.400.700
## 196.7
```

6.11 Task: calculate the contribution of different regions of a spectrum to photon irradiance

The function `split_photon_irradiance` takes the same arguments as the equivalent function for photon irradiance, consequently only one code example is provided here (see section 6.10 for more details):

```
with(sun.data, split_photon_irradiance(w.length, s.e.irrad, c(400, 500, 600, 700), scale = "percent"))

## range.400.500 range.500.600 range.600.700
##          29.41          35.14          35.45
```


CHAPTER 7

Weighted and effective irradiance

Abstract

In this chapter we explain how to .

7.1 Task:

CHAPTER 8

Colour

Abstract

In this chapter we explain how to .

8.1 Task:

CHAPTER 9

Photoreceptors

Abstract

In this chapter we explain how to .

9.1 Task:

CHAPTER 10

Radiation sources

Abstract

In this chapter we explain how to .

10.1 Task:

Filters

Abstract

In this chapter we explain how to .

11.1 Task:

CHAPTER 12

Plotting spectra

Abstract

In this chapter we explain how to .

12.1 Task:

Calibration

Abstract

In this chapter we explain how to .

13.1 Task:

Simulation

Abstract

In this chapter we explain how to .

14.1 Task:

CHAPTER 15

Measurement

Abstract

In this chapter we explain how to .

15.1 Task:

CHAPTER 16

Optimizing performance

Abstract

In this chapter we explain how to .

16.1 Task: