

# Homework Sheet 2

Author: Abdullah Oğuz Topçuoğlu

## Problem 5

We are given the function

$$f(x_1, x_2) := |x_1| + |x_2|.$$

We are looking for points  $x^0 \in \mathbb{R}^2$  where  $\frac{\partial f}{\partial x_1^0}$  and  $\frac{\partial f}{\partial x_2^0}$  exist. First things first that we realize the function is symmetric in both variables, so we can just focus on one of them and the other will follow the same logic.

Let's consider the partial derivative with respect to  $x_1^0$ :

$$\begin{aligned}\frac{\partial f}{\partial x_1^0} &= \lim_{h \rightarrow 0} \frac{f(x_1^0 + h, x_2^0) - f(x_1^0, x_2^0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{|x_1^0 + h| + |x_2^0| - (|x_1^0| + |x_2^0|)}{h} \\ &= \lim_{h \rightarrow 0} \frac{|x_1^0 + h| - |x_1^0|}{h}.\end{aligned}$$

Now we have to consider different cases for  $x_1^0$ :

- If  $x_1^0 > 0$ :

$$\frac{\partial f}{\partial x_1^0} = \lim_{h \rightarrow 0} \frac{(x_1^0 + h) - x_1^0}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1.$$

- If  $x_1^0 < 0$ :

$$\frac{\partial f}{\partial x_1^0} = \lim_{h \rightarrow 0} \frac{-(x_1^0 + h) + x_1^0}{h} = \lim_{h \rightarrow 0} \frac{-h}{h} = -1.$$

- If  $x_1^0 = 0$ :

$$\begin{aligned}\frac{\partial f}{\partial x_1^0} &= \lim_{h \rightarrow 0^+} \frac{|h| - 0}{h} = \lim_{h \rightarrow 0^+} \frac{h}{h} = 1, \\ \frac{\partial f}{\partial x_1^0} &= \lim_{h \rightarrow 0^-} \frac{|-h| - 0}{h} = \lim_{h \rightarrow 0^-} \frac{-h}{h} = -1.\end{aligned}$$

Since the left-hand limit and right-hand limit are not equal, the partial derivative does not exist at this point.

So partial derivatives exist for all points where  $x_1^0 \neq 0$ . By symmetry, the same applies for  $x_2^0$ .

**Partial Derivatives:**

$$\frac{\partial f}{\partial x_1^0} = \begin{cases} 1, & x_1^0 > 0 \\ -1, & x_1^0 < 0 \end{cases},$$

$$\frac{\partial f}{\partial x_2^0} = \begin{cases} 1, & x_2^0 > 0 \\ -1, & x_2^0 < 0 \end{cases}$$

## Problem 6

We are given the set

$$D := \{(x_1, x_2) \in \mathbb{R}^2 : x_1 \neq x_2^2, x_2 > 0\}$$

and the function

$$f : D \rightarrow \mathbb{R}, \quad f(x_1, x_2) := \frac{x_1 \ln(x_2)}{(x_1 - x_2^2)x_2}.$$

(i)

We need to show that for any point  $x \in D$  we can find an open ball around it.




Figure 1: The set  $D$

So if we pick a point in  $D$  we can get the distance to  $x_1$  axis and distance to the parabola in the image and if we choose the radius of the open ball to be the smaller of both then every point in that open ball will be in  $D$  because we are guaranteed to not touch the  $x_1$  axis and the parabola. That's why  $D$  is open.

(ii)

We want to show that for any point in the set  $D$  the partial derivatives exist.  
Let's start with  $\frac{\partial f}{\partial x_1}$ :

Define  $f_{x_2}(x_1) = f(x_1, x_2)$ . Now we want to show that  $\frac{df_{x_2}}{dx_1}$  exists.

And yes it does because  $f_{x_2}(x_1)$  is a rational function where the numerator and denominator are polynomials in  $x_1$  (with  $x_2$  being constant) and denominator is never zero.

Now we want to show that  $\frac{\partial f}{\partial x_2}$  exists:

Define  $f_{x_1}(x_2) = f(x_1, x_2)$ . Now we want to show that  $\frac{df_{x_1}}{dx_2}$  exists.

And yes it does again because the numerator is differentiable and the denominator is also differentiable and never zero.

**Partial derivatives:**

$$\frac{\partial f}{\partial x_1} = \frac{df_{x_2}}{dx_1}$$

apply quotient rule

$$\begin{aligned} &= \frac{\ln(x_2)((x_1 - x_2^2)x_2) - x_1 \ln(x_2)x_2}{(x_1 - x_2^2)^2 x_2^2}, \\ &= \frac{\ln(x_2)x_2(x_1 - x_2^2 - x_1)}{(x_1 - x_2^2)^2 x_2^2}, \\ &= \frac{\ln(x_2)x_2(-x_2^2)}{(x_1 - x_2^2)^2 x_2^2}, \\ &= \frac{-\ln(x_2)x_2^3}{(x_1 - x_2^2)^2 x_2^2}, \\ &= \frac{-\ln(x_2)x_2}{(x_1 - x_2^2)^2}, \end{aligned}$$

$$\frac{\partial f}{\partial x_2} = \frac{df_{x_1}}{dx_2}$$

apply quotient rule

$$\begin{aligned}
&= \frac{x_1/x_2((x_1 - x_2^2)x_2) - (x_1(\ln(x_2)(x_1 - 3x_2^2)))}{(x_1 - x_2^2)^2 x_2^2} \\
&= \frac{x_1(x_1 - x_2^2) - (x_1 \ln(x_2)(x_1 - 3x_2^2))}{(x_1 - x_2^2)^2 x_2^2} \\
&= \frac{x_1(x_1 - x_2^2 - \ln(x_2)(x_1 - 3x_2^2))}{(x_1 - x_2^2)^2 x_2^2}
\end{aligned}$$

This is the furthest i could simplify :)

**(iii)**

The gradient is simply the vector of partial derivatives which we just calculated in (ii):

$$\begin{aligned}
\nabla f(x_1, x_2) &= \begin{pmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \end{pmatrix} \\
&= \begin{pmatrix} \frac{-\ln(x_2)x_2}{(x_1 - x_2^2)^2} \\ \frac{x_1(x_1 - x_2^2 - \ln(x_2)(x_1 - 3x_2^2))}{(x_1 - x_2^2)^2 x_2^2} \end{pmatrix}
\end{aligned}$$