

Mathematics Homework Sheet 7

Authors: Abdullah Oguz Topcuoglu & Ahmed Waleed Ahmed Badawy
Shora

Problem 1

A transformation T is linear if and only if it satisfies the following two properties for all vectors u, v and scalar c :

1. $T(u + v) = T(u) + T(v)$
 2. $T(cu) = cT(u)$
1. $T : \mathbb{R}^2 \rightarrow \mathbb{R}, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto x + 2y$ is linear.
 2. $T : \mathbb{R}^2 \rightarrow \mathbb{R}, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto x + y^2$ is not linear.
Because rule (1) is not satisfied.
 3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto xy$ is not linear.
Because rule (2) is not satisfied.
 4. $T : \mathbb{C} \rightarrow \mathbb{C}, z \mapsto \bar{z}$ is linear.
 5. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x + 1 \\ y - 1 \end{pmatrix}$ is not linear.
Because rule (1) is not satisfied.
 6. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x - y \\ x + 2y \end{pmatrix}$ is linear.
 7. $T : \mathbb{R}^n[x] \rightarrow \mathbb{R}, p(x) \mapsto p(1)$ is linear.
 8. $T : \mathbb{R}^n[x] \rightarrow \mathbb{R}^{n+2}[x], p(x) \mapsto x^2 p(x)$ is linear.

Problem 2

(a)

The way we construct the matrix given a linear transformation is that we take the each basis vector and apply the transformation to it and write the result to corresponding column of the matrix.

$$\begin{aligned}
T(e_1) &= T(1, 0, 0) = (1 - 0, 1 + 2 \cdot 0 - 0, 2 \cdot 1 + 0 + 0) = (1, 1, 2) \\
T(e_2) &= T(0, 1, 0) = (0 - 1, 0 + 2 \cdot 1 - 0, 2 \cdot 0 + 1 + 0) = (-1, 2, 1) \\
T(e_3) &= T(0, 0, 1) = (0 - 0, 0 + 2 \cdot 0 - 1, 2 \cdot 0 + 0 + 1) = (0, -1, 1)
\end{aligned}$$

Put each result as a column in the matrix A :

$$A = \begin{pmatrix} 1 & -1 & 0 \\ 1 & 2 & -1 \\ 2 & 1 & 1 \end{pmatrix}$$

(b)

The standard basis looks like this: $\{1, x, \dots, x^n\}$

$$\begin{aligned}
T(1) &= 1 \\
T(x) &= x + 1 \\
T(x^2) &= (x + 1)^2 = x^2 + 2x + 1 \\
&\vdots \\
T(x^k) &= (x + 1)^k = \sum_{i=0}^k \binom{k}{i} x^i \\
&\vdots \\
T(x^n) &= (x + 1)^n
\end{aligned}$$

$$m_{i,j} = \begin{cases} \binom{j}{i} & \text{if } 0 \leq i \leq j \leq n \\ 0 & \text{if } i > j \end{cases}$$

Each column of the matrix has the coefficients of a binomial expansion of $(x + 1)^k$ for $k = 0, 1, \dots, n$. For example the first column looks like this:

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

The second column looks like this:

$$\begin{pmatrix} 1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

(c)

The standard basis of $\mathbb{R}^{2 \times 2}$ is:

$$E_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad E_{12} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad E_{21} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad E_{22} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

We compute $T(E_{ij})$ for each basis element:

$$T(E_{11}) = T\left(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}\right) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = E_{11}$$

$$T(E_{12}) = T\left(\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}\right) = \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix} = 2E_{12}$$

$$T(E_{21}) = T\left(\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}\right) = \begin{pmatrix} 0 & 0 \\ 3 & 0 \end{pmatrix} = 3E_{21}$$

$$T(E_{22}) = T\left(\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}\right) = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix} = 4E_{22}$$

Thus, the matrix of T with respect to the basis $\{E_{11}, E_{12}, E_{21}, E_{22}\}$ is:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 4 \end{pmatrix}$$