

Mathematics Homework Sheet 2

Authors: Abdullah Oguz Topcuoglu & Ahmed Waleed Ahmed
Badawy Shora

Problem 1

(a)

Z_5 :

\oplus	0	1	2	3	4
0	0	1	2	3	4
1	1	2	3	4	0
2	2	3	4	0	1
3	3	4	0	1	2
4	4	0	1	2	3

\times	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	1	3
3	0	3	1	4	2
4	0	4	3	2	1

Z_7 :

\oplus	0	1	2	3	4	5	6
0	0	1	2	3	4	5	6
1	1	2	3	4	5	6	0
2	2	3	4	5	6	0	1
3	3	4	5	6	0	1	2
4	4	5	6	0	1	2	3
5	5	6	0	1	2	3	4
6	6	0	1	2	3	4	5

\times	0	1	2	3	4	5	6
0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6
2	0	2	4	6	1	3	5
3	0	3	6	2	5	1	4
4	0	4	1	5	2	6	3
5	0	5	3	1	6	4	2
6	0	6	5	4	3	2	1

(b)

Z_4 :

\oplus	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

\times	0	1	2	3
0	0	0	0	0
1	0	1	2	3
2	0	2	0	2
3	0	3	2	1

(Z_4, \oplus, \times) is not a field because $[2]$ does not have a multiplicative inverse.

Problem 2

(a)

Let $f : Z_n \rightarrow Z_n$ be:

$$f([i]) = [i] + [m], \quad [m] \in Z_n$$

We want to show that f is a bijection.

- **Injective:** Suppose $f([i_1]) = f([i_2])$:

$$[i_1] + [m] = [i_2] + [m]$$

Thus,

$$[i_1] = [i_2]$$

Thus, f is injective.

- **Surjective:** Let $[j] \in Z_n$. We want to show that there exists an $[x]$ such that $f([x]) = [j]$:

$$f([j - m]) = [j - m] + [m] = [j]$$

Thus, f is surjective.

Thus, f is a bijection.

(b)

$$\begin{aligned} [i] &\rightarrow [i] + [1]: (01234567) \\ [i] &\rightarrow [i] + [2]: (0246)(1357) \\ [i] &\rightarrow [i] + [3]: (03614725) \\ [i] &\rightarrow [i] + [4]: (04)(15)(26)(37) \end{aligned}$$

Problem 3

(a)

Let $f : Z_p \setminus \{[0]\} \rightarrow Z_p \setminus \{[0]\}$ be:

$$f([i]) = [m].[i], \quad [m] \in Z_p \setminus \{[0]\}$$

We want to show that f is a bijection.

- **Injective:** Suppose $f([i_1]) = f([i_2])$:

$$[m].[i_1] = [m].[i_2]$$

Thus,

$$[i_1] = [i_2]$$

Thus, f is injective.

- **Surjective:** Let $[j] \in Z_p \setminus \{[0]\}$. We want to show that there exists an $[x]$ such that $f([x]) = [j]$:

$$f([j.m^{-1}]) = [j.m^{-1}].[m] = [j]$$

Thus, f is surjective.

Thus, f is a bijection.

(b)

$$\begin{aligned} [i] &\rightarrow [6].[i]: (0)(16)(25)(34) \\ [i] &\rightarrow [2].[i]: (0)(124)(365) \end{aligned}$$

Problem 4

Let I be an arbitrary interval. We will show that I and \mathbb{R} have the same cardinality by constructing a bijection between them.

Problem 5

(a)

Let $(p, n), (q, m) \in N_0 \times N_0$. We want to show that \sim is an equivalence relation on $N_0 \times N_0$:

- **Reflexive:** We want to show that $(p, n) \sim (p, n)$:

$$p + n = p + n$$

Thus, \sim is reflexive.

- **Symmetric:** We want to show that if $(p, n) \sim (q, m)$, then $(q, m) \sim (p, n)$:

$$p + m = q + n$$

Thus,

$$q + n = p + m$$

Thus, \sim is symmetric.

- **Transitive:** We want to show that if $(p, n) \sim (q, m)$ and $(q, m) \sim (r, s)$, then $(p, n) \sim (r, s)$:

$$p + m = q + n$$

and

$$q + s = r + m$$

Thus,

$$p + s = r + n$$

Thus, \sim is transitive.

Thus, \sim is an equivalence relation on $N_0 \times N_0$.

(b)

Let $k \in N_0$. We want to show that $(p, n) \sim (k + p, k + n)$:

$$p + (k + n) = (k + p) + n$$

Thus, $(p, n) \sim (k + p, k + n)$.

(c)

Let $[(k, 0)]$ be the equivalence class of $(k, 0)$. We want to find an equivalence class $-[(k, 0)]$ such that

$$-[(k, 0)] + [(k, 0)] = [(0, 0)]$$

Define $-[(k, 0)]$ as $[(0, k)]$. Then,

$$[(0, k)] + [(k, 0)] = [(0 + k, k + 0)] = [(k, k)] = [(0, 0)]$$

Thus, we have

$$-[(k, 0)] + [(k, 0)] = [(0, 0)]$$

Thus, $-[(k, 0)]$ is the equivalence class with the property that $-[(k, 0)] + [(k, 0)] = [(0, 0)]$.