

# Homework Sheet 7

Author: Abdullah Oğuz Topçuoğlu

## Exercise 25

We choose the probability space

$$\begin{aligned}\Omega &= \{1, 2, 3, 4, 5, 6\}^3, \\ \mathcal{F} &= \text{Power set of } \Omega, \\ P(A) &= U_\Omega\end{aligned}$$

The events are

$$\begin{aligned}A_1 &= \{(a, b, c) \in \Omega \mid a + b < 5\}, \\ A_2 &= \{(a, b, c) \in \Omega \mid b \cdot c = 6\}, \\ A_3 &= \{(a, b, c) \in \Omega \mid b = 2\} \\ A_1 \cap A_3 &= \{(a, 2, c) \in \Omega \mid a < 3\}, \\ (A_1 \cup A_2) \setminus A_3 &= \{(a, b, c) \in \Omega \mid (a + b < 5) \text{ or } (b \cdot c = 6) \text{ and } b \neq 2\}\end{aligned}$$

Since  $P$  is uniform we count the number of elements in each event.

Calculating the probabilities:

6 many (a,b) pairs times 6 choices for c

$$P(A_1) = \frac{6 \cdot 6}{216} = \frac{36}{216} = \frac{1}{6},$$

4 many (b,c) pairs times 6 choices for a

$$P(A_2) = \frac{6 \cdot 4}{216} = \frac{24}{216} = \frac{1}{9},$$

6 choices for a times 1 choice for b times 6 choices for c

$$P(A_3) = \frac{36}{216} = \frac{1}{6},$$

2 choices for a times 1 choice for b times 6 choices for c

$$P(A_1 \cap A_3) = \frac{2 \cdot 6}{216} = \frac{1}{18},$$

For the last one we need to calculate  $|A_1 \cup A_2|$  first:

$$\begin{aligned}|A_1 \cup A_2| &= |A_1| + |A_2| - |A_1 \cap A_2|, \\ |A_1| &= 36, \\ |A_2| &= 24,\end{aligned}$$

Calculating  $|A_1 \cap A_2|$ :

if  $b = 1, c = 6 : a + 1 < 5 \Rightarrow a < 4 \Rightarrow 3$  choices

if  $b = 2, c = 3 : a + 2 < 5 \Rightarrow a < 3 \Rightarrow 2$  choices

if  $b = 3, c = 2 : a + 3 < 5 \Rightarrow a < 2 \Rightarrow 1$  choice

if  $b = 6, c = 1 : a + 6 < 5 \Rightarrow$  no choices.

So  $|A_1 \cap A_2| = 3 + 2 + 1 + 0 = 6$ .

$$|A_1 \cup A_2| = 36 + 24 - 6 = 54,$$

$$P(A_1 \cup A_2) = \frac{54}{216} = \frac{1}{4},$$

$$P((A_1 \cup A_2) \setminus A_3) = P(A_1 \cup A_2) - P(A_3) + P(A_1 \cap A_3) = \frac{1}{4} - \frac{1}{6} + \frac{1}{18} = \frac{5}{36}.$$