

Homework Sheet 3

Author: Abdullah Oğuz Topçuoğlu

Exercise 9

We are given the function:

$$f(x_1, x_2) := \begin{cases} \frac{x_1^2 x_2}{x_1^2 + x_2^2}, & (x_1, x_2) \neq (0, 0) \\ 0, & (x_1, x_2) = (0, 0) \end{cases}$$

(i)

To show that f is continuous at the point $(0, 0)$, we need to verify that every sequence $(x_1^{(n)}, x_2^{(n)})$ converging to $(0, 0)$, $f(x_1^{(n)}, x_2^{(n)})$ also converges to $f(0, 0) = (0, 0)$.

Fix a sequence $(x_1^{(n)}, x_2^{(n)})$ such that $(x_1^{(n)}, x_2^{(n)}) \rightarrow (0, 0)$.

Then, we have:

If $(x_1^{(n)}, x_2^{(n)}) \neq (0, 0)$

$$\begin{aligned} |f(x_1^{(n)}, x_2^{(n)}) - f(0, 0)| &= \left| \frac{(x_1^{(n)})^2 x_2^{(n)}}{(x_1^{(n)})^2 + (x_2^{(n)})^2} - 0 \right| \\ &= \left| \frac{(x_1^{(n)})^2 x_2^{(n)}}{(x_1^{(n)})^2 + (x_2^{(n)})^2} \right| \\ &\leq \left| \frac{(x_1^{(n)})^2 x_2^{(n)}}{(x_1^{(n)})^2} \right| \quad (\text{since } (x_1^{(n)})^2 + (x_2^{(n)})^2 \geq (x_1^{(n)})^2) \\ &= |x_2^{(n)}| \end{aligned}$$

$|x_2^{(n)}| \rightarrow 0$ since $(x_1^{(n)}, x_2^{(n)}) \rightarrow (0, 0)$. Therefore $f(x_1^{(n)}, x_2^{(n)}) \rightarrow (0, 0)$.

If $(x_1^{(n)}, x_2^{(n)}) = (0, 0)$

$$|f(x_1^{(n)}, x_2^{(n)}) - f(0, 0)| = |0 - 0| = 0$$

which also converges to 0.

That's what we wanted to show.

(ii)

We need to show that the following limit doesn't converge to zero

$$\lim_{x \rightarrow 0} \frac{f(x) - f(0) - Jf(0)x}{\|x\|}$$

The J is the Jacobian matrix.

$$J = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \frac{\partial f}{\partial x_2} \end{bmatrix}$$

Lets compute the partials at $(0, 0)$

$$\begin{aligned} \frac{\partial f}{\partial x_1}(0, 0) &= \lim_{h \rightarrow 0} \frac{f(h, 0) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0 - 0}{h} = 0 \\ \frac{\partial f}{\partial x_2}(0, 0) &= \lim_{h \rightarrow 0} \frac{f(0, h) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0 - 0}{h} = 0 \end{aligned}$$

Thus, the Jacobian matrix at $(0, 0)$ is:

$$Jf(0, 0) = \begin{bmatrix} 0 & 0 \end{bmatrix}$$

Substituting this into our limit we get

$$\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{\|x\|}$$

We also know that $f(0) = 0$, so we can simplify this to:

$$\lim_{x \rightarrow 0} \frac{f(x)}{\|x\|}$$

So it is enough if i can show a vector x where this limit is not zero.
Lets choose $x = (x_1, x_1)$. Then we have:

$$\begin{aligned} \lim_{x_1 \rightarrow 0} \frac{f(x_1, x_1)}{\|(x_1, x_1)\|} &= \lim_{x_1 \rightarrow 0} \frac{\frac{x_1^2 x_1}{x_1^2 + x_1^2}}{\sqrt{x_1^2 + x_1^2}} \\ &= \lim_{x_1 \rightarrow 0} \frac{\frac{x_1^3}{2x_1^2}}{\sqrt{2}|x_1|} \\ &= \lim_{x_1 \rightarrow 0} \frac{x_1}{2\sqrt{2}|x_1|} \\ &= \lim_{x_1 \rightarrow 0} \frac{1}{2\sqrt{2}} \cdot \frac{x_1}{|x_1|} \end{aligned}$$

This limit doesnt even exist so it cant be equal to zero.
Thats what we wanted to show.

Exercise 10

We are given the function:

$$f(x_1, x_2, x_3) := \begin{bmatrix} x_1^4 \ln(3 + 2x_2^2) \\ x_1 \sin(x_2 x_3) e^{x_1} \end{bmatrix}$$

(i)

To show that f is a C^1 function we need to verify that all partial derivatives of f exist and are continuous.

Lets compute the partial derivatives of each component of f which corresponds to the entries of Jacobian matrix.

$$Jf = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \frac{\partial f_1}{\partial x_3} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \frac{\partial f_2}{\partial x_3} \end{bmatrix}$$

Calculating the partial derivatives:

$$\begin{aligned} \frac{\partial f_1}{\partial x_1} &= 4x_1^3 \ln(3 + 2x_2^2) \\ \frac{\partial f_1}{\partial x_2} &= x_1^4 \cdot \frac{4x_2}{3 + 2x_2^2} \\ \frac{\partial f_1}{\partial x_3} &= 0 \\ \frac{\partial f_2}{\partial x_1} &= \sin(x_2 x_3) e^{x_1} + x_1 \sin(x_2 x_3) e^{x_1} \\ \frac{\partial f_2}{\partial x_2} &= x_1 x_3 \cos(x_2 x_3) e^{x_1} \\ \frac{\partial f_2}{\partial x_3} &= x_1 x_2 \cos(x_2 x_3) e^{x_1} \end{aligned}$$

All these partial derivatives are polynomials, logarithmic, exponential, sine and cosine functions which are continuous everywhere in \mathbb{R}^3 Thus, all partial derivatives of f exist and are continuous.

(we can explicitly use the remark 1.3.13 in lecture notes to prove continuity but thats trivial so i cut it short with one sentence. Can i cut it short like this in the exam?)

Thats what we wanted to show.

(ii)

Yes it is because we know that continuously partially differentiability implies total differentiability. In (i) we showed that f is continuously partially differentiable function.

Exercise 11

We are given the functions:

$$f_1(x) := \begin{bmatrix} x^2 \\ 1 \end{bmatrix}, \quad f_2(y_1, y_2) := \begin{bmatrix} \sin(y_1) \\ \cos(y_2) \\ \sin(y_1) + \cos(y_2) \end{bmatrix}, \quad f_3(z_1, z_2, z_3) := e^{z_1 + z_2 + z_3}$$

(i)

Chain rule says that if f and g are totally differentiable then $g \circ f$ is also totally differentiable. And it is obvious that f_1, f_2, f_3 are totally differentiable since they are polynomials, sine, cosine and exponential functions. (all the partial derivatives are continuous everywhere and as we showed in exercise 10 (ii) that implies total differentiability)

This is why $f_3 \circ f_2 \circ f_1$ is also totally differentiable.