

Mathematics Homework Sheet 5

Author: Abdullah Oguz Topcuoglu & Yousef Farag

Problem 1

Problem 1(b)

Problem 1(b)(i)

We want to prove $\bigcap_{i \in I} U_i$ is closed.

We are given $(\forall i \in I \ U_i \subseteq R)$ is closed.

A set being closed means that its complement is open. So we want to prove that $\bigcup_{i \in I} U_i^c$ is open.

Since each U_i is closed, we know that U_i^c is open.

And from the lecture we know that union or intersection of open sets is open.

Thus $\bigcup_{i \in I} U_i^c$ is open. Which means that $\bigcap_{i \in I} U_i$ is closed.

And this completes the proof.

Problem 1(b)(ii)

We want to prove $\bigcup_{i=1}^n U_i$ is closed.

We are given $(U_1, \dots, U_n \subseteq R)$ are closed.

A set being closed means that its complement is open. So we want to prove that $\bigcap_{i=1}^n U_i^c$ is open.

Since each U_i is closed, we know that U_i^c is open.

And from the lecture we know that union or intersection of open sets is open.

Thus $\bigcap_{i=1}^n U_i^c$ is open. Which means that $\bigcup_{i=1}^n U_i$ is closed.

And this completes the proof.

Problem 3

Problem 3(a)

$$a_n := (-1)^n$$

a_n is not convergent. Because, for example, if we take $\epsilon = 1/10$ then there is no N that satisfies

$$\forall n \geq N \quad |a_n - a| < 1/10$$

a_n alternates between -1 and 1. So we can't find a value a that stays in the neighborhood of both -1 and 1. For example, when $\epsilon = 1/10$ there is no $a \in R$

that satisfies

$$|1 - a| < 1/10 \quad \text{and} \quad |-1 - a| < 1/10$$

No matter what you choose N to be you will always get for some $j \in N$ $a_j = 1$ and $a_j = -1$.

Thus a_n is divergent.

Problem 3(b)

$$b_n := \frac{(-1)^n}{n}$$

b_n is convergent.

Because, we can find an N that satisfies

$$\forall \epsilon > 0 \quad \forall n \geq N \quad |b_n - 0| < \epsilon$$

We are trying to find an N such that this inequality holds for any choice of ϵ .

$$\left| \frac{(-1)^n}{n} \right| < \epsilon \quad \text{when } n \geq N$$

When n is even, we have

$$\begin{aligned} \left| \frac{1}{n} \right| &< \epsilon && \text{when } n \geq N \\ \frac{1}{n} &< \epsilon && \text{when } n \geq N \\ n &> \frac{1}{\epsilon} && \text{when } n \geq N \end{aligned}$$

So if we choose N to be the any integer greater than $\frac{1}{\epsilon}$ then the inequality holds for even n . So, such N exist when n is even.

When n is odd, we have

$$\begin{aligned} \left| \frac{-1}{n} \right| &< \epsilon && \text{when } n \geq N \\ \frac{1}{n} &< \epsilon && \text{when } n \geq N \\ n &> \frac{1}{\epsilon} && \text{when } n \geq N \end{aligned}$$

Basically, we have the same thing for odd n . So, such N exist when n is odd too.

Thus, we can find an N that satisfies the inequality for any choice of ϵ if we choose a to be 0.

Since $a = 0$, the limit is zero.

$$\lim_{n \rightarrow \infty} b_n = \frac{(-1)^n}{n} = 0$$