Homework Sheet 1

Author: Abdullah Oğuz Topçuoğlu

Problem 1

We are given the function:

$$f: \mathbb{R}^2 \to \mathbb{R}$$
 $f(x_1, x_2) = x_1^2 + 2x_2^2$

We want to show that f is continuous on \mathbb{R}^2 . We want to find a delta that satisfies:

$$\forall \varepsilon > 0, \exists \delta > 0 : \forall x \in \mathbb{R}^2 ||x - x^0|| < \delta \implies |f(x) - f(x^0)| < \varepsilon$$

for an arbitrary but fixed x^0 .

Fix $x^0 = (x_1^0, x_2^0)$. Fix $\varepsilon > 0$. We want to find a delta that makes this inequality satisfied always:

$$|f(x) - f(x^0)| < \varepsilon$$

We are gonna find a relation between delta and $|f(x)-f(x^0)|$ and from there we are look for values of delta where it is less than epsilon.

$$|f(x) - f(x^{0})| = |x_{1}^{2} + 2x_{2}^{2} - (x_{1}^{0})^{2} - 2(x_{2}^{0})^{2}|$$

$$= |(x_{1}^{2} - (x_{1}^{0})^{2}) + 2(x_{2}^{2} - (x_{2}^{0})^{2})|$$

$$\leq |x_{1}^{2} - (x_{1}^{0})^{2}| + 2|x_{2}^{2} - (x_{2}^{0})^{2}|$$

$$= |(x_{1} - x_{1}^{0})(x_{1} + x_{1}^{0})| + 2|(x_{2} - x_{2}^{0})(x_{2} + x_{2}^{0})|$$

$$\leq |x_{1} - x_{1}^{0}||x_{1} + x_{1}^{0}| + 2|x_{2} - x_{2}^{0}||x_{2} + x_{2}^{0}|$$

(replace invidual components of the vector with the vector itself. that makes the overall value larger)

$$\leq \|x - x^0\| \|x + x^0\| + 2\|x - x^0\| \|x + x^0\|$$

$$= 3\|x - x^0\| \|x + x^0\|$$

$$\leq 3\delta \|x + x^0\|$$

and now assume that $\delta \leq 1$, then we can say that:

$$||x|| \le ||x^0|| + 1$$

$$|f(x) - f(x^{0})| \le 3\delta ||x + x^{0}||$$

$$\le 3\delta (||x|| + ||x^{0}||)$$

$$\le 3\delta (||x^{0}|| + 1 + ||x^{0}||)$$

$$= 3\delta (2||x^{0}|| + 1)$$

We are almost done. Now if we chose a delta that satisfies:

$$3\delta(2||x^0||+1) < \varepsilon$$
$$\delta < \frac{\varepsilon}{3(2||x^0||+1)}$$

And at the top we assumed that $\delta \leq 1$, so we can chose:

$$\delta = \min\left(1, \frac{\varepsilon}{3(2\|x^0\| + 1)}\right)$$

So found a delta meaning that f is continuous at x^0 . Since x^0 was arbitrary, f is continuous on \mathbb{R}^2 .