

Laplacian Subdivision Surface

Alexander Pinzón Fernandez

Advisor: Eduardo Romero

Universidad Nacional de Colombia

October 11, 2012

Outline



- 1 Introduction
- 2 Normalized LBO over triangles and quads
- 3 GSOC 2012 Mesh Smoothing



Laplacian smooth



The functional used in many laplacian smoothing approach to constrain energy minimization is based on a total curvature of a surface S .

$$E(S) = \int_S \kappa_1^2 + \kappa_2^2 dS \quad (1)$$



Gradient of Voronoi Area

Consider a surface S compound by a set of triangles around vertex v_i . We can define the *Voronoi region* of v_i , the change in area produced by move v_i is named gradient of *Voronoi region*.

$$\nabla A = \frac{1}{2} \sum_j (\cot \alpha_j + \cot \beta_j) (v_i - v_j) \quad (2)$$




Figure: Area of Voronoi region around v_i in dark blue. v_j 1-ring neighbors around v_i . α_j and β_j opposite angles to edge $\overrightarrow{v_j - v_i}$.

Laplace Beltrami Operator



If we normalize this gradient in equation (2) by the total area in 1-ring around v_i , we have the *discrete mean curvature normal* of a surface S .

$$2\kappa\mathbf{n} = \frac{\nabla A}{A} \quad (3)$$

The LBO has desirable features, one feature of the LBO is in direction of surface area minimization, allowing us to minimize energy using it on a total curvature of a surface S equation (1).

$$\Delta_g S = 2\kappa\mathbf{n} \quad (4)$$



Normalized LBO over triangles and quads



$$L(i, j) = \begin{cases} -\frac{w_{ij}}{\sum_{j \in N(v_i)} w_{ij}} & \text{if } j \in N(v_i) \\ \delta_{ij} & \text{otherwise} \end{cases}$$

Where $N(v_i)$ is the 1-ring neighbors with shared face to v_i , and δ_{ij} being the Kronecker delta function.



The 5 basic triangle-quad cases

$$w_{ij} = \begin{cases} \frac{1}{2} (\cot \alpha_j + \cot \beta_j) & \text{case a.} \\ \frac{1}{4} (\cot \theta_{(j-1)1} + \cot \theta_{(j-1)4} + \cot \theta_{j3} + \cot \theta_{j6}) & \text{case b.} \\ \frac{1}{4} (\cot \theta_{j2} + \cot \theta_{j5}) & \text{case c.} \\ \frac{1}{4} (\cot \theta_{j3} + \cot \theta_{j6}) + \frac{1}{2} \cot \beta_j & \text{case d.} \\ \frac{1}{4} (\cot \theta_{(j-1)1} + \cot \theta_{(j-1)4}) + \frac{1}{2} \cot \alpha_j & \text{case e.} \end{cases}$$






Figure: The 5 basic triangle-quad cases with common vertex V_i and the relationship with V_j and V'_j . (a) Two triangles. (b) (c) Two quads and one quad. (d) (e) Triangles and quads.


Results

Enhancing Features




Results

Enhancing Features




Results

Enhancing Features




Blender



- Blender is the free open source 3D content creation suite, available for all major operating systems under the GNU General Public License..
- Use OpenNL for solve sparse system

Results



Bibliography



Alexa, M. and Boubekeur, T. 2008.

Subdivision shading.

In *ACM SIGGRAPH Asia 2008 papers*. SIGGRAPH Asia '08. ACM, New York, NY, USA, 142:1–142:4.



Belkin, M., Sun, J., and Wang, Y. 2008.

Discrete laplace operator on meshed surfaces.

In *Proceedings of the twenty-fourth annual symposium on Computational geometry*. SCG '08. ACM, New York, NY, USA, 278–287.



Biermann, H., Levin, A., and Zorin, D. 2000.

Piecewise smooth subdivision surfaces with normal control.

In *Proceedings of the 27th annual conference on Computer graphics and interactive techniques*. SIGGRAPH '00. ACM Press/Addison-Wesley Publishing Co., New York, NY, USA, 113–120.



Botsch, M., Pauly, M., Rossli, C., Bischoff, S., and Kobbelt, L. 2006.

Geometric modeling based on triangle meshes.

In *ACM SIGGRAPH 2006 Courses*. SIGGRAPH '06. ACM, New York, NY, USA.