

SOME TOPICS IN KNOT THEORY

A REPORT

*submitted in partial fulfillment of the requirements
for the award of the dual degree of*

Bachelor of Science-Master of Science

in

MATHEMATICS

by

APOORV POTNIS

(18343)

DEPARTMENT OF MATHEMATICS,
INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH
BHOPAL,
BHOPAL - 462 066

April 2023

CERTIFICATE

This is to certify that **Apoorv Potnis**, BS-MS (Mathematics), has worked on the project entitled '**Some topics in knot theory**' under my supervision and guidance. The content of this report is original and has not been submitted elsewhere for the award of any academic or professional degree.

April 2023
IISER Bhopal

Dr. Dheeraj Kulkarni
Prof. Subhash Chaturvedi

Committee Member	Signature	Date
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

ACADEMIC INTEGRITY AND COPYRIGHT DISCLAIMER

I hereby declare that this project is my own work and, to the best of my knowledge, it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the award of any other degree or diploma at IISER Bhopal or any other educational institution, except where due acknowledgement is made in the document.

I certify that all copyrighted material incorporated into this document is in compliance with the Indian Copyright Act (1957) and that I have received written permission from the copyright owners for my use of their work, which is beyond the scope of the law. I agree to indemnify and save harmless IISER Bhopal from any and all claims that may be asserted or that may arise from any copyright violation.

**April 2023
IISER Bhopal**

Apoorv Potnis

ACKNOWLEDGEMENT

Acknowledgement is here.

Apoorv Potnis

ABSTRACT

We shall study some selected topics in knot theory from the book *Knots and Physics*, 4th ed. by Louis Kauffman [1]. The basics of knot theory have been studied from the book *Knots and Links* by Peter Cromwell [2].

A knot is an embedding of the circle in three-dimensional space. We discuss the state model for bracket polynomial, Jones polynomial and some of its generalisations. One can apply these knot invariants to show that various knots are not ambient isotopic, which is the notion we shall use to distinguish knots. We shall see a resolution of some of the old conjectures in knot theory using the bracket polynomial. Vaughn Jones found the polynomial named after him while he was studying towers of von Neumann algebras. We shall follow a different route following Kauffman to understand the relation amongst braids, links, Temperley–Lieb algebras, which arose in statistical physics, and the Jones algebras.

Contents

Certificate	i
Academic Integrity and Copyright Disclaimer	ii
Acknowledgement	iii
Abstract	iv
1 The Jones polynomial and its Generalisations	2
1.1 Jones polynomial	2
1.2 Alexander–Conway polynomial	4
1.3 HOMFLYPT polynomial	4

Chapter 1

The Jones polynomial and its Generalisations

1.1 Jones polynomial

Definition 1 (Jones polynomial). The Jones polynomial invariant $V(t, K)$ is defined as

$$V(t, K) := L\left(\frac{1}{t^{1/4}}, K\right)$$

for an oriented link K .

Here, $t = \frac{1}{A^4}$. $V(t, K)$ is a Laurent polynomial in $t^{1/2}$; $V(K) \in \mathbb{Z}\left[t^{1/2}, \frac{1}{t^{1/2}}\right]$. With the above definition, we have the following two properties of the Jones polynomial.

1. $V\left(t, \bigcirc\right) = 1$.
2. $\frac{1}{t} V\left(t, \begin{array}{c} \nearrow \\ \searrow \end{array}\right) - t V\left(t, \begin{array}{c} \nwarrow \\ \swarrow \end{array}\right) = \left(t^{1/2} - \frac{1}{t^{1/2}}\right) V\left(\begin{array}{c} \nearrow \\ \nwarrow \end{array}\right)$.

We can define the Jones polynomial axiomatically as a Laurent polynomial in $t^{1/2}$ satisfying the above two properties as well. But then we would have to check well-definedness and invariance under the Reidemeister moves. We derive these properties using the definition. The first property follows evaluating the bracket polynomial on the trivial knot. Orientation of a knot does not matter in this case. Just as with the normalised bracket polynomial, we shall drop the variable in which the polynomial is based if it is clear from the context.

Proof of the second property. We know that

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = A \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} + B \begin{array}{c} \diagup \diagdown \\ \diagup \diagup \end{array}$$

and

$$\begin{array}{c} \diagup \diagup \\ \diagdown \diagdown \end{array} = B \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array} + A \begin{array}{c} \diagup \diagdown \\ \diagup \diagup \end{array}.$$

Hence,

$$B^{-1} \begin{array}{c} \diagup \diagup \\ \diagdown \diagdown \end{array} - A^{-1} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = \left(\frac{A}{B} - \frac{B}{A}\right) \begin{array}{c} \diagup \diagdown \\ \diagup \diagup \end{array}.$$

Thus,

$$A \langle \text{crossing} \rangle - A^{-1} \langle \text{crossing} \rangle = (A^2 - A^{-2}) \langle \text{crossing} \rangle.$$

Let $w = w(\text{crossing})$ so that $w(\text{crossing}) = w - 1$ and $w(\text{crossing}) = w + 1$. Also, let $\alpha = -A^3$. Then

$$A\alpha \langle \text{crossing} \rangle \alpha^{-w+1} - A^{-1}\alpha^{-1} \langle \text{crossing} \rangle \alpha^{-w+1} = (A^2 - A^{-2}) \langle \text{crossing} \rangle \alpha^{-w}.$$

Thus,

$$-A^4 L(\text{crossing}) + A^{-4} L(\text{crossing}) = (A^2 - A^{-2}) L(\text{crossing}).$$

Substituting $A = t^{-1/4}$ yields the desired result. \square

The Jones polynomial follows a reversing property.

Theorem 2 (Reversing property). Let K be an oriented link. Let K' be such that K' is obtained by reversing the orientation of a component $K_i \subset K$.

Let $\lambda = \text{lk}(K_i, K - K_i)$ denote the total linking number of K_i with the remaining components of K . Then

$$V(t, K') = t^{-3\lambda} \cdot V(t, K).$$

Proof. We see that $w(K') = w(K) - 4\lambda$.

$$\begin{aligned} L(A, K') &= (-A^3)^{-w(K')} \langle K' \rangle \\ &= (-A^3)^{-w(K)+4\lambda} \langle K \rangle. \end{aligned}$$

Therefore $L(A, K') = (-A^3)^{4\lambda} L(A, K)$. Substituting $A = t^{-1/4}$ yields the desired result. \square

Example 3. Using the skein relation, we have that

$$\frac{1}{t} V(\text{link}) - t V(\text{link}) = \left(t^{1/2} - \frac{1}{t^{1/2}}\right) V(\text{link}).$$

But since both link and link are trivial knots,

$$V(\text{link}) = V(\text{link}) = V(\text{link}) = 1.$$

Thus,

$$V(\text{link}) = \frac{t^{-1} - t}{t^{1/2} - 1/t^{1/2}} = \frac{(t^{-1} - t)(t^{1/2} + 1/t^{1/2})}{t - t^{-1}} = -(t^{1/2} + 1/t^{1/2}) = \delta.$$

The third property is surprising when viewed with respect to the skein relation [3, 4]. Skein relations of this sort form a general defining feature for various polynomials.

1.2 Alexander–Conway polynomial

Definition 4 (Alexander–Conway polynomial). Let K be an oriented link diagram. Then the Alexander–Conway polynomial invariant $\nabla(z, K) \in \mathbb{Z}[z]$ is defined by the rules:

1. $\nabla\left(z, \begin{array}{c} \circlearrowright \end{array}\right) = 1.$
2. $\nabla\left(z, \begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) - \nabla\left(z, \begin{array}{c} \nearrow \nwarrow \\ \searrow \nearrow \end{array}\right) = \nabla\left(z, \begin{array}{c} \nearrow \nwarrow \\ \nearrow \nwarrow \end{array}\right).$

This polynomial is a generalisation and reformulation of the original Alexander polynomial.

1.3 HOMFLYPT polynomial

The HOMFLYPT polynomial was discovered independently by two groups, one group consisting of Jim Hoste, Adrian Ocneanu, Kenneth Millett, Peter Freyd, William Lickorish, and David Yetter and the other group consisting of Polish mathematicians Józef Przytycki and Paweł Traczyk.

Definition 5 (HOMFLYPT polynomial). Let K be an oriented link diagram. Then the HOMFLYPT polynomial invariant $P(\alpha, z, K)$ is defined by the rules:

1. $P\left(\alpha, z, \begin{array}{c} \circlearrowright \end{array}\right) = 1.$
2. $\alpha P\left(\alpha, z, \begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) - \frac{1}{\alpha} P\left(\alpha, z, \begin{array}{c} \nearrow \nwarrow \\ \searrow \nearrow \end{array}\right) = z P\left(\alpha, z, \begin{array}{c} \nearrow \nwarrow \\ \nearrow \nwarrow \end{array}\right).$

Bibliography

- [1] Louis H. Kauffman. *Knots and physics*. 4th ed. Knots and Everything 53. Singapore: World Scientific, 2013. ISBN: 978-981-4383-00-4.
- [2] Peter R. Cromwell. *Knots and links*. Cambridge, UK: Cambridge University Press, 2004. ISBN: 0-521-83947-5.
- [3] William Bernard Lickorish and Kenneth Millett. “The reversing result for the Jones polynomial”. In: *Pacific Journal of Mathematics* 124.1 (Sept. 1986), pp. 173–176. DOI: [10.2140/pjm.1986.124.173](https://doi.org/10.2140/pjm.1986.124.173).
- [4] Hugh R. Morton. “The Jones polynomial for unoriented links”. In: *The Quarterly Journal of Mathematics* 37.1 (1986), pp. 55–60. DOI: [10.1093/qmath/37.1.55](https://doi.org/10.1093/qmath/37.1.55).