

SOME TOPICS IN KNOT THEORY

A REPORT

*submitted in partial fulfillment of the requirements
for the award of the dual degree of*

Bachelor of Science-Master of Science

in

MATHEMATICS

by

APOORV POTNIS

(18343)

**DEPARTMENT OF MATHEMATICS,
INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH
BHOPAL,
BHOPAL - 462 066**

April 2023

CERTIFICATE

This is to certify that **Apoorv Potnis**, BS-MS (Mathematics), has worked on the project entitled '**Some topics in knot theory**' under my supervision and guidance. The content of this report is original and has not been submitted elsewhere for the award of any academic or professional degree.

April 2023
IISER Bhopal

Dr. Dheeraj Kulkarni
Prof. Subhash Chaturvedi

Committee Member	Signature	Date
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

ACADEMIC INTEGRITY AND COPYRIGHT DISCLAIMER

I hereby declare that this project is my own work and, to the best of my knowledge, it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the award of any other degree or diploma at IISER Bhopal or any other educational institution, except where due acknowledgement is made in the document.

I certify that all copyrighted material incorporated into this document is in compliance with the Indian Copyright Act (1957) and that I have received written permission from the copyright owners for my use of their work, which is beyond the scope of the law. I agree to indemnify and save harmless IISER Bhopal from any and all claims that may be asserted or that may arise from any copyright violation.

**April 2023
IISER Bhopal**

Apoorv Potnis

ACKNOWLEDGEMENT

Acknowledgement is here.

Apoorv Potnis

ABSTRACT

We shall study some selected topics in knot theory from the book *Knots and Physics*, 4th ed. by Louis Kauffman [1]. The basics of knot theory have been studied from the book *Knots and Links* by Peter Cromwell [2].

A knot is an embedding of the circle in three-dimensional space. We discuss the state model for bracket polynomial, Jones polynomial and some of its generalisations. One can apply these knot invariants to show that various knots are not ambient isotopic, which is the notion we shall use to distinguish knots. We shall see a resolution of some of the old conjectures in knot theory using the bracket polynomial. Vaughn Jones found the polynomial named after him while he was studying towers of von Neumann algebras. We shall follow a different route following Kauffman to understand the relation amongst braids, links, Temperley–Lieb algebras, which arose in statistical physics, and the Jones algebras.

Contents

Certificate	i
Academic Integrity and Copyright Disclaimer	ii
Acknowledgement	iii
Abstract	iv
1 The Jones polynomial and its Generalisations	2
1.0.1 Jones polynomial	2

Chapter 1

The Jones polynomial and its Generalisations

In the previous chapters, we defined Kauffman's bracket polynomial using the state model and then normalised it to get a link invariant. Instead of using the state model, one can define the bracket polynomial in an axiomatic way as well. One defines the polynomial as a function on link diagrams which satisfies certain properties. We would need to check the well-definedness and invariance of such a function under the Reidemeister moves. We thus define the bracket polynomial in the following way.

Definition 1 (Kauffman's bracket polynomial). Let K an unoriented link diagram. Then the bracket $\langle K \rangle \in \mathbb{Z}[A, A^{-1}]$ is defined by the rules:

1. $\langle \bigcirc \rangle = 1$.
2. $\langle \bigcirc \cup K \rangle = (-A^2 - A^{-2})\langle K \rangle$.
3. $\langle \text{X} \rangle = A \langle \text{X} \rangle + A^{-1} \langle \text{X} \rangle$.

This definition is consistent with the state model. $\langle K \rangle$ is invariant under the type II and type III moves. In order to gain type I invariance, we normalise it by adding an orientation and multiplying $\langle K \rangle$ with $(-A^3)^{-w(K)}$.

1.0.1 Jones polynomial

We now give an axiomatic definition of the Jones polynomial.

$$F \langle \text{X} \rangle G \langle \text{X} \rangle H \langle \text{X} \rangle I \langle \text{X} \rangle J \langle \text{X} \rangle K \langle \text{X} \rangle L \langle \bigcirc \rangle$$

Definition 2 (Jones polynomial). Let K be an oriented link diagram. Then the Jones polynomial $V(K) \in \mathbb{Z} \left[\sqrt{t}, \frac{1}{\sqrt{t}} \right]$ is defined by the rules:

1. $V \left(\bigcirc \right) = 1$.

$$2. \frac{1}{t} V \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) - t V \left(\begin{array}{c} \nwarrow \\ \swarrow \end{array} \right) = \left(\sqrt{t} - \frac{1}{\sqrt{t}} \right).$$

Bibliography

- [1] Louis H. Kauffman. *Knots and physics*. 4th ed. Knots and Everything 53. Singapore: World Scientific, 2013. ISBN: 978-981-4383-00-4.
- [2] Peter R. Cromwell. *Knots and links*. Cambridge, UK: Cambridge University Press, 2004. ISBN: 0-521-83947-5.