

The Quantum Mechanical Two-body Problem


These are lecture notes by Apoorv Potnis of the lecture ‘Quantenmechanisches Zweikörperproblem’ or ‘The Quantum Mechanical Two-body Problem’, given by **Prof. Frederic Paul Schuller**, as the seventeenth lecture in the course ‘Theoretische Physik 2: Theoretische Quantenmechanik’ in 2014/15 at the Friedrich-Alexander-Universität Erlangen-Nürnberg. While the original lecture is in German, these notes are in English and have been prepared using YouTube’s automatic subtitle translation tool. The lecture is available at https://www.youtube.com/watch?v=mcM4S3IMMvI&list=PLP05pgr_frzTeqa_thbltYjyw8F9ehw7v&index=17 and at <https://www.fau.tv/clip/id/44891>.

Contents


1 Introduction	1
2 Moving to spherical co-ordinates	4
3 Simultaneous eigenvectors	6
References	7

1 Introduction

Consider a classical system consisting of two interacting particles of masses m_1 and m_2 , such that the interaction is completely determined by the potential $V(r)$, and the potential depends only on the distance r between the particles. If we consider the central force problem, like a planet orbiting the sun, then we know we have bound states corresponding to *all* the negative energy values, and scattering states corresponding to all the positive energy values. It turns out that if consider the system quantum-mechanically, then bound states cannot admit all the negative energy solutions. Only certain



Classical picture



Yukawa and Coulomb potentials

energy levels are allowed. This can be seen experimentally from the discrete lines in the spectra of atoms, demonstrating the quantised nature of energy levels. In this lecture, we shall consider the quantum-mechanical case of the two-body problem.

An example of a potential as described above would be the *Yukawa potential*, defined as

$$V_{\text{Yukawa}}(r) := a \frac{\exp(-kmr)}{r},$$

where k , m and a are constants. $a \in \mathbb{R} \setminus \{0\}$, $m \geq 0$. According to quantum field theory, very roughly speaking, interaction between particles takes place via a ‘mediating particle’. If the interaction is mediated by a ‘scalar field’¹, then the mass associated to the particle of that scalar field is the mass m appearing in the Yukawa potential. If we plot a graph of the Yukawa potential for a massive scalar field, then we see that the magnitude of the potential becomes very close to zero after a certain distance. Thus, these interactions are short-ranged. If instead we have $m = 0$, corresponding to a

¹Whatever that means

photon, then we get the familiar long-range Coulomb potential

$$V_{\text{Coulomb}}(r) := a \frac{1}{r}.$$

We also have the finite wall potential $V_{\text{finite wall}}(r) := a\Theta(r - r_0)$ and the isotropic harmonic oscillator $V_{\text{ihc}}(r) := ar^2$.

The Hilbert space of the individual particles is $L^2(\mathbb{R}^3)$ and the Hilbert space of the composite two-body system is $L^2(\mathbb{R}^3) \otimes L^2(\mathbb{R}^3)$, which is naturally isomorphic to $L^2(\mathbb{R}^6)$. This can be seen from Theorem II.10 of the book of Reed and Simon [1, p. 52]. If $f, g \in L^2(\mathbb{R}^3)$, then we can define an isomorphism $L^2(\mathbb{R}^3) \otimes L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^6)$ by $f \otimes g \mapsto fg \in L^2(\mathbb{R}^6)$.

The Hamiltonian for our quantum-mechanical two-body system is

$$H := -\frac{\hbar^2}{2m_1}\Delta_{(1)} - \frac{\hbar^2}{2m_2}\Delta_{(2)} + V(\|\mathbf{x}^{(1)} - \mathbf{x}^{(2)}\|),$$

where $\Delta_{(i)}$ is the Laplacian operator which acts on the Hilbert space $L^2(\mathbb{R}^3)$ of the particle i , and $\mathbf{x}^{(i)} \in \mathbb{R}^3$ denotes the position vector of the i^{th} particle. We are interested in finding out the spectrum of this Hamiltonian.

As is the case in classical mechanics, we shall move to the center-of-mass co-ordinates. We can move the center-of-mass of the system in any direction by any value, i.e. the spectrum of the ‘center-of-mass’ position operator is \mathbb{R} . The discrete energy levels are due to the angular momentum about the center-of-mass. We thus introduce


$$\begin{aligned} \mathbf{y} &:= \frac{m_1\mathbf{x}^{(1)} + m_2\mathbf{x}^{(2)}}{m_1 + m_2}, \\ \mathbf{r} &:= \mathbf{x}^{(1)} - \mathbf{x}^{(2)}, \\ \mu &:= \frac{m_1m_2}{m_1 + m_2}. \end{aligned}$$

The Hamiltonian then takes the form

$$H := -\frac{\hbar^2}{2(m_1 + m_2)}\Delta_{\mathbf{y}} - \frac{\hbar^2}{2\mu}\Delta_{\mathbf{r}} + V(\|\mathbf{r}\|).$$

The wave function of the system $\psi(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})$ expressed in the new co-ordinates becomes

$$\phi(\mathbf{y}, \mathbf{r}) := \psi(\mathbf{x}^{(1)}(\mathbf{y}, \mathbf{r}), \mathbf{x}^{(2)}(\mathbf{y}, \mathbf{r})).$$



Spherical co-ordinates

The Hamiltonian H can be separated into two parts, H_{free} describing the movement of the center-of-mass and H_{rel} , describing the movement of the particles around the center-of-mass. We have

$$H_{\text{free}} := -\frac{\hbar^2}{2(m_1 + m_2)}\Delta_y \quad \text{and} \quad H_{\text{rel}} := -\frac{\hbar^2}{2\mu}\Delta_r + V(\|r\|).$$

We know that $\text{spec}(H_{\text{free}}) = \mathbb{R}_{\geq 0}$, so the task that remains is to find $\text{spec}(H_{\text{rel}})$. We see that H_{rel} is spherically symmetric. Thus, we shall move to the spherical co-ordinates to separate the angular and radial dependencies. This shall reduce our three-dimensional problem to a one-dimensional one.

2 Moving to spherical co-ordinates

Expressed in spherical co-ordinates, we have the wave-function as

$$F(r, \theta, \phi) := \psi(r(r, \theta, \phi)),$$

where

$$\begin{aligned} r_1 &= r \cos \phi \sin \theta, \\ r_2 &= r \sin \phi \sin \theta, \\ r_3 &= r \cos \theta, \end{aligned}$$

and r_1, r_2 and r_3 denote the x -, y - and z -components of r respectively. We now express the angular momentum operators in spherical co-ordinates as

well.

$$\begin{aligned}\tilde{L}_1 &:= i \cos \phi \cot \theta \frac{\partial}{\partial \phi} + i \sin \phi \frac{\partial}{\partial \theta}, \\ \tilde{L}_2 &:= i \sin \phi \cot \theta \frac{\partial}{\partial \phi} - i \cos \phi \frac{\partial}{\partial \theta}, \\ \tilde{L}_3 &:= -i \frac{\partial}{\partial \phi}.\end{aligned}$$

These can be derived by the application of the chain rule. If L_a denotes a^{th} ($a \in \{1, 2, 3\}$) angular momentum operator in Cartesian co-ordinates, then we have

$$(\tilde{L}_a F)(r, \theta, \phi) = (L_a \psi)(\mathbf{r}(r, \theta, \phi)).$$

We shall use the same letter for operators in Cartesian co-ordinates and spherical co-ordinates now on. We also have

$$L^2 F = -\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} F \right) - \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} \left(\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} F \right)$$

and

$$H_{\text{rel}} F = -\frac{\hbar^2}{2\mu} \left(\frac{1}{r} \left(\frac{\partial}{\partial r} \right)^2 (rF) - \frac{L^2}{r^2} F \right) + V(r)F.$$

In order to fully determine $\text{spec}(H_{\text{rel}})$, along with the continuous spectrum, i.e. the part containing the generalised eigenvalues, we would need to lift the eigenvalue equation to the distribution space and solve it.² However, we already know from experiments that the hydrogen atom has discrete lines in its observed spectrum. Thus, we shall first solve for the point spectrum

$$\text{spec}_p(H_{\text{rel}}) = \{E \in \mathbb{R} \mid H_{\text{rel}} F = EF\},$$

which was historically a major validation for the quantum theory.

²This can be achieved using Gelfand's rigged Hilbert space formalism, which was introduced in an earlier lecture. The lecture itself can be found on this link: https://www.youtube.com/watch?v=FNJOyxOp3Ik&list=PLP05pgr_frzTega_thb1tYjw8F9ehw7v&index=8. The notes for this lecture can be found here: https://github.com/apo-orvpotnis/schuller_rigged_hilbert_spaces/blob/main/schuller_rigged_hilbert_spaces.pdf.

3 Simultaneous eigenvectors

We shall find it using a technique which was employed earlier as well. We considered the simultaneous eigenvectors and eigenvalues of \mathbf{L}^2 and L_a , as they commute with each other. We thus consider the simultaneous eigenvalues of H_{rel} , \mathbf{L}^2 and L_3 in the hope that this will give us more equations to work with and make the job easier. It can be seen that all three operators commute with each other pairwise.

We know that the common eigenvectors of \mathbf{L}^2 and L_3 come as families ψ_l^m , where $m = -l, -l + 1, \dots, l - 1, l$ and $l \in \mathbb{N}_0$. We have

$$\begin{aligned}\mathbf{L}^2 \psi_l^m &= l(l+1) \psi_l^m, \\ L_3 \psi_l^m &= m \psi_l^m.\end{aligned}$$

Note that l can take only integer values, not half-integer values as we are dealing with the orbital angular momentum operator. Also recall the raising and lowering operators $L_+ := L_1 + iL_2$ and $L_- := L_1 - iL_2$. We have

$$L_{\pm} \psi_l^m = \sqrt{l(l+1) - m(m \pm 1)} \psi_l^{m \pm 1}.$$

In spherical co-ordinates, it turns out that the solutions ψ_l^m are given by

$$\psi_l^m(\mathbf{x}(r, \theta, \phi)) = k \cdot Y_l^m(\theta, \phi) \cdot f(r),$$

where

$$Y_l^m(\theta, \phi) := \frac{(-1)^m}{2^l l!} \left(\frac{(2l+1)(l-m)!}{4\pi(l+m)!} \right)^{\frac{1}{2}} e^{im\phi} (\sin \theta)^m \left(\frac{\partial}{\partial \cos \theta} \right)^{l+m} (\cos^2 \theta - 1)^l.$$

These Y_l^m 's are called as *spherical harmonic functions*. Since the operators \mathbf{L}^2 and L_3 act only on the angular components and not the radial components when expressed in spherical co-ordinates, we have that

$$\begin{aligned}\mathbf{L}^2 Y_l^m &= l(l+1) Y_l^m, \\ L_3 Y_l^m &= m Y_l^m,\end{aligned}$$

i.e. the spherical harmonic functions are eigenvectors of \mathbf{L}^2 and L_3 . These actually form an orthonormal basis of the Hilbert space $L^2(\mathbb{S}^2)$ consisting of complex square-integrable functions defined on the 2-sphere.

References

- [1] Michael Reed and Barry Simon. *Methods of Modern Mathematical Analysis I: Functional Analysis*. Revised and Enlarged edition. Vol. 1. Academic Press, Inc. London, 1980. ISBN: 978-0-080-57048-8.
- [2] Frederic Schuller, Simon Rea, and Richie Dadhley. ‘Lectures on Quantum Theory’. Lecturer: Prof. Frederic Paul Schuller. 2019. URL: https://docs.wixstatic.com/ugd/6b203f_a94140db21404ae69fd8b367d9fcd360.pdf.
- [3] Frederic Schuller. *Lectures on Quantum Theory*. 2015. URL: https://youtube.com/playlist?list=PLPH7f_7Z1zxQVx5jRjbfRGEzWY_upS5K6.
- [4] Philip Bowers. *Lectures on Quantum Mechanics*. Cambridge University Press, Cambridge, 2020. ISBN: 978-1-108-42976-4.

The source code, updates and corrections to this document can be found on this GitHub repository: https://github.com/apoorvpotnis/schuller_two-body_problem. The source code, along with the .bib file is embedded in this PDF. Comments and corrections can be mailed at apoorvpotnis@gmail.com.