

# The Quantum Mechanical Two-body Problem

These are lecture notes by Apoorv Potnis of the lecture ‘Quantenmechanisches Zweikörperproblem’ or ‘The Quantum Mechanical Two-body Problem’, given by **Prof. Frederic Paul Schuller**, as the seventeenth lecture in the course ‘Theoretische Physik 2: Theoretische Quantenmechanik’ in 2014/15 at the Friedrich-Alexander-Universität Erlangen-Nürnberg. While the original lecture is in German, these notes are in English and have been prepared using YouTube’s automatic subtitle translation tool. The lecture is available at [https://www.youtube.com/watch?v=mcM4S3IMMvI&list=PLP05pgr\\_frzTeqa\\_thbltYjyw8F9ehw7v&index=17](https://www.youtube.com/watch?v=mcM4S3IMMvI&list=PLP05pgr_frzTeqa_thbltYjyw8F9ehw7v&index=17) and at <https://www.fau.tv/clip/id/4489>.


The source code, updates and corrections to this document can be found on this GitHub repository: [https://github.com/apoorvpotnis/schuller\\_two-body\\_problem](https://github.com/apoorvpotnis/schuller_two-body_problem). The source code, along with the .bib file is embedded in this PDF. Comments and corrections can be mailed at [apoorvpotnis@gmail.com](mailto:apoorvpotnis@gmail.com).

## Contents


<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Moving to center-of-mass co-ordinates</b>	<b>3</b>
<b>3</b>	<b>Spectrum of angular momentum operators</b>	<b>4</b>
3.1	Justification of simultaneous eigenvectors . . . . .	5
3.2	Moving to spherical co-ordinates . . . . .	9
3.3	Spherical harmonics . . . . .	12
<b>4</b>	<b>Spectrum of the Hamiltonian</b>	<b>13</b>
	<b>References</b>	<b>13</b>

## 1 Introduction

Consider a classical system consisting of two interacting particles of masses  $m_1$  and  $m_2$ , such that the interaction is completely determined by the potential



Classical picture



Yukawa and Coulomb potentials

$V(r)$ , and the potential depends only on the distance  $r$  between the particles. If we consider the central force problem, like a planet orbiting the sun, then we know we have bound states corresponding to *all* the negative energy values, and scattering states corresponding to all the positive energy values. It turns out that if consider the system quantum-mechanically, then bound states cannot admit all the negative energy solutions. Only certain energy levels are allowed. This can be seen experimentally from the discrete lines in the spectra of atoms, demonstrating the quantised nature of energy levels. In this lecture, we shall consider the quantum-mechanical case of the two-body problem.

An example of a potential as described above would be the *Yukawa potential*, defined as

$$V_{\text{Yukawa}}(r) := a \frac{\exp(-kmr)}{r},$$

where  $k$ ,  $m$  and  $a$  are constants.  $a \in \mathbb{R} \setminus \{0\}$ ,  $m \geq 0$ . According to quantum field theory, very roughly speaking, interaction between particles takes place via a ‘mediating particle’. If the interaction is mediated by a ‘scalar field’<sup>1</sup>, then the mass associated to the particle of that scalar field is the mass  $m$  appearing in the

---

<sup>1</sup>Whatever that means

Yukawa potential. If we plot a graph of the Yukawa potential for a massive scalar field, then we see that the magnitude of the potential becomes very close to zero after a certain distance. Thus, these interactions are short-ranged. If instead we have  $m = 0$ , corresponding to a photon, then we get the familiar long-range Coulomb potential

$$V_{\text{Coulomb}}(r) := a \frac{1}{r}.$$

We also have the finite wall potential  $V_{\text{finite wall}}(r) := a\Theta(r - r_0)$ , where  $\Theta$  denotes the Heaviside function, and the isotropic harmonic oscillator  $V_{\text{ihc}}(r) := ar^2$ .

## 2 Moving to center-of-mass co-ordinates

The Hilbert space of the individual particles is  $L^2(\mathbb{R}^3)$  and the Hilbert space of the composite two-body system is  $L^2(\mathbb{R}^3) \otimes L^2(\mathbb{R}^3)$ , which is naturally isomorphic to  $L^2(\mathbb{R}^6)$ . This can be seen from theorem II.10 of the book of Reed and Simon [1, p. 52]. If  $f, g \in L^2(\mathbb{R}^3)$ , then we can define an isomorphism  $L^2(\mathbb{R}^3) \otimes L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^6)$  by  $f \otimes g \mapsto fg \in L^2(\mathbb{R}^6)$ , where  $fg$  denotes the point-wise product:  $(fg)(\mathbf{x}, \mathbf{y}) := f(\mathbf{x})g(\mathbf{y})$ . Tensor products of Hilbert spaces have been discussed in the fourteenth lecture ‘14: Composite Systems’ in the English quantum mechanics series [2, 3].

The Hamiltonian for our quantum-mechanical two-body system is

$$H := -\frac{\hbar^2}{2m_1}\Delta_{(1)} - \frac{\hbar^2}{2m_2}\Delta_{(2)} + V(\|\mathbf{x}^{(1)} - \mathbf{x}^{(2)}\|),$$

where  $\Delta_{(i)}$  is the Laplacian operator which acts on the Hilbert space  $L^2(\mathbb{R}^3)$  of the particle  $i$ , and  $\mathbf{x}^{(i)} \in \mathbb{R}^3$  denotes the position vector of the  $i^{\text{th}}$  particle. Assuming that this Hamiltonian is self-adjoint, we are interested in finding out its spectrum.

As is the case in classical mechanics, we shall move to the center-of-mass co-ordinates. We can move the center-of-mass of the system in any direction by any value, i.e. the spectrum of the ‘center-of-mass’ position operator is  $\mathbb{R}$ . The discrete energy levels are due to the angular momentum about the center-of-mass. We thus introduce

$$\begin{aligned}\mathbf{y} &:= \frac{m_1\mathbf{x}^{(1)} + m_2\mathbf{x}^{(2)}}{m_1 + m_2}, \\ \mathbf{r} &:= \mathbf{x}^{(1)} - \mathbf{x}^{(2)}, \\ \mu &:= \frac{m_1m_2}{m_1 + m_2}.\end{aligned}$$

The Hamiltonian then takes the form

$$H := -\frac{\hbar^2}{2(m_1 + m_2)}\Delta_{\mathbf{y}} - \frac{\hbar^2}{2\mu}\Delta_{\mathbf{r}} + V(\|\mathbf{r}\|).$$

The wave function of the system  $\psi(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})$  expressed in the new co-ordinates becomes

$$\phi(\mathbf{y}, \mathbf{r}) := \psi(\mathbf{x}^{(1)}(\mathbf{y}, \mathbf{r}), \mathbf{x}^{(2)}(\mathbf{y}, \mathbf{r})).$$

The Hamiltonian  $H$  can be separated into two parts,  $H_{\text{free}}$  describing the movement of the center-of-mass and  $H_{\text{rel}}$ , describing the movement of the particles around the center-of-mass. We have

$$H_{\text{free}} := -\frac{\hbar^2}{2(m_1 + m_2)}\Delta_{\mathbf{y}} \quad \text{and} \quad H_{\text{rel}} := -\frac{\hbar^2}{2\mu}\Delta_{\mathbf{r}} + V(\|\mathbf{r}\|).$$

We know that  $\text{spec}(H_{\text{free}}) = \mathbb{R}_{\geq 0}$ , so the task that remains is to find  $\text{spec}(H_{\text{rel}})$ , provided that  $H_{\text{rel}}$  is self-adjoint, depending on  $V(\|\mathbf{r}\|)$ . We see that  $H_{\text{rel}}$  is spherically symmetric. Thus, we shall shortly move on to the spherical co-ordinates to separate the angular and radial dependencies. This shall reduce our three-dimensional problem to a one-dimensional one.

### 3 Spectrum of angular momentum operators

We know that the orbital angular momentum operators are self-adjoint and defined on their Stone domains. They are defined as the generators of a rotation about an axis. We shall instead consider the restrictions of these operators on a common core, the Schwartz space  $\mathcal{S}(\mathbb{R}^3)$ , to avoid running into technical issues.  $\mathcal{S}(\mathbb{R}^3)$  is dense in the Stone domains and  $L^2(\mathbb{R}^3)$ . These operators are essentially self-adjoint on the Schwartz space. This machinery has been explained in the twelfth lecture ‘12: Stone’s Theorem and Construction of Observables’ in the English quantum mechanics series [2, 3].

Let  $i \in \{1, 2, 3\}$ . We thus have the Cartesian angular momentum operators

$$\begin{aligned} L_i &: \mathcal{S}(\mathbb{R}^3) \rightarrow \mathcal{S}(\mathbb{R}^3), \\ L_i &: \psi \mapsto \epsilon_{ijk} Q_j P_k \psi, \end{aligned}$$

where  $Q_j$  and  $P_k$  denote the  $j^{\text{th}}$  and  $k^{\text{th}}$  position and momentum operators respectively. The indices run over 1, 2 and 3. We thus have

$$\begin{aligned} L_1 &= -i\hbar(y\partial_z - z\partial_y), \\ L_2 &= -i\hbar(z\partial_x - x\partial_z), \\ L_3 &= -i\hbar(x\partial_y - y\partial_x). \end{aligned}$$

Consider a Lie algebra over  $\mathbb{C}$  with the basis  $\{J_1, J_2, J_3\}$ , where  $J_i$ 's are linear operators on a *finite-dimensional* Hilbert space and the Lie bracket is given by the commutator. Furthermore, let  $[J_i, J_j] = i\epsilon_{ijk}J_k$  for all  $i, j, k \in \{1, 2, 3\}$ . In the thirteenth lecture ‘13: Spin’ of the English quantum mechanics series [3, 2], we analysed the spectrum of such a system. But it should be noted that without any further justifications, such an analysis *cannot* be done when the vector spaces in question are infinite-dimensional. The Schwartz space, which we need to consider for a spectral analysis of the orbital angular momentum operators, is infinite-dimensional. The justification we need is something Schuller briefly alludes to in the end of the twelfth lecture, but does not elaborate on it any further in the lecture series. We need to consider a number of finite-dimensional subspaces of the Schwartz space, each of which is invariant under the action of the group corresponding to rotations along an axis.<sup>2</sup> Our treatment here is based on an answer of Valter Moretti [4]. The following discussion for this section requires some rudimentary definitions from topology and representation theory; the reader unfamiliar with these is referred to ‘The Geometrical Anatomy of Physics’ lectures of Frederic Schuller [5, 6].<sup>3</sup>

### 3.1 Justification of simultaneous eigenvectors

Consider a Lie algebra over  $\mathbb{C}$  with the basis  $\{J_1, J_2, J_3\}$ , where  $J_i$ 's are linear operators on a Hilbert space  $\mathcal{H}$  and the Lie bracket is given by the commutator. Furthermore, we assume that the following conditions are satisfied.

1. Each  $J_i$  is essentially self-adjoint on a common core.
2.  $\mathbf{J}^2$  is also essentially self-adjoint on the same core, with  $\mathbf{J}^2 := \sum_i J_i \circ J_i$ .
3.  $[J_i, J_j] = i\epsilon_{ijk}J_k$  for all  $i, j, k \in \{1, 2, 3\}$ .

We denote the Lie algebra by  $\mathcal{A} = \text{span}_{\mathbb{C}}\{J_1, J_2, J_3\}$ . We need to show that each  $J_i$  has a pure point spectrum.

We now state a theorem of Edward Nelson, as stated in a book of Moretti [7, p. 291].

---

<sup>2</sup>Let  $D_j(\alpha): \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the operator which describes a counter-clockwise rotation about the  $j^{\text{th}}$  axis by an angle  $\alpha$ . The corresponding group here then is  $\{D_j(\alpha) \mid \alpha \in [0, 2\pi)\}$  with the group operation being executing the rotations about the axis one after other.

<sup>3</sup>More precisely, one needs to know the following notions and concepts. Topological: Hausdorffness, compactness and simple connectedness. Representation theory: Irreducible unitary representations, the Lie groups  $\text{SU}(2)$ ,  $\text{SO}(3)$  and  $\text{SL}(2; \mathbb{C})$ , and the Lie algebras  $\mathfrak{su}(2)$  and  $\mathfrak{sl}(2; \mathbb{C})$ .

**Theorem 1** (Nelson). Consider a real  $n$ -dimensional Lie algebra  $V$  of operators  $-iS_1, \dots, -iS_n$ , where each  $S_i$  with  $i \in \{1, \dots, n\}$  is symmetric on the Hilbert space  $\mathcal{H}$ , defined on a common invariant and dense subspace  $\mathcal{D} \subset \mathcal{H}$ , with the usual commutator as the Lie bracket. Let  $-iS_1, \dots, -iS_n \in V$  be a basis of  $V$  and define Nelson's operator with domain  $\mathcal{D}$  as

$$\Delta_N := \sum_{k=1}^n S_k^2.$$

If  $\Delta_N$  is essentially self-adjoint, then there exists a strongly-continuous unitary representation

$$\begin{aligned} U: G_V &\rightarrow V, \\ U: g &\mapsto U_g \end{aligned}$$

on  $\mathcal{H}$  of the unique simply-connected Lie group  $G_V$  with the Lie algebra  $V$ .  $U$  is uniquely determined by the fact that the closures  $\overline{S}$ , for every  $-iS \in V$ , are the self-adjoint generators of the representations of the one-parameter subgroups of  $G_V$ . In particular, the symmetric operators  $S$  are essentially self-adjoint on  $\mathcal{D}$ .

A proof of the above theorem can be found in the original paper of Nelson [8]. In our case, we see that the conditions of Nelson's theorem are satisfied, with the real Lie algebra  $V = \text{span}_{\mathbb{R}}\{-iJ_1, -iJ_2, -iJ_3\}$ , with the commutator relation

$$[-iJ_i, -iJ_j] = \epsilon_{ijk}(-iJ_k),$$

the Hilbert space being  $L^2(\mathbb{R}^3)$ , and the common domain or core being the Schwartz space  $\mathcal{S}(\mathbb{R}^3)$ .  $V$  is isomorphic to

$$\mathfrak{su}(2) = \text{span}_{\mathbb{R}} \left\{ \frac{i\sigma_1}{2}, -\frac{i\sigma_2}{2}, -\frac{i\sigma_3}{2} \right\}$$

as a Lie algebra, where  $\sigma_i$  denote the Pauli matrices, with the commutator relation

$$\left[ -\frac{i\sigma_i}{2}, -\frac{i\sigma_j}{2} \right] = \epsilon_{ijk} \left( -\frac{i\sigma_k}{2} \right).$$

$\mathfrak{su}(2)$  is the Lie algebra of both  $\text{SU}(2)$  and  $\text{SO}(3)$ . Since  $\mathfrak{su}(2) + i\mathfrak{su}(2) = \mathfrak{sl}(2; \mathbb{C})$ , we have that  $\mathcal{A} = V + iV$ .  $\mathcal{A}$  is isomorphic to  $\mathfrak{sl}(2; \mathbb{C})$  as a Lie algebra.

The closures of  $J_i$ 's are thus generators of a unique strongly continuous representation of  $\text{SU}(2)$  in  $L^2(\mathbb{R})$ . We now state a version of the Peter-Weyl theorem from Moretti [7, p. 287].

**Theorem 2** (Peter–Weyl). Let  $G$  be a compact Hausdorff group, a compact Lie group in particular, and let

$$\begin{aligned} U: G &\rightarrow B(\mathcal{H}), \\ U: g &\mapsto U_g, \end{aligned}$$

be a strongly-continuous unitary representation of  $G$  on the Hilbert space  $\mathcal{H} \neq 0$ , with  $B(\mathcal{H})$  denoting the set of all bounded linear functions on  $\mathcal{H}$ . The following statements then hold.

1. If  $U$  is irreducible, then  $\mathcal{H}$  is finite.
2. If  $U$  is not irreducible, then the orthogonal Hilbert decomposition

$$\mathcal{H} = \bigoplus_{k \in K} \mathcal{H}_k$$

holds, where  $\mathcal{H}_k$  are pairwise orthogonal non-trivial closed subspaces of finite dimension, all invariant under  $U$ .  $K$  is a possibly infinite subset of the natural numbers.

3. Every map  $U|_{\mathcal{H}_k} : \mathcal{H}_k \rightarrow \mathcal{H}_k$  is an irreducible representation of  $G$ .

$\mathrm{SU}(2)$  is a compact group. And thus, strongly-continuous unitary representations of  $\mathrm{SU}(2)$  on  $L^2(\mathbb{R}^3)$  decompose  $L^2(\mathbb{R}^3)$  into finite dimensional subspaces.  $L^2(\mathbb{R}^3)_n$  denotes such a subspace. Thus,

$$L^2(\mathbb{R}^3) = \bigoplus_{n \in \mathbb{N}_0} L^2(\mathbb{R}^3)_n.$$

Define  $\mathcal{S}(\mathbb{R}^3)_n := \mathcal{S}(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)_n$ , for all  $n \in \mathbb{N}_0$ . Thus,

$$\mathcal{S}(\mathbb{R}^3) = \bigoplus_{n \in \mathbb{N}_0} \mathcal{S}(\mathbb{R}^3)_n.$$

Each  $\mathcal{S}(\mathbb{R}^3)_n$  is invariant under the action of  $J_i$ 's. This was the procedure Schuller briefly mentioned. Now, since

$$J_i|_{\mathcal{S}(\mathbb{R}^3)_n} : \mathcal{S}(\mathbb{R}^3)_n \rightarrow \mathcal{S}(\mathbb{R}^3)_n$$

acts on finite dimensional spaces, its spectrum consists of only eigenvalues, i.e. each  $J_i|_{\mathcal{S}(\mathbb{R}^3)_n}$  has a pure-point spectrum. The spectrum of  $J_i$ 's is just the union of these individual eigenvalues, i.e.

$$\mathrm{spec}(J_i) = \bigcup_{n \in \mathbb{N}_0} \mathrm{spec}\left(J_i|_{\mathcal{S}(\mathbb{R}^3)_n}\right).$$

Thus, the usual spectral analysis for angular momentum operators using ladder operators holds, since all  $\mathcal{S}(\mathbb{R}^3)_n$  are finite-dimensional. Thus, using the theorems of Nelson and Peter–Weyl, and the subsequent ladder operator analysis, we get that  $J_i$ ’s have a pure-point spectrum, and that the common eigenvectors of  $\mathbf{J}^2$  and  $J_3$  come as families  $\psi_{j(j+1),m}$ , where  $m = -j, -j+1, \dots, j-1, +j$ . The eigenvalue  $j(j+1)$  is associated to  $\mathbf{J}^2$  and  $m$  is associated to  $J_3$  with  $j \in \mathbb{N}_0/2$ .

Actually, as we only discuss the orbital angular momentum operators, i.e. the  $L_i$ ’s in this lecture, we could have used a slightly different approach to justify the procedure using ladder operators. We know that the spectrum of each  $L_i$  is entirely pure point and contained within the integers, i.e. for all  $i$ ,

$$\text{spec}(L_i) \subseteq \mathbb{Z}.$$

We state a version of the spectral mapping theorem, referring to theorem 3.17 of the book of Teschl [9, p. 118].

**Theorem 3** (Spectral Mapping). Let  $\Xi$  be a self-adjoint operator on a Hilbert space  $\mathcal{H}$ , and let  $f: \mathbb{R} \rightarrow \mathbb{C}$  be a continuous function. Then

$$\text{spec}((f(\Xi))) = f(\text{spec}(\Xi)),$$

if  $|f(\lambda)| \rightarrow \infty$  as  $|\lambda| \rightarrow \infty$ .

Here,  $f(\Xi)$  is defined using the spectral calculus. By applying the spectral mapping theorem, we get that complex polynomials in  $L_i$  have the same spectrum as  $L_i$ , i.e. if  $p$  is a complex polynomial, then

$$\text{spec}(p(L_i)) = p(\text{spec}(L_i)).$$

If we take  $p(x) = x^2$ , then we get that


$$\text{spec}(L_i^2) \subseteq \mathbb{Z},$$

as  $\text{spec}(L_i) \subseteq \mathbb{Z}$ . We now need the following result

**Theorem 4.** If two commuting self-adjoint operators on a complex Hilbert space have a pure point spectrum, then there exists a common orthonormal eigenbasis of both the operators.

This result has been asked as problem 4.5 in the book of Teschl [9, p. 138]. This justifies the assumption of considering simultaneous eigenvectors of  $\mathbf{L}^2 := L_1^2 + L_2^2 + L_3^2$  and  $L_3$  in the usual procedure using ladder operators.





Spherical co-ordinates

### 3.2 Moving to spherical co-ordinates

Let

$$A := [0, +\infty) \times [0, \pi] \times [0, 2\pi).$$

Let  $T$  be defined as

$$\begin{aligned} T: A &\rightarrow \mathbb{R}^3, \\ T: (r, \theta, \phi) &\mapsto (x, y, z), \end{aligned}$$

where

$$\begin{aligned} x &= r \cos \phi \sin \theta, \\ y &= r \sin \phi \sin \theta, \\ z &= r \cos \theta. \end{aligned}$$

Thus,

$$T(r, \theta, \phi) := (r \cos \phi \sin \theta, r \sin \phi \sin \theta, r \cos \theta).$$

If  $F \in \mathcal{S}(A)$  corresponds to  $\psi$  in spherical co-ordinates<sup>4</sup>, we have

$$F = \psi \circ T,$$

and the chain rule gives us

$$F'(r, \theta, \phi) = \psi'(x, y, z) \cdot T'(r, \theta, \phi),$$

where  $(x, y, z) = T(r, \theta, \phi)$ . We thus have

$$\partial_\phi = (\partial_\phi x) \partial_x + (\partial_\phi y) \partial_y + (\partial_\phi z) \partial_z.$$

---

<sup>4</sup> Note that even though  $T$  is only surjective and not injective,  $T^{-1}(\psi)$  is a measure zero set for any  $\psi \in \mathcal{S}(\mathbb{R}^3)$ . Thus, we don't need to make any further identifications in  $\mathcal{S}(A)$ .

Substituting  $x = r \cos \phi \sin \theta$ ,  $y = r \sin \phi \sin \theta$  and  $z = r \cos \theta$ , we get

$$\begin{aligned}\partial_\phi &= -(r \sin \theta \sin \phi) \partial_x + (r \sin \theta \cos \phi) \partial_y \\ \partial_\phi &= x \partial_2 - y \partial_1.\end{aligned}$$

Thus, as expected on physical grounds, we get

$$\tilde{L}_3 = -i\hbar \partial_\phi,$$

where  $\tilde{L}_3: \mathcal{S}(A) \rightarrow \mathcal{S}(A)$  denotes the angular momentum operator in spherical co-ordinates. We now restrict the domain of  $T$  suitably such that it becomes injective and  $T^{-1}$  becomes a well defined function. More precisely, we take the domain to be  $\tilde{A} := (0, +\infty) \times (0, \pi) \times (0, 2\pi)$ . This does not cause any problems as we are excluding only measure zero sets which are irrelevant when considering wave-functions. If we define  $\tilde{\mathbb{R}}^3$  as  $\mathbb{R}^3$  excluding the axes, we have  $T^{-1}: \tilde{\mathbb{R}}^3 \rightarrow \tilde{A}$ ,  $T^{-1}: (x, y, z) \mapsto (r, \theta, \phi)$  as

$$\begin{aligned}r &= \sqrt{x^2 + y^2 + z^2}, \\ \theta &= \cos^{-1} \left( \frac{z}{\sqrt{x^2 + y^2 + z^2}} \right), \\ \phi &= \cos^{-1} \left( \frac{x}{\sqrt{x^2 + y^2}} \right).\end{aligned}$$

Applying chain rule to  $\psi = F \circ T^{-1}$  gives us

$$\partial_i = (\partial_i r) \partial_r + (\partial_i \theta) \partial_\theta + (\partial_i \phi) \partial_\phi.$$

Calculating the partial derivatives and substituting the above in  $L_1 = -i\hbar(y\partial_z - z\partial_y)$  and  $L_2 = -i\hbar(z\partial_x - x\partial_z)$  finally gives us

$$\begin{aligned}\tilde{L}_1 &:= i\hbar \left( \cos \phi \cot \theta \frac{\partial}{\partial \phi} + \sin \phi \frac{\partial}{\partial \theta} \right), \\ \tilde{L}_2 &:= i\hbar \left( -\sin \phi \cot \theta \frac{\partial}{\partial \phi} + \cos \phi \frac{\partial}{\partial \theta} \right), \\ \tilde{L}_3 &:= -i\hbar \frac{\partial}{\partial \phi}.\end{aligned}$$

We also have

$$\tilde{L}^2 F = -\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} F \right) - \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} F \right),$$

and

$$\tilde{H}_{\text{rel}}F = -\frac{\hbar^2}{2\mu} \left( \frac{1}{r} \left( \frac{\partial}{\partial r} \right)^2 (rF) - \frac{\tilde{L}^2}{r^2} F \right) + V(r)F.$$

We now observe that  $\tilde{L}^2 F$  is dependent only on the angular part of  $F$ . So, if we multiply  $F$  by a function  $f(r)$  of  $r$  alone, without any dependencies on  $\theta$  and  $\phi$ , then we have that  $\tilde{L}^2(f \cdot F) = f \cdot \tilde{L}^2 F$ . Because of this observation, we attempt to decompose the Hilbert space into radial and angular parts.

We know that

$$\mathbb{R}^3 = \mathbb{S}^2 \times [0, +\infty)$$

as sets. The standard Lebesgue measure on  $\mathbb{R}^3$  can be seen as the tensor product of the measures on  $\mathbb{S}^2$  and  $[0, \infty)$ . Let  $\mathcal{B}(\mathbb{S}^2)$  be the Borel  $\sigma$ -algebra on  $\mathbb{S}^2$ , generated by the standard induced subset topology on  $\mathbb{S}^2$ , and let  $\mathcal{B}([0, +\infty))$  be the Borel  $\sigma$ -algebra on  $[0, +\infty)$  generated by the standard induced subset topology on  $[0, +\infty)$ . We know that  $\mathbb{S}^2 = [0, \pi] \times [0, 2\pi)$ , as sets. Every element of  $\mathcal{B}(\mathbb{S}^2)$ , i.e. every measurable set of  $\mathbb{S}^2$ , can be expressed as a countable union of sets  $\Theta \times \Phi$ , where  $\Theta := (\theta_1, \theta_2)$  and  $\Phi := (\phi_1, \phi_2)$  are open sets in  $[0, \pi]$  and  $[0, 2\pi)$ . Let

$$\nu: \mathcal{B}(\mathbb{S}^2) \rightarrow \mathbb{R}$$

be the measure defined on  $\mathbb{S}^2$  by

$$\nu((\Theta, \Phi)) := (\phi_2 - \phi_1) \int_{\theta_1}^{\theta_2} \sin \theta \, d\theta.$$

$\nu$  is known as the standard spherical measure on  $\mathbb{S}^2$ . It can be easily seen that  $\nu((\Theta, \Phi))$  is nothing but the area of a surface element on the sphere which subtends the angles  $\Theta$  and  $\Phi$  at the origin in the polar and azimuthal directions respectively. We define the measure

$$\xi: \mathcal{B}([0, +\infty)) \rightarrow \mathbb{R}$$

on  $[0, +\infty)$  by

$$\xi((a, b)) := \int_a^b r^2 dr,$$

with  $a, b \in (0, +\infty)$ . Note that in both the cases, we have defined the measure on open sets. Every element of the Borel  $\sigma$ -algebra can be expressed as a countable union or a countable intersection of the open sets. The standard Lebesgue measure  $\mu$  on  $\mathbb{R}^3$  can be expressed as

$$\mu = \nu \otimes \xi.$$

Applying theorem II.10 of the book of Reed and Simon [1, p. 52] again, we see that

$$L^2(\mathbb{R}^3, \mu) = L^2(\mathbb{S}^2, \nu) \otimes L^2([0, +\infty), \xi). \quad (1)$$

We define the operators  $\tilde{L}_{\mathbb{S}^2}^2 : C^\infty(\mathbb{S}^2, \nu) \rightarrow C^\infty(\mathbb{S}^2, \nu)$  and  $\tilde{L}_{3, \mathbb{S}^2} : C^\infty(\mathbb{S}^2, \nu) \rightarrow C^\infty(\mathbb{S}^2, \nu)$ . These are simply the angular parts of the  $\tilde{L}^2$  and  $\tilde{L}_3$  operators, which anyway act on  $C^\infty(\mathbb{S}^2, \nu)$  since they did not have any radial dependence.

$$\begin{aligned} \tilde{L}_{\mathbb{S}^2}^2 &= -i\hbar \partial_\phi, \\ \tilde{L}_{3, \mathbb{S}^2} &= -\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) - \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \right). \end{aligned}$$

We use the fact that  $C^\infty(\mathbb{S}^2, \nu)$  is dense in  $L^2(\mathbb{S}^2, \nu)$ . It can be seen that these operators are symmetric.

We need the fact that  $\tilde{L}^2$  is essentially self-adjoint on  $C^\infty(\mathbb{S}^2)$  [10]. In order to fully determine  $\text{spec}(H_{\text{rel}})$ , along with the continuous spectrum, i.e. the part containing the generalised eigenvalues, we would need to lift the eigenvalue equation to the distribution space and solve it.<sup>5</sup> However, we already know from experiments that the hydrogen atom has discrete lines in its observed spectrum. Thus, we shall first try to find the point spectrum

$$\text{spec}_p(H_{\text{rel}}) = \{E \in \mathbb{R} \mid H_{\text{rel}}F = EF\},$$

assuming it exists, which was historically a major validation for the quantum theory.

### 3.3 Spherical harmonics

$\mathcal{S}(\mathbb{R}^3)$ ,  $\mathcal{S}(\tilde{\mathbb{R}}^3)$ ,  $\mathcal{S}(A)$ , and  $\mathcal{S}(\tilde{A})$  are all isomorphic to each other. Thus, the eigenvalues and eigenvectors of  $L^2$  and  $L_i$ 's, and  $\tilde{L}^2$  and  $\tilde{L}_i$ 's are the same, respectively. The self-adjointness is unaffected by these co-ordinate changes as well. Thus, we know that the common eigenvectors of  $\tilde{L}^2$  and  $\tilde{L}_3$  come as families  $\psi_l^m \in \mathcal{S}(\tilde{A})$ , where  $m = -l, -l+1, \dots, l-1, l$  and  $l \in \mathbb{N}_0$ . We have

$$\begin{aligned} \tilde{L}^2 \psi_l^m &= l(l+1) \psi_l^m, \\ \tilde{L}_3 \psi_l^m &= m \psi_l^m. \end{aligned}$$

---

<sup>5</sup> This can in principle be achieved using Gelfand's rigged Hilbert space formalism, which was introduced in an earlier lecture. The lecture itself can be found on this link: [https://www.youtube.com/watch?v=FNJ0yxOp3Ik&list=PLP05pgr\\_frzTeqa\\_thbltYjyw8F9ehw7v&index=8](https://www.youtube.com/watch?v=FNJ0yxOp3Ik&list=PLP05pgr_frzTeqa_thbltYjyw8F9ehw7v&index=8). The notes for this lecture can be found here: [https://github.com/apoorvpotnis/schuller\\_rigged\\_hilbert\\_spaces/blob/main/schuller\\_rigged\\_hilbert\\_spaces.pdf](https://github.com/apoorvpotnis/schuller_rigged_hilbert_spaces/blob/main/schuller_rigged_hilbert_spaces.pdf).

Note that  $l$  can take only integer values, not half-integer values as we are dealing with the orbital angular momentum operator. In spherical co-ordinates, it turns out that the solutions  $\psi_l^m$  are given by

$$\psi_l^m(\tilde{\mathbf{x}}(r, \theta, \phi)) = k \cdot Y_l^m(\theta, \phi) \cdot f(r),$$

where

$$Y_l^m(\theta, \phi) := \frac{(-1)^m}{2^l l!} \left( \frac{(2l+1)(l-m)!}{4\pi(l+m)!} \right)^{\frac{1}{2}} \cdot e^{im\phi} \cdot (\sin \theta)^m \left( \frac{\partial}{\partial \cos \theta} \right)^{l+m} (\cos^2 \theta - 1)^l$$

and  $\tilde{\mathbf{x}} = T^{-1}(\mathbf{x})$ . These  $Y_l^m$ 's are called as *spherical harmonic functions*. Since the operators  $\tilde{L}^2$  and  $\tilde{L}_3$  act only on the angular components and not the radial components when expressed in spherical co-ordinates, we have that

$$\begin{aligned} \tilde{L}^2 Y_l^m &= l(l+1) Y_l^m, \\ \tilde{L}_3 Y_l^m &= m Y_l^m, \end{aligned}$$

i.e. the spherical harmonic functions are eigenvectors of  $\tilde{L}^2$  and  $\tilde{L}_3$ . These functions actually form an orthonormal basis of the Hilbert space  $L^2(\mathbb{S}^2)$  consisting of complex square-integrable functions defined on the 2-sphere.

$$\langle Y_l^m, Y_{l'}^{m'} \rangle = \delta_{l,l'} \cdot \delta_{m,m'}.$$

We request that the reader consult section 14.2 of the book of Bowers for a derivation of these spherical harmonic functions [11, p. 200].

## 4 Spectrum of the Hamiltonian

Depending on  $H_{\text{rel}}$ , a common orthonormal eigenbasis of  $L^2$ ,  $L_i$ , and  $H_{\text{rel}}$  can possibly exist, as all three operators commute with each other pairwise. We for the moment *assume* that this holds.<sup>6</sup>

## References

- [1] Michael Reed and Barry Simon. *Methods of Modern Mathematical Analysis I: Functional Analysis*. Revised and Enlarged editon. Vol. 1. Academic Press, Inc. London, 1980. ISBN: 978-0-080-57048-8 (cit. on pp. 3, 12).

---

<sup>6</sup>Clarify this later.

- [2] Frederic Schuller. *Lectures on Quantum Theory*. Video lectures on YouTube. 2015. URL: [https://youtube.com/playlist?list=PLPH7f\\_7ZlzxQVx5jRjbfrGEzWY\\_upS5K6](https://youtube.com/playlist?list=PLPH7f_7ZlzxQVx5jRjbfrGEzWY_upS5K6) (cit. on pp. 3–5).
- [3] Frederic Schuller, Simon Rea, and Richie Dadhley. ‘Lectures on Quantum Theory’. Lecture notes in .pdf format. Lecturer: Prof. Frederic Paul Schuller. 2019. URL: [https://docs.wixstatic.com/ugd/6b203f\\_a94140db21404ae69fd8b367d9fcd360.pdf](https://docs.wixstatic.com/ugd/6b203f_a94140db21404ae69fd8b367d9fcd360.pdf) (cit. on pp. 3–5).
- [4] Valter Moretti (<https://physics.stackexchange.com/users/35354/valter-moretti>). *Answer to the question ‘Angular momentum Lie algebra for infinite-dimensional Hilbert spaces’*. Physics Stack Exchange (<https://physics.stackexchange.com>). (version: 2024-05-17 21:03:48Z). URL: <https://physics.stackexchange.com/a/814882/81224> (cit. on p. 5).
- [5] Frederic Schuller. *Lectures on the Geometric Anatomy of Theoretical Physics*. Video lectures on YouTube. 2016. URL: [https://www.youtube.com/playlist?list=PLPH7f\\_7ZlzxTi6kS4vCmv4ZKm9u8g5yic](https://www.youtube.com/playlist?list=PLPH7f_7ZlzxTi6kS4vCmv4ZKm9u8g5yic) (cit. on p. 5).
- [6] Frederic Schuller, Simon Rea, and Richie Dadhley. *Lectures on the Geometric Anatomy of Theoretical Physics*. Lecture notes in .pdf format. Lecturer: Prof. Frederic Paul Schuller. 2017. URL: <https://drive.google.com/file/d/1nchF1fRGSY3R3rP1QmjUg7fe28tAS428/view> (cit. on p. 5).
- [7] Valter Moretti. *Fundamental Mathematical Structures of Quantum Theory: Spectral Theory, Foundational Issues, Symmetries, Algebraic Formulation*. Springer Nature Switzerland AG, 20189. ISBN: 978-3-030-18345-5 (cit. on pp. 5, 6).
- [8] Edward Nelson. *Analytic Vectors*. Vol. 70. 3. Annals of Mathematics, Trustees of Princeton University on Behalf of the Annals of Mathematics, Mathematics Department, Princeton University, 1959, pp. 572–615. DOI: <https://doi.org/10.2307/1970331> (cit. on p. 6).
- [9] Gerald Teschl. *Mathematical methods in quantum mechanics: With applications to Schrödinger operators*. 2nd ed. The book is freely available in .pdf format on the author’s website at <https://www.mat.univie.ac.at/~gerald/ftp/book-schroe/schroe.pdf>. American Mathematical Society, Providence, Rhode Island, 2014. ISBN: 978-1-4704-1704-8 (cit. on p. 8).
- [10] Valter Moretti (<https://physics.stackexchange.com/users/35354/valter-moretti>). *Answer to the question ‘What are the proper domains of the position and squared angular momentum operator?’* Physics Stack Exchange (<https://physics.stackexchange.com>). (version: 2014-07-23 00:40:35Z).

URL: <https://physics.stackexchange.com/a/127629/81224> (cit. on p. 12).

- [11] Philip Bowers. *Lectures on Quantum Mechanics*. Cambridge University Press, Cambridge, 2020. ISBN: 978-1-108-42976-4 (cit. on p. 13).
- [12] Valter Moretti. *Spectral Theory and Quantum Mechanics: Mathematical Foundations of Quantum Theories, Symmetries and Introduction to the Algebraic Formulation*. 2nd ed. Springer Nature Switzerland AG, 2018. ISBN: 978-3-319-70705-1. Translated from Italian by Simon G. Choissi.
- [13] Eduard Prugovečki. *Quantum Mechanics in Hilbert Space*. 2nd ed. Academic Press, Inc. London, 1981. ISBN: 0-12-566060-X.
- [14] Brian C. Hall. *Quantum Theory for Mathematicians*. Springer Science+Business Media New York, 2013. ISBN: 978-1-4614-7115-8.