


1 **Supplementary Figure 1**



2

3 **Supplementary Figure 1. Donor bootstrap uncertainty in genetic candidate expression. Points**

4 **show mean log2(CP10k + 1) across eligible CIDP donors and bars show percentile 95%**

5 **intervals from 5000 resamples within each gene–cell group. These are descriptive intervals for**

6 **expression means, not tests of cell-type specificity or genetic mediation.**

7 **Alt text: Four expression plots show gene-specific mean expression and donor-bootstrap**


8 **intervals across nerve cell groups.**

9

10 **Supplementary Figure 2**

Published CIDP nerve evidence

52/55 biopsies: 94.5% (Wilson 95% CI 85.1–98.1%)



Additional source observations

Increased C3 and C6 abundance in nerve proteomics

CD68-positive macrophages and CD8-positive T cells

Source: Stascheit et al. 2025; DOI: 10.1007/s00401-025-02936-w

11

12 Supplementary Figure 2. Published nerve observations and a calculated complement-

13 deposition interval. Stascheit et al. reported C5b-9 deposition on endoneurial capillaries in 52

14 of 55 CIDP biopsies and additional histological and proteomic observations.³⁷ The proportion

15 (94.5%) and Wilson 95% interval (85.1–98.1%) were calculated from that published count.

16 The other observations are qualitative source findings associated with PXD056286; no biopsy


17 or proteome was reanalysed.

18 Alt text: A point and interval summarize published C5b-9 deposition in 52 of 55 biopsies; text

19 lists additional source observations.


20

21 Supplementary Figure 3



23 Supplementary Figure 3. Fc module components and donor sensitivity analyses. Panel A
 24 shows differences in mean $\log_2(\text{CPM} + 0.5)$ for the 11 measured genes. FCGR2C is
 25 unavailable. Displayed q values use the 11 gene family. Panel B shows module score
 26 differences under demographic and composition sensitivity analyses. Points and bars are
 27 mean differences and 95% intervals from ordinary least squares with HC3 covariance and
 28 residual degrees of freedom. Standardisation in 20 donors changes the score scale. The exact
 29 permutation test within centres has $P = 0.0048$ and is reported separately because it does not
 30 supply a model interval. The original module q value is 0.0087 within eight macrophage
 31 modules and 0.087 across all 80 eligible module tests. These post hoc analyses are not
 32 independent validation.
 33 Alt text: Gene effects vary in direction. Demographic model intervals exclude zero. The
 34 composition adjusted interval includes zero.

36 Supplementary Figure 4



37

38 Supplementary Figure 4. Fc scores under alternative annotations. The original and refined
 39 annotations are distinct classification levels. The refined union contains Macro1 through
 40 Macro18. Eligibility requires 20 nuclei per donor and three donors per diagnosis. Original
 41 Macro1 is ineligible. Points and bars show mean score differences and HC3 95% intervals.
 42 Genes are standardized within each eligible annotation. Parentheses give CIDP and CIAP
 43 donor counts. Subtype q values use nine tests. None meets $q < 0.05$. The three broader
 44 definition checks form a separate family.

45 Alt text: Broader macrophage definitions retain negative differences. Individual refined
 46 subtype estimates have limited donor counts, and none survives correction across nine
 47 subtypes.