

1 **Table 3. Descriptive evidence map across blood, CSF and peripheral nerve**

Program	Blood evidence	CSF evidence	Peripheral nerve evidence	Interpretation
CXCL8 recruitment	Positive in 3 cohorts; pooled P = 0.114; whole-blood q < 0.05	Published innate immune and inflammatory protein findings	Myeloid localization and endothelial trafficking context	Shared recruitment hypothesis
Complement	Positive in 3 cohorts; pooled P = 0.234; whole-blood q < 0.01	Published complement protein findings	Macrophage localization	Complement-related evidence; amplification remains a hypothesis
Fc receptor	Whole blood q < 0.05, with mixed directions across cohorts	Not directly evaluated	Macrophage q = 0.0087 within eight modules and q = 0.087 across 80 nerve comparisons	Exploratory association. Attenuated after composition adjustment.
Interferon response	Whole blood q < 0.05, with mixed directions across cohorts	Published evidence of antiviral defense	Macrophage, endothelial and Schwann cell states	Response varies by compartment
Endothelial adhesion	Positive blood direction	Not directly evaluated	ICAM1 and VCAM1 localization to BNB endothelial cells	Endothelial context for functional study
Schwann repair	Not evaluated	Published evidence of axonal domain peptide degradation	Repair and stress programs localized to Schwann cells	Proposed link between structural injury and repair

- 2 q denotes the Benjamini–Hochberg-adjusted P value within the specified testing family. Blood comparisons concern acute GBS versus controls;
3 the nerve comparison concerns CIDP versus CIAP. This map does not constitute a direct paired comparison between compartments or a test of
4 causal mediation.