

Boosting Variational Inference: An Optimization Perspective


Francesco Locatello^{1,2}, Rajiv Khanna³, Joydeep Ghosh³, Gunnar Rätsch¹



Variational Inference & Boosting

Variational Inference

$$\min_{q \in \mathcal{Q}} D^{KL}(q || p_x)$$



Boosting Variational Inference

$$\min_{q \in \text{conv}(\mathcal{Q})} D^{KL}(q || p_x)$$


- Efficient approximate inference
- Poor approximation of the posterior
- Can't trade time with accuracy

- Efficient approximate inference
- Highly expressive family
- Greedily adding components: convex optimization problem

Frank-Wolfe [Frank&Wolfe 1956]

Linearization

$$\min_{\mathbf{z} \in \mathcal{A}} f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{z} - \mathbf{x} \rangle$$



$\text{conv}(\mathcal{A})$

[Jaggi]


Frank-Wolfe [Frank&Wolfe 1956]

Linearization

$$\min_{\mathbf{z} \in \mathcal{A}} f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{z} - \mathbf{x} \rangle$$

The Linear Minimization Oracle

$$\text{LMO}_{\mathcal{A}}(\mathbf{d}) := \arg \min_{\mathbf{z} \in \mathcal{A}} \langle \mathbf{d}, \mathbf{z} \rangle$$



$\text{conv}(\mathcal{A})$

[Jaggi]

Frank-Wolfe [Frank&Wolfe 1956]


Linearization

$$\min_{\mathbf{z} \in \mathcal{A}} f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{z} - \mathbf{x} \rangle$$

The Linear Minimization Oracle

$$\text{LMO}_{\mathcal{A}}(\mathbf{d}) := \arg \min_{\mathbf{z} \in \mathcal{A}} \langle \mathbf{d}, \mathbf{z} \rangle$$

-
- 1: **init** $\mathbf{x}_0 \in \text{conv}(\mathcal{A})$
 - 2: **for** $t = 0 \dots T$
 - 3: Find $\mathbf{z}_t := (\text{Approx-}) \text{LMO}_{\mathcal{A}}(\nabla f(\mathbf{x}_t))$
 - 4: Variant 0: $\gamma = \frac{2}{t+2}$
 - 5: Update $\mathbf{x}_{t+1} := (1 - \gamma) \mathbf{x}_t + \gamma \mathbf{z}_t$
 - 6: **end for**




[Jaggi]

Frank-Wolfe [Frank&Wolfe 1956]

$$\min_{\mathbf{x} \in \text{conv}(\mathcal{A})} f(\mathbf{x})$$

```
1: init  $\mathbf{x}_0 \in \text{conv}(\mathcal{A})$ 
2: for  $t = 0 \dots T$ 
3:   Find  $\mathbf{z}_t := (\text{Approx-}) \text{LMO}_{\mathcal{A}}(\nabla f(\mathbf{x}_t))$ 
4:   Variant 0:  $\gamma = \frac{2}{t+2}$ 
5:   Update  $\mathbf{x}_{t+1} := (1 - \gamma)\mathbf{x}_t + \gamma\mathbf{z}_t$ 
6: end for
```



Frank-Wolfe [Frank&Wolfe 1956]

$$\min_{\mathbf{x} \in \text{conv}(\mathcal{A})} f(\mathbf{x})$$

```


1: init  $\mathbf{x}_0 \in \text{conv}(\mathcal{A})$ 
2: for  $t = 0 \dots T$ 
3:   Find  $\mathbf{z}_t := (\text{Approx-}) \text{LMO}_{\mathcal{A}}(\nabla f(\mathbf{x}_t))$ 
4:   Variant 0:  $\gamma = \frac{2}{t+2}$ 
5:   Update  $\mathbf{x}_{t+1} := (1 - \gamma)\mathbf{x}_t + \gamma\mathbf{z}_t$ 
6: end for

```

Sparse Iterate

$$\mathbf{x}_t = \sum_{i=0}^t \gamma_i \mathbf{z}_i$$

$$\sum_{i=0}^t \gamma_i = 1$$



Outlook

- ❖ How to adapt the FW framework to boost variational inference
- ❖ Converge analysis in the Variational Inference case

Thank you

- ✿ Poster #31
- ✿ P.S. We are hiring PostDocs!

