# Natural Gradients via the Variational Predictive Distribution

Da Tang<sup>1</sup> and Rajesh Ranganath<sup>2</sup>


Columbia University<sup>1</sup> New York University<sup>2</sup>

December 8, 2017



## Pathological Curvature of the ELBO

- The curvature of the ELBO may be pathological
- Example: A bivariate Gaussian model with unknown mean and known covariance  $\Sigma=\begin{pmatrix}1&1-arepsilon\\1-arepsilon&1\end{pmatrix}$ ,  $0<arepsilon\ll1$



• The natural gradient (Hoffman et al., 2013) fails to help



#### The Variational Predictive Natural Gradient

• Approximate the posterior predictive distribution:


$$r(\mathbf{x}'|\mathbf{x}_i; \boldsymbol{\lambda}) = \int p(\mathbf{x}'|\mathbf{z}_i, \boldsymbol{\beta}) q(\mathbf{z}_i|\mathbf{x}_i, \boldsymbol{\beta}; \boldsymbol{\lambda}) q(\boldsymbol{\beta}; \boldsymbol{\lambda}) d\mathbf{z}_i d\boldsymbol{\beta}$$

• The variational predictive Fisher information:

$$F_r = \mathbb{E}_{Q_{\mathbf{x}_i}, r(\mathbf{x}'|\mathbf{x}_i; \boldsymbol{\lambda})} [\nabla_{\boldsymbol{\lambda}} \log r(\mathbf{x}'|\mathbf{x}_i; \boldsymbol{\lambda}) \cdot \nabla_{\boldsymbol{\lambda}} \log r(\mathbf{x}'|\mathbf{x}_i; \boldsymbol{\lambda})^{\top}]$$

Eigenspace comparison:








- (a) Precision mat  $\Sigma^{-1}$
- (b) *q*-Fisher info  $F_q$  (c) Our Fisher info  $F_r$

### Experiment: Learning a VAE

- Training on the MNIST dataset (Lecun et al., 1998)
- Learning curves:



# Thank you!