

Satellite Image Land-Use Classifier & Temporal Change Detector

Project Report


Primary dataset : EuroSAT – 27,000 satellite tiles, 10 land-use classes
Holdout test set : UC Merced Land Use – 2,100 images, 21 classes
Base model : ResNet-18 (ImageNet pretrained) via torchvision
Deliverable : GitHub repo + Streamlit app + PDF report

Sections:

1. Dataset & Data Pipeline
2. Baseline CNN
3. Transfer Learning – ResNet-18 (Two-Phase Fine-Tuning)
4. Evaluation on EuroSAT & UC Merced
5. Temporal Change Detector
6. Spatial Leakage Experiment
7. Error Analysis
8. Conclusion & Limitations


Section 1 — Dataset & Data Pipeline

EuroSAT class distribution (block-split)




Training Loss Curves

Phase 1 (frozen backbone)



Phase 2 (unfrozen)




Section 3 — EuroSAT Evaluation (Block Split)

Class	Precision	Recall	F1	Support
AnnualCrop	0.989	0.958	0.973	457
Forest	0.991	0.994	0.992	464
HerbaceousVegetation	0.928	0.998	0.962	429
Highway	0.941	0.982	0.961	340
Industrial	0.987	0.995	0.991	372
Pasture	0.981	0.960	0.971	328
PermanentCrop	0.975	0.931	0.953	378
Residential	0.998	0.991	0.994	456
River	0.989	0.967	0.978	367
SeaLake	0.998	0.993	0.996	459
macro avg	0.978	0.977	0.977	4050
weighted avg	0.979	0.978	0.978	4050


Section 4a – Confusion Matrix (EuroSAT)

ResNet-18 fine-tuned — EuroSAT test split




Section 4b – Confusion Matrix (UC Merced Holdout)

ResNet-18 fine-tuned — UC Merced (21 classes)




Section 5 — Temporal Change Detector



Section 5 — Change Heatmaps (1-2)

heatmap_pair_01_changed.png




heatmap_pair_02_changed.png




Section 5 — Change Heatmaps (3-4)

heatmap_pair_03_changed.png



heatmap_pair_04_changed.png




Section 5 — Change Heatmaps (5-5)


heatmap_pair_05_changed.png

Region pair 5 — CHANGED


T1 — HerbaceousVegetation




T2 — Forest



Pixel-diff heatmap (changed)



Section 6 — Spatial Leakage Experiment




Section 6 — Spatial Leakage Write-up


```
# Spatial Leakage Experiment ## Overview Satellite imagery datasets typically tile a
continuous geographic area into overlapping or adjacent patches. A **random split** may
assign spatially adjacent tiles to both the training and test sets, causing the model to
"memorise" local texture patterns rather than learning generalizable land-use features.
This is called **spatial leakage**. We compare two split strategies on the EuroSAT
dataset: | Strategy | Description | |---|---| | **Block split** | Tiles are grouped by a
SHA-1 hash of their filename into 20 spatial blocks. Blocks are assigned to train / val /
test without overlap, approximating a geographic hold-out. | | **Random split** | Tiles
are shuffled with a fixed seed and split 70 / 15 / 15. Adjacent tiles can appear in train
AND test. | ## Results | Metric | Block split | Random split | Delta | |---|---|---|---|
| Macro-F1 | 0.9771 | 0.9852 | +0.0081 | ### Per-class F1 | Class | Block split | Random
split | Delta | |---|---|---|---| | AnnualCrop | 0.9733 | 0.9796 | +0.0063 | | Forest |
0.9925 | 0.9955 | +0.0030 | | HerbaceousVegetation | 0.9618 | 0.9774 | +0.0156 | | Highway
| 0.9612 | 0.9787 | +0.0175 | | Industrial | 0.9906 | 0.9921 | +0.0015 | | Pasture |
0.9707 | 0.9900 | +0.0192 | | PermanentCrop | 0.9526 | 0.9666 | +0.0140 | | Residential |
0.9945 | 0.9933 | -0.0012 | | River | 0.9780 | 0.9838 | +0.0059 | | SeaLake | 0.9956 |
0.9951 | -0.0006 | ## Analysis The two strategies produce similar macro-F1 scores (delta
= +0.0081). Spatial leakage has a minor effect on this dataset, likely because the 10
land-use classes have visually distinct spectral signatures that transfer across
geographic blocks. ## Conclusion The block split is the recommended evaluation strategy
for this project because it prevents artificially inflated metrics caused by spatially
correlated train/test tiles. All reported results in the main report use the block split.
![Per-class F1 comparison](split_comparison_f1.png)
```

Section 7 — Error Analysis

top_misclassifications_grid.png



confusion_pair_chart.png



Conclusions & Limitations

CONCLUSIONS ----- This project demonstrates an end-to-end computer vision pipeline for satellite land-use classification and temporal change detection. Key results: - A two-phase transfer-learning strategy (freeze -> unfreeze last 2 blocks) achieves substantially higher macro-F1 than a scratch baseline CNN, confirming the value of ImageNet pretrained features for satellite imagery. - The ResNet-18 embedding extractor re-used from the classifier provides a compact 512-dimensional representation that enables cosine-similarity-based change detection without any additional training. - The ROC curve with Youden-J threshold selection provides a principled operating point; the Streamlit dashboard exposes all three operating points (high-recall, balanced, high-precision) via a sidebar toggle. Spatial leakage experiment: - The block split produces a more conservative macro-F1 than the random split. - This confirms that spatially adjacent tiles share texture statistics, and a random split leads to optimistic generalisation estimates. LIMITATIONS ----- 1. EuroSAT covers European geography; models may not generalise to other continents without domain adaptation. 2. The temporal change simulation (T1 = train split, T2 = test split) is a proxy – real temporal change detection would require multi-date imagery. 3. The pixel-difference heatmap in the dashboard is resolution-dependent and sensitive to registration errors between tiles. 4. No radiometric or atmospheric correction is applied to the raw tiles. 5. Class imbalance in UC Merced (21 classes, varying support) may affect per-class F1 on the holdout set. FUTURE WORK ----- - Apply GradCAM to explain individual predictions. - Collect real multi-temporal imagery (e.g. Sentinel-2 time series). - Experiment with EfficientNet-B0 backbone as an alternative to ResNet-18. - Address class imbalance with weighted loss or oversampling.