

Calculus Notes

Jack Gong
Pinetree Secondary
`haotiangong@hotmail.com`

November 9, 2019

Contents

1	Intro	3
1.1	Ideas about Sets	3
1.2	Sets We Use Frequently	3
1.3	More definitions	3
1.4	Properties of Reals	4
1.5	The Least Upper Bound Property	4
1.5.1	Least Upper Bound and \mathbb{Q}	5
1.6	Proof by Contradiction	5
1.7	Open and Closed Intervals	6
1.8	Function Definitions	6
1.8.1	Graphical Representation	7
1.9	Injectivity and Surjectivity	8
1.10	Function interactions	9
1.11	Triangle inequality	9
1.12	Polynomial	9
1.13	Rational Function	10
2	Limits	10
2.1	Definition	10
2.1.1	Negating statements involving quantifiers	11
2.1.2	Proving limits false	12
2.2	Limit Rules	12
2.3	Squeeze Theorem	16
2.4	One-sided Limits	17
2.5	Limits at Infinity	18
2.6	Infinite Limits	19
2.7	Existence of Limits	20
2.7.1	Use of product rule	20
2.8	Asymtotes	20
2.9	Limit dependency on Power of Polynomial	22
2.10	Limit Laws	25
2.11	Limit Calculations	26
2.11.1	Direct Substitution Property	26
3	Continuity	27

1 Intro

1.1 Ideas about Sets

Definition of a set: a collection of distinct objects. These objects are called the elements of the set.

We use $\{\}$ to enclose elements of the set. Two sets are the same if they contain the same objects. The size, or cardinality of a set is the number of (distinct) elements in that set. It may be infinite.

Sets can be “arbitrarily declared” without following the rule of distinct elements. For example,

$$\{1, 1, 2, 2, 3, 3, 4, 4, 5, 5\} = \{1, 2, 3, 4, 5\}$$

1.2 Sets We Use Frequently

- $\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$
- $\mathbb{Z} = \{-2, -1, 0, 1, 2, \dots\}$
- \mathbb{Q} How do we define rational numbers?
- \mathbb{R} What about real numbers?

Trying to define rationals

1st attempt: Set of all numbers that can be expressed in the form $\frac{a}{b}$, where a and b are integers and $b \neq 0$.

But how do we define “numbers”?

2nd attempt (to avoid explaining the definition): Define S to be the set of all pairs of integers (a, b) , where $b \neq 0$; and declare that the pair (a, b) is the same as the pair (a', b') if $ab' = a'b$.

\mathbb{Q} is the set of results of $\frac{a}{b}$.

1.3 More definitions

If S & T are sets, their union is the set consisting of all elements that are elements of S or are elements of T , or both. We write this as: $S \cup T$.

Different notation:

$$S \cup T = \{x : x \in S \text{ or } x \in T\}$$

If S and T are sets, the intersection of S and T is the set consisting of all elements that are in S and T .

$$S \cap T = \{x : x \in S \text{ and } x \in T\}$$

If S and T are sets and every element of S is also an element of T , then we say T contains S and write this as

$$S \subset T \text{ or } T \supset S$$

If S and T are sets, we also define the set difference operation:

$$S \setminus T = \{x : x \in S \text{ and } x \notin T\}$$

How do we express the elements in $\{x : x \in S \text{ or } x \in T, \text{ but not both}\}$

We can express it as $(S \setminus T) \cup (T \setminus S)$ or $(S \cup T) \setminus (S \cap T)$. (It can also be expressed as $S \triangle T$)

How do we prove that these sets actually equal to each other? We can prove that \subset and \supset goes both ways. If a set includes the other and the other includes the set, they are equal.

Empty (null) set: $\{\} = \emptyset$

If S and T are sets and $S \cap T = \emptyset$, we say S and T are **disjoint**.

1.4 Properties of Reals

Defining the reals is a tedious task. Rather than defining the real numbers, we will describe some of their properties.

- \mathbb{R} is a set. It contains the national numbers.
- They can be added and subtracted.
- They can be multiplied and divided.
- The real numbers are ordered. If $x, y \in \mathbb{R}$, then either $x < y$, $x = y$, or $x > y$.
- Product of 2 positives or negatives is positive; product of positive and negative is negative.
- The \mathbb{R} s follow the **Least upper bound property**.

1.5 The Least Upper Bound Property

If $S \subset \mathbb{R}$, a real number \mathbb{R} is called an upper bound for S if:

$$\text{For all } (\forall)y \in S, \text{ we have } y \leq x.$$

Is the upper bound of $T = \{x \in \mathbb{R}, x < 0\}$ in T ?

Prove: For every $x \in T$, there exists a number $y \in T$ with $y > x$.

Proof: Let $y = \frac{x}{2}$. We need to show ($y \in T$ and $y > x$)

① Since $\frac{1}{2}$ is positive, $y = \frac{1}{2}x$ has the same sign as x .
 Since $x < 0$, we conclude $y < 0$. Hence $y \in T$.

② Since $x < 0$, and $0 < \frac{1}{2} < 1$, we have $y = \frac{x}{2} > x$.

We conclude if $x \in T$, then x is not an upper bound for T .

Some sets $S \subset \mathbb{R}$ have an upper bound, some don't.

If $S \subset \mathbb{R}$ has an upper bound, we say it is **bounded above**; otherwise it is not bounded above.

Let $S \subset \mathbb{R}$. A number $x \in \mathbb{R}$ is called a least upper bound for S if

1. x is an upper bound for S .
2. If $z \in \mathbb{R}, z < x$, then z is not an upper bound for S .


Least upper bound property: if $S \subset \mathbb{R}$ and S is bounded above, then S has a least upper bound.

1.5.1 Least Upper Bound and \mathbb{Q}

How do we show the least upper bound prop. isn't true for all \mathbb{Q} s?

Thought: construct a set whose least upper bound is a real number.

Let $S = \{x \in \mathbb{Q}, x > 0, \text{ and } x^2 < 2\}$



We will show:

1. S is bounded above. Indeed, if $x \in \mathbb{Q}, x^2 > 2$, then x is an upper bound and $x > 0$.
2. If $y \in \mathbb{Q}$ is an upper bound for S , then there exists $z \in \mathbb{Q}$, with $z < y$ and z is also an upper bound.


1.6 Proof by Contradiction

The goal is to prove some mathematical statement.

1. Suppose the statement is false.
2. Using the assumption that the statement is false, and its relevant deductions,
3. arrive at the conclusion it is clearly false.
4. Conclude the original statement is false.

1.7 Open and Closed Intervals

We can think of real numbers as points on a number line.



An **open interval** is a subset of \mathbb{R} of one of the following forms:

$$1. \begin{cases} x \in \mathbb{R}, x > a \text{ and } x < b, \text{ where } a, b \in \mathbb{R}, a < b \\ x \in \mathbb{R} : a < x < b \\ (a, b) \end{cases}$$

We have finite length open intervals with length $b - a$.

$$2. \begin{cases} x \in \mathbb{R} : x > a, \text{ where } a \in \mathbb{R} \\ (a, \infty) \end{cases}$$

$$3. \begin{cases} x \in \mathbb{R} : x < b, \text{ where } b \in \mathbb{R} \\ (-\infty, b) \end{cases}$$

$$4. \begin{cases} x \in \mathbb{R} \\ (-\infty, \infty) \end{cases}$$

A **closed interval** is a subset of \mathbb{R} of one of the following forms:

$$1. \begin{cases} x \in \mathbb{R} : a \leq x \leq b \text{ } a, b \in \mathbb{R}, a < b \\ [a, b] \end{cases}$$

$$2. \begin{cases} x \in \mathbb{R} : a \leq x \\ [a, \infty] \end{cases}$$

$$3. \begin{cases} x \in \mathbb{R} : x \leq b \\ [-\infty, b] \end{cases}$$

$$4. \mathbb{R}$$

What about the sets $(a, b]$ or $[a, b)$? “Half-open” interval.

1.8 Function Definitions

A function is a rule f that assigns to each element x of a set D , called the domain, a unique element $f(x)$ of a set S , called the co-domain.

The domain and co-domain can be the same set.

We write:

$$x \in D, f(x) \in S, \text{ and } f : D \rightarrow S$$

Example: $D : \{1, 2, 3, 4\}$, $S \subset \mathbb{R}$, and f is given by $f(1) = 1, f(2) = 1, f(3) = \pi, f(4) = -3$

- $D = \mathbb{R}, f(x) = 0, f(x) = x, f(x) = x^2 + x + 1$ - **Polynomials**
- $D = \mathbb{R}, f(x) = \sin x, f(x) = \cos x, f(x) = \sin(2x)$ - **Trigonometric functions**
- $D = \mathbb{R}, f(x) = e^x, f(x) = e^2 x$ - **Exponential functions**
- $D = (0, \infty), f(x) = \log x, f(x) = \frac{1}{2} \log x$ - **Logarithmic functions**

We choose to study these special cases of functions because most mathematical aspects of the natural world are reflected by them.

Domain convention : If the domain of a function isn't specified, we assume the domain is the largest subset of \mathbb{R} for which $f(x)$ "makes sense" as a real number. We will write this as $D(f)$ or D of f .

ex. $f(x) = x^2 + 2, D(f) = \mathbb{R}$
 $f(x) = \log(x), D(f) = (0, \infty)$
 $f(x) = \log|x|, D(f) = (-\infty, 0) \cup (0, \infty)$.


* Domains including the complex number set is hard to define since defining a domain in one interval can affect numbers in other intervals.

If f is a function, the **range** of f is the set $\{f(x) : x \in D(f)\}$. We write this as $R(f)$.


ex. $f(x) = \sin x \quad D(f) : \mathbb{R} \quad R(f) : [-1, 1] \quad \text{Codomain} : \mathbb{R}$

1.8.1 Graphical Representation

If f is a function, the *graph* of f is the set of ordered pairs $\{(x, f(x)) : x \in D(f)\}$. This is a subset of the x-y plane $\{(x, y) : x, y \in \mathbb{R}\} = \mathbb{R}^2$.



(a) $f(x) = \log|x|$



(b) $f(x) = \sin x$

For every subset of the plane (curve), is there a function that yields that subset?

No. Consider the **vertical line test**. A subset $T \subset \mathbb{R}$ is the graph of a function if and only if every vertical line in the xy plane intersects T at most once.

Consider $\begin{cases} 1, & \text{if } x \in \mathbb{Q} \\ 0, & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$ Is this a function?

1.9 Injectivity and Surjectivity

A function f is called **one-to-one** or **injective** if whenever

$$x \neq y, f(x) \neq f(y)$$

Equivalently, for all $x, y \in D(f)$, $x = y$ if and only if $f(x) = f(y)$.

Ex. $f(x) = 2x + 1$ injective?

Yes. Proof: let x, y be in the domain of $f(x)$ and suppose $f(x) = f(y)$. We need to prove $x = y$.

$$\begin{aligned} f(x) &= f(y) \\ 2x + 1 &= 2y + 1 \\ 2x &= 2y \\ x &= y \end{aligned}$$

Horizontal Line Test: A function f is one-to-one if and only if every horizontal line in the xy plane intersects the graph of f in at most one point.

Ex. $f(x) = x^2 + 1, D = [0, \infty)$ is this one-to-one?

Proof by contradiction: suppose f is not one-to-one. There exists real numbers $x, y \in D$ so that $x \neq y$, but $f(x) = f(y)$.

Without loss of generality, we can assume $x < y$. This implies $x^2 \leq x \cdot y < y \cdot y = y^2$. This is because $x \in [0, \infty)$ implies $x \geq 0$, $y > x$ implies $y > 0$, hence $x \cdot y < y \cdot y$, and it follows from $x \geq 0$.

Since $x^2 < y^2$, we have $x^2 + 1 < y^2 + 1$, $f(x) < f(y)$, but this contradicts assumption.

A function $f : D \rightarrow S$ is called **onto** or **surjective** if we let $R(f) = S$, and for every $y \in S$, there exists $x \in D$ such that $f(x) = y$.

$$\forall y \in S, \exists x \in D \text{ s.t. } f(x) = y$$

If a function is **both** one-to-one and onto, we say it is “one-to-one and onto”, or “**bijective**”.

For every $y \in S$, there exists a unique $x \in D$ such that $f(x) = y$.

$$\forall y \in S, \exists! x \in D \text{ s.t. } f(x) = y$$

1.10 Function interactions

If f and g are functions, with the domains $D(f)$ and $D(g)$ respectively, then we can define the functions $f + g$, $f - g$, and $f \cdot g$ in obvious ways.

The domain of each of these new functions is $D(f) \cap D(g)$.
ex. $f(x) = x^2$ and $g(x) = 1/x$. $f \cdot g(x) = x$ with Domain $D = \mathbb{R}/0$.

We can also define $(f/g)x = \frac{f(x)}{g(x)}$

$$\begin{aligned} D(f/g) &= (D(f) \cap D(g)) \setminus \{x \in D(g), g(x) = 0\} \\ &= \{x \in D(f) \cap D(g) : g(x) \neq 0\} \end{aligned}$$

If f and g are functions, we define the compositions $f \circ g = f(g(x))$. We first apply g then apply f .

$$D(f \circ g) = \{x \in D(g) : g(x) \in D(f)\}$$

ex. $g(x) = 1/x$ $f(x) = 1/x$
 $f \circ g(x) = x$ if $x \neq 0$.

1.11 Triangle inequality

$$|x + y| \leq |x| + |y|$$

$$||x| - |y|| \leq |x - y|$$

1.12 Polynomial

A **polynomial** is a function of the form

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

where $a_n \dots a_0$ are real numbers, and $a_n \neq 0$.

We define the degree of f to be n .

What about $f(x) = 0$?

Additionally, we say the function $f(x) = 0$ is also a polynomial, and we define its degree to be -1 .

1.13 Rational Function

A **rational function** is a function of the form

$$f(x) = \frac{P(x)}{Q(x)}$$

where P and Q are polynomials, and Q is not the zero polynomial.

Every polynomial is a rational function.

Ex. $f(x) = \frac{x^2+1}{2x+4}$, $D(f) = \mathbb{R} \setminus \{-2\}$.

2 Limits

2.1 Definition


Let f be a function, let $a \in \mathbb{R}$, and let $L \in \mathbb{R}$. We Say:

$$\lim_{x \rightarrow a} f(x) = L$$

If the following holds:

① There exists a number $t > 0$ so that

$$\{x \in \mathbb{R} : 0 < |x - a| < t\} \subset D(f)$$



* a doesn't have to be defined.

② For every real number $\epsilon > 0$, there exists a real number $\delta > 0$ so that the following is true: For every real number x satisfying the inequality $0 < |x - a| < \delta$, we have $|f(x) - L| < \epsilon$.

$$\forall \epsilon > 0, \exists \delta > 0 \text{ s.t. } \forall x \in \mathbb{R} \text{ with } 0 < |x - a| < \delta, \text{ we have } |f(x) - L| < \epsilon$$

Prove $\lim_{x \rightarrow 3} 2x = 6$.

① Find a $t > 0$ so that:

$$\{x \in \mathbb{R} : 0 < |x - 3| < t\} \subset D(f)$$

Any $t > 0$ works.

② Let $\epsilon > 0$, set $\delta = \epsilon/2$

Let $x \in \mathbb{R}$ with $0 < |x - 3| < \delta$.

Then $|2x - 6| = |2(x - 3)| = 2|x - 3| < 2\delta = \epsilon = \epsilon$.

Prove $\lim_{x \rightarrow 4} x^2 = 16$.

Since $D(f) = \mathbb{R}$, the domain requirement is satisfied.

Let $\epsilon > 0$, select $\delta = \min(\epsilon/9, 1)$.

Let $x \in \mathbb{R}$ with

$$0 < |x - 4| < \delta$$

$$|f(x) - L| = |x^2 - 16| = |(x - 4)(x + 4)| = |x + 4||x - 4|$$

$$\begin{aligned} |x + 4||x - 4| &< |x + 4|\delta \\ &\leq (|x| + 4)\delta \text{ We specified that } \delta \leq 1, \text{ then } |x| + 4 \leq 5 \\ &\leq (5 + 4)\delta \\ &= 9\delta \leq \epsilon \end{aligned}$$

Prove $\lim_{x \rightarrow 0} 2x \sin(1/x) = 0$

Since $D(f) = \mathbb{R} \setminus 0$, we can select $t = 1$, and $(-1, 0) \cup (0, 1) \subset D(f)$.

Let $\epsilon > 0$, select $\delta = \epsilon/2$

Let $x \in \mathbb{R}$ with $0 < |x - 0| < \delta$.

$$\begin{aligned} |f(x) - L| &= |2x \cdot \sin(1/x) - 0| \\ &= |2x \cdot \sin(1/x)| \\ &= 2|x| |\sin(1/x)| \quad \text{Since } 0 \leq |\sin(1/x)| < 1 \\ &\leq 2|x| = \epsilon \end{aligned}$$

2.1.1 Negating statements involving quantifiers

Every well crafted mathematical statement is either true or false. If S is a statement, either S is true, or S is false (*which is equivalent to “ S is false” is true)

$$\forall x \in \mathbb{R}, x^2 \geq 0 \text{ is true.} \quad \left| \quad \begin{array}{l} \exists x \in \mathbb{R}, \text{ s.t. } x^2 < 0. \\ (\forall x \in \mathbb{R}, x^2 \geq 0 \text{ is false.}) \end{array} \right.$$

General rules for negation: $\forall \rightarrow \exists, \exists \rightarrow \forall$, statement false.

$$\begin{array}{l} \forall x \in \mathbb{R}, \exists y \in \mathbb{R} \text{ s.t. } y > x \\ \\ \forall x \in \mathbb{R}, \exists y \in \mathbb{R} \text{ s.t. } \forall z \in \mathbb{R}, z > y > x \end{array} \quad \left| \quad \begin{array}{l} \text{There exists } x \in \mathbb{R}, \text{ s.t. } \forall y \in \mathbb{R}, y \leq x. \\ \\ \exists x \in \mathbb{R}, \forall y \in \mathbb{R}, \exists z \in \mathbb{R} \text{ s.t. } z \leq y \text{ or } y \leq x, \text{ or both.} \end{array} \right.$$

Definition of Limits: $\forall \epsilon > 0, \exists \delta > 0$ s.t. $\forall x \in \mathbb{R}$ with $0 < |x - a| < \delta$, we have $|f(x) - L| < \epsilon$.

⇓ Negation

$\exists \epsilon > 0, \forall \delta > 0, \exists x \in \mathbb{R}$ with $0 < |x - a| < \delta$, s.t. $|f(x) - L| \geq \epsilon$.

2.1.2 Proving limits false

Lets prove $f(x) = 2x$, $a = 1, L = 3, \lim_{x \rightarrow 1} f(x) = 3$ is false.

The domain req. holds. (*unfortunately*)

Choose $\epsilon = 1/2$.

Let $\delta > 0$.

Choose $x = \min(1.1, 1 + \delta/2)$ (We chose $1 + \delta/2$ to make $x > 1$)

Then $|f(x) - L| = |2x - 3|$

Since $0 < |x - a| < \delta$,

$|2x - 3| > |2 \cdot 1.1 - 3| = 0.8 \geq 1/2 = \epsilon$.

Let $f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$ Prove $\lim_{x \rightarrow 0} f(x)$ does not exist as a real number.

For every $L \in \mathbb{R}$, the statement " $\lim_{x \rightarrow 0} f(x) = L$ " is false.

⇓

$\forall L \in \mathbb{R}, \exists \delta > 0$ s.t. $\forall \delta > 0, \exists x \in \mathbb{R}$ with $0 < |x| < \delta$, s.t. $|f(x) - L| \geq \epsilon$.

Let $L \in \mathbb{R}$, select $\epsilon = 1/2$. Let $\delta > 0$.

If $L \geq 1/2$, select a number $x \in (0, \delta)$ with $x \notin \mathbb{Q}$. (The existence of x was proved in Hw)

Then $|f(x) - L| = |0 - L| = |L| \geq 1/2 = \epsilon$.

If $L < 1/2$, choose $x \in (0, \delta)$ with $x \in \mathbb{Q}$.

Then $|f(x) - L| = |1 - L| > 1/2 = \epsilon$.

2.2 Limit Rules

1 If $f(x) = k$ for all $x \in \mathbb{R}$, then for every $a \in \mathbb{R}$, $\lim_{x \rightarrow a} f(x) = k$. We write this as

$$\lim_{x \rightarrow a} k = k$$

2 If $f(x) = x$ for all $x \in \mathbb{R}$, then for every $a \in \mathbb{R}$, $\lim_{x \rightarrow a} f(x) = a$. We write this as

$$\lim_{x \rightarrow a} x = a$$

Proof: Let $a \in \mathbb{R}$, domain req. is met.

Let $\epsilon > 0$, choose $\delta = \epsilon/2$.

Then let $x \in \mathbb{R}$ with $0 < |x - a| < \delta$.

Then $|f(x) - a| = |x - a| < \delta = \epsilon/2 < \epsilon$.

3 Sum Rule: let $a \in \mathbb{R}$, If $\lim_{x \rightarrow a} f(x) = L$, and $\lim_{x \rightarrow a} g(x) = M$, then

$$\lim_{x \rightarrow a} f(x) + g(x) = L + M$$

Proof: Since $\lim_{x \rightarrow a} f(x) = L$, the domain requirement is met, thus there exists $t_1 > 0$ so that

$$\{x \in \mathbb{R} : 0 < |x - a| < t_1\} \subset D(f)$$

Similarly,

$$\{x \in \mathbb{R} : 0 < |x - a| < t_2\} \subset D(g)$$

Hence, if we define $t = \min(t_1, t_2)$, then

- $t > 0$
- $\{x \in \mathbb{R} : 0 < |x - a| < t\} \subset D(f)$ and
 $\{x \in \mathbb{R} : 0 < |x - a| < t\} \subset D(g)$

$$x \in \mathbb{R} : \{0 < |x - a| < t\} \subset D(f) \cap D(g) = D(f + g)$$

Let $\epsilon > 0$.

Define $\epsilon_1 = \epsilon/2$, $\epsilon_2 = \epsilon/2$.

Since we know $\lim_{x \rightarrow a} f(x) = L$, there exists a number $\delta_1 > 0$ so that whenever $0 < |x - a| < \delta_1$, we have $|f(x) - L| < \epsilon_1$.

Similarly, since $\lim_{x \rightarrow a} g(x) = M$, there exists $\delta_2 > 0$ so that for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta_2$, we have $|g(x) - M| < \epsilon_2$.

Let $\delta = \min(\delta_1, \delta_2)$. Then, satisfying all the requirements above, we have $x \in \mathbb{R}$ with $0 < |x - a| < \delta$, yielding

$$|f(x) - L| < \epsilon_1 = \epsilon/2$$

and

$$|g(x) - M| < \epsilon_2 = \epsilon/2$$

$$\begin{aligned} |(f + g)(x) - (L + M)| &\leq |f(x) - L| + |g(x) - M| \quad (\text{Triangle inequality}) \\ &< \epsilon/2 + \epsilon/2 = \epsilon \end{aligned}$$

4 Difference Rule: let $a \in \mathbb{R}$, If $\lim_{x \rightarrow a} f(x) = L$, and $\lim_{x \rightarrow a} g(x) = M$, then

$$\lim_{x \rightarrow a} f(x) - g(x) = L - M$$

Similar to the sum rule proof, we can prove the difference rule with the key step:

$$\begin{aligned} |[f(x) - g(x)] - (L - M)| &= |[f(x) - L] + [L - M + M - g(x)]| \\ &\leq |f(x) - L| + |L - M + M - g(x)| \\ &= |f(x) - L| + |g(x) - M| \end{aligned}$$

5 Product Rule: let $a \in \mathbb{R}$, If $\lim_{x \rightarrow a} f(x) = L$, and $\lim_{x \rightarrow a} g(x) = M$, then

$$\lim_{x \rightarrow a} (f \cdot g)(x) = L \cdot M$$

Proof: Domain requirement can be satisfied similar to sum rule.
Let $\epsilon > 0$, Choose $\epsilon_1 = ?$ $\epsilon_2 = ?$

Consider (scratch work): $f(x)g(x) - LM$
 $f(x)g(x) - LM \stackrel{?}{=} (f(x) - L)(g(x) - M)$

$$\begin{aligned} f(x)g(x) - LM &= (f(x) - L)(g(x) - M) + (f(x) - L)M + (g(x) - M)L \\ |f(x)g(x) - LM| &\leq |(f(x) - L)(g(x) - M)| + |(f(x) - L)M| + |(g(x) - M)L| \end{aligned}$$

Can all of these terms be $\epsilon/3$?

- If ϵ_1 and ϵ_2 are $\sqrt{\epsilon/3}$, their product $< \epsilon/3$.
- If $L \neq 0$, $\epsilon_1 = \frac{\epsilon}{3|L|}$, $L = 0$, $\epsilon_1 = ?$
- If $M \neq 0$, $\epsilon_2 = \frac{\epsilon}{3|M|}$, $M = 0$, $\epsilon_2 = ?$

$$\begin{aligned} \text{Choose } \epsilon_1 &= \min(\sqrt{\epsilon/3}, \frac{\epsilon}{3|L| + 1}) \\ \epsilon_2 &= \min(\sqrt{\epsilon/3}, \frac{\epsilon}{3|M| + 1}) \end{aligned}$$

Then we have $\epsilon_1 \leq \sqrt{\epsilon/3}$, $\epsilon_1 < \frac{\epsilon}{3|L|}$ if $L \neq 0$.
 $\epsilon_2 \leq \sqrt{\epsilon/3}$, $\epsilon_2 < \frac{\epsilon}{3|M|}$ if $M \neq 0$.

Since $\lim_{x \rightarrow a} f(x) = L$, there exists $\delta_1 > 0$ such for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta_1$, we have $|f(x) - L| < \epsilon_1$.

Similarly, we have $\delta_2 > 0$ such for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta_2$, we have $|g(x) - M| < \epsilon_2$.

Let $\delta = \min(\delta_1, \delta_2)$. Then for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta$, we have $|f(x) - L| < \epsilon$ and $|g(x) - M| < \epsilon$, so:

1. $|f(x) - L| < \sqrt{\epsilon/3}$
2. $|(f(x) - L)M| < \epsilon/3$
3. $|g(x) - M| < \sqrt{\epsilon/3}$
4. $|(g(x) - M)L| < \epsilon/3$

Finally,

$$\begin{aligned}
|(f \cdot g)(x) - LM| &= |(f(x) - L)(g(x) - M) + (f(x) - L)M + (g(x) - M)L| \\
&\leq |(f(x) - L)(g(x) - M)| + |(f(x) - L)M| + |(g(x) - M)L| \\
&< |\sqrt{\epsilon/3} \cdot \sqrt{\epsilon/3}| + \epsilon/3 + \epsilon/3 \\
&= \epsilon
\end{aligned}$$

6 Reciprocal Rule: let $a \in \mathbb{R}$, If $\lim_{x \rightarrow a} g(x) = M$ and $M \neq 0$, then

$$\lim_{x \rightarrow a} \frac{1}{g(x)} = \frac{1}{M}$$

Proof: Check the domain requirement:

Let $\epsilon' = |M|$,

Since $\lim_{x \rightarrow a} g(x) = M$, there exists δ' such that for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta'$, we have $|g(x) - M| = |M|$.

In particular, $g(x) \neq 0$. (Since if $g(x) = 0$ then $|g(x) - M| = |M|$, which isn't allowed).

Let $t = \delta'$, then $\{x \in \mathbb{R} : 0 < |x - a| < t\} \subset D(1/g)$.

The domain requirement is met.

In order to get the expression involving $|g(x) - M|$, we find the common denominator:

$$\left| \frac{1}{g(x)} - \frac{1}{M} \right| = \left| \frac{M - g(x)}{g(x)M} \right| = \left| \frac{g(x) - M}{|g(x)||M|} \right|$$

We can construct bound: Let $\epsilon_1 = \frac{|M|}{2}$. Since $\lim_{x \rightarrow a} g(x) = M$, there exists δ_1 such that if $0 < |x - a| < \delta_1$, then $|g(x) - M| < \frac{|M|}{2}$.

$$\begin{aligned}
|M| &= |M - g(x) + g(x)| \\
&\leq |M - g(x)| + |g(x)| \\
&= |g(x) - M| + |g(x)| \\
|M| &< \left| \frac{|M|}{2} \right| + |g(x)| \\
\frac{|M|}{2} &< |g(x)|
\end{aligned}$$

\Downarrow

$$\frac{1}{g(x)} < \frac{2}{|M|}, \text{ where } 0 < |x - a| < \delta_1 \text{ and } |M| \neq 0$$

Now we can move onto the entire expression.

Since $\lim_{x \rightarrow a} g(x) = M$, there exists δ_2 such if $0 < |x - a| < \delta_2$, then

$$|g(x) - M| < \frac{|M|^2 \epsilon}{2}$$

Let $\epsilon > 0$. We can choose $\delta = \min(\delta_1, \delta_2)$. For all x , $0 < |x - a| < \delta$, we have

$$\left| \frac{1}{g(x)} - \frac{1}{M} \right| = \left| \frac{g(x) - M}{|M|g(x)} \right| < \frac{M^2 \epsilon}{2|M|} \cdot \frac{1}{|h(x)|} < \frac{M^2 \epsilon}{2|M|} \cdot \frac{2}{|M|} < \epsilon$$

7 Quotient Rule: let $a \in \mathbb{R}$, if $\lim_{x \rightarrow a} f(x) = L$, $\lim_{x \rightarrow a} g(x) = M$ and $M \neq 0$,

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{L}{M}$$

Proof: We can apply the product rule to f and $\frac{1}{g}$.

8 “Limits are a local property”: Let f and g be functions and let $a \in \mathbb{R}$. If there exists $t > 0$ so that $f(x) = g(x)$ for all $x \in (a - t, a) \cup (a, a + t)$ (and in particular both functions are defined),

Then, if $\lim_{x \rightarrow a} f(x) = L$, we have $\lim_{x \rightarrow a} g(x) = L$

ex. $f(x) = x$, $g(x) = x$, but $D(g) = [-1, 1] \setminus \{0\}$

Since $\lim_{x \rightarrow 0} f(x) = 0$ and $f(x) = g(x)$ for all $x \in [-1, 0] \cup [0, 1]$,
by LAALP, $\lim_{x \rightarrow a} g(x) = 0$.

2.3 Squeeze Theorem

Let f, g and h be functions and let $a \in \mathbb{R}$. suppose there exists $t > 0$ so that

$$f(x) \leq g(x) \leq h(x)$$

For all $x \in (a - t, a) \cup (a, a + t)$ (for all $x \in \mathbb{R}$ with $0 < |x - a| < t$,

Suppose also that $\lim_{x \rightarrow a} f(x) = L$ and $\lim_{x \rightarrow a} h(x) = L$.


Then $\lim_{x \rightarrow a} g(x) = L$.

ex. $g(x) = 2x \sin(\frac{1}{x})$

$$\begin{aligned} f(x) &= -2|x|, \quad h(x) = 2|x|. \\ |g(x) = 2|x| &< |\sin \frac{1}{x}| < 2|x| \text{ for all } x \neq 0 \\ |g(x)| &\leq h(x) \Rightarrow g(x) \leq h(x). \end{aligned}$$

and $|g(x)| < -f(x) \Rightarrow g(x) \geq f(x)$.

Then, by the squeeze theorem, $\lim_{x \rightarrow 0} g(x) = 0$.



Proof: Domain req.: By the hypothesis of the theorem there exists $t > 0$ so that $\{x \in \mathbb{R}, 0 < |x - a| < t\} \subset D(g)$.

Let $\epsilon > 0$. Select $\epsilon_1 = \epsilon/3$, $\epsilon_2 = \epsilon/3$.

Since $\lim_{x \rightarrow a} f(x) = L$, there exists $\delta_1 > 0$ so that for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta_1$, we have $|f(x) - L| < \epsilon_1$.

Similarly, there exists $\delta_2 > 0$ so that for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta_2$, we have $|f(x) - L| < \epsilon_2$.

Let $\delta = \min(\delta_1, \delta_2, t)$, then, for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta$, we have:

$$\begin{aligned} |g(x) - L| &= |g(x) - f(x) + f(x) - L| \\ &\leq |g(x) - f(x)| + |f(x) - L| \\ &\leq |h(x) - f(x)| + |f(x) - L| \text{ for all } x \in (a - t, a) \cup (a, a + t) \\ &= |h(x) - L + L - f(x)| + |f(x) - L| \\ &= |h(x) - L| + |L - f(x)| + |f(x) - L| \\ &= |h(x) - L| + 2|f(x) - L| \\ &< \epsilon_2 + \epsilon_1 + \epsilon_1 = \epsilon \end{aligned}$$

2.4 One-sided Limits

Let f be a function and $a \in \mathbb{R}$, $L \in \mathbb{R}$. We say $\lim_{x \rightarrow a^+} f(x) = L$ if the following holds:

- There exists $t > 0$ so that $(a, a + t) \subset D(f)$.
- $\forall \epsilon > 0, \exists \delta > 0$ s.t. $\forall x \in (a, a + \delta)$, we have $|f(x) - L| < \epsilon$.

Similarly, we can define $\lim_{x \rightarrow a^-} f(x) = L$.

Let f be a function and $a \in \mathbb{R}, L \in \mathbb{R}$. We say $\lim_{x \rightarrow a^-} f(x) = L$ if the following holds:

- There exists $t > 0$ so that $(a - t, a) \subset D(f)$.
- $\forall \epsilon > 0, \exists \delta > 0$ s.t. $\forall x \in (a - \delta, a)$, we have $|f(x) - L| < \epsilon$.

1

$$\lim_{x \rightarrow a} f(x) = L$$

if and only if

- $\lim_{x \rightarrow a^-} f(x) = L$
- $\lim_{x \rightarrow a^+} f(x) = L$

2.5 Limits at Infinity

Let f be a function and $L \in \mathbb{R}$. We say $\lim_{x \rightarrow \infty} f(x) = L$ if:

- There exists $T \in \mathbb{R}$ so that $(T, \infty) \subset D(f)$
- For all $\epsilon > 0$, there exists a number $R \in \mathbb{R}$ such that if $x \in \mathbb{R}$ and $x > R$, then $|f(x) - L| < \epsilon$

Similarly,

Let f be a function and $L \in \mathbb{R}$. We say $\lim_{x \rightarrow -\infty} f(x) = L$ if:

- There exists $T \in \mathbb{R}$ so that $(-\infty, T) \subset D(f)$
- For all $\epsilon > 0$, there exists a number $R \in \mathbb{R}$ such that if $x \in \mathbb{R}$ and $x < R$, then $|f(x) - L| < \epsilon$

As seen in figure a) and b), both functions have a limit approaching infinity. We write their limits as


$$\lim_{x \rightarrow a} f(x) = (\pm)\infty$$

Thus, we say $x = a$ are their **asymptotes**.


E.x. Let $f(x) = 1/x$, show $\lim_{x \rightarrow \infty} f(x) = 0$

Let $\epsilon > 0$, select $R = 1/\epsilon$.

If $x \in (R, \infty)$, then $|f(x) - L| = |f(x)| = |1/x| = 1/x < 1/R = \epsilon$



(a) $y = \frac{1}{x^2}$



(b) y approaches $-\infty$ as x approaches 2

E.x. $f(x) = \sin(x)$, show for every $L \in \mathbb{R}$, $\lim_{x \rightarrow \infty} f(x) = L$ is false.

Proof: Let $L \in \mathbb{R}$,

Since $D(f) = \mathbb{R}$, the domain requirement is met.

Select $\epsilon = 1$. Let $R \in \mathbb{R}$.

If $L \leq 0$, select a number x with $x > R$ of the form $x = \frac{\pi}{2} + 2\pi n, n \in \mathbb{Z}$.

Then $f(x) = 1$, so $|f(x) - L| = |1 - L| = 1 - L \geq 1 = \epsilon$.

If $L > 0$ select a number $x \in \mathbb{R}$ with $x > R$ of the form $x = \frac{3\pi}{2} + 2\pi n, n \in \mathbb{Z}$.

Then $f(x) = -1$, so $|f(x) - L| = |-1 - L| = 1 + L \geq 1 = \epsilon$.

2.6 Infinite Limits

Let f be a function and $a \in \mathbb{R}$. We say $\lim_{x \rightarrow a} f(x) = \infty(-\infty)$, if:

- Exists $t > 0$ such that $\{x \in \mathbb{R} : 0 < |x - a| < t\} \subset D(f)$
- For all $M \in \mathbb{R}$, exists $\delta > 0$, such that for all $x \in \mathbb{R}$ with $0 < |x - a| < \delta$, we have $f(x) > M$ ($f(x) < M$).

Ex. $f(x) = \frac{1}{x^2}$, show $\lim_{x \rightarrow 0} f(x) = \infty$.

Since $D(f) = \mathbb{R} \setminus \{0\}$, any $t > 0$ will work.

Next, let $M \in \mathbb{R}$, select $\delta = \frac{1}{\sqrt{|M|+1}}$.

If $0 < |x - a| < \delta$, $0 < |x| < \delta$, then

$$\begin{aligned} f(x) &= \frac{1}{x^2} \\ &> \frac{1}{\frac{1}{|M+1|}} \\ &= |M| + 1 \geq M \end{aligned}$$

We can also define $\lim_{x \rightarrow a^+} f(x) = \infty$, or $\lim_{x \rightarrow a^-} f(x) = \infty$ in the obvious way.

We say $\lim_{x \rightarrow \infty} f(x) = \infty$ if:

- $T \in \mathbb{R}$ s.t. $(T, \infty) \subset D(f)$.
- For all $M \in \mathbb{R}$, there exists $R \in \mathbb{R}$, such that for all $x \in \mathbb{R}$ with $x > R$, we have $f(x) > M$.

We can also define $\lim_{x \rightarrow \infty} f(x) = -\infty$, $\lim_{x \rightarrow -\infty} f(x) = \infty$, $\lim_{x \rightarrow \infty} f(x) = -\infty$, $\lim_{x \rightarrow -\infty} f(x) = -\infty$ in the obvious way.

2.7 Existence of Limits

If f is a function and $a \in \mathbb{R}$ for all $L \in \mathbb{R}$, the statement “ $\lim_{x \rightarrow a} f(x) = L$ ” is false, then we say $\lim_{x \rightarrow a} f(x)$ does not exist as a real number.

If f is a function, $a \in \mathbb{R}$, we say $\lim_{x \rightarrow a} f(x)$ doesn't exist, if for all $L \in \mathbb{R}$:

$$\begin{cases} \lim_{x \rightarrow a} f(x) = L \text{ is false, and} \\ \lim_{x \rightarrow a} f(x) = \infty \text{ is false, and} \\ \lim_{x \rightarrow a} f(x) = -\infty \text{ is false} \end{cases}$$

2.7.1 Use of product rule

Despite that limits at infinity are defined as limits that exist, we cannot simply apply the product rules to their product. We need to give individual proof.

Example: $\lim_{x \rightarrow a} \frac{1}{(x-a)^2} \cdot \lim_{x \rightarrow a} (x-a)^2$.

2.8 Asymptotes


When at least one of the following is true:

$$\lim_{x \rightarrow a} f(x) = \infty, \lim_{x \rightarrow a^-} f(x) = \infty, \lim_{x \rightarrow a^+} f(x) = \infty,$$

$$\lim_{x \rightarrow a} f(x) = -\infty, \quad \lim_{x \rightarrow a^-} f(x) = -\infty, \quad \lim_{x \rightarrow a^+} f(x) = -\infty,$$

$x = a$ is a vertical asymptote.

Example Limit of $y = \tan(x)$



The vertical asymptotes are:

$$x = \frac{\pi}{2} + n\pi \quad (n \in \mathbb{Z})$$

2.9 Limit dependency on Power of Polynomial

Let P and Q be non-zero polynomials, of degree n and m respectively, i.e.

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$$

$$Q(x) = b_n x^n + b_{n-1} x^{n-1} + \dots + b_0$$

Let $f(x) = \frac{P(x)}{Q(x)}$.

1 If $\deg P < \deg Q$ ($n < m$), then

$$\lim_{x \rightarrow \infty} f(x) = 0$$

2 If $\deg P = \deg Q$ ($n = m$), then

$$\lim_{x \rightarrow \infty} f(x) = \frac{a_n}{b_m}$$

3 If $\deg P > \deg Q$ ($n > m$), then

$$\lim_{x \rightarrow \infty} f(x) = \infty \text{ if } \frac{a_n}{b_m} > 0$$

$$\lim_{x \rightarrow \infty} f(x) = -\infty \text{ if } \frac{a_n}{b_m} < 0$$

Proof of 1 To satisfy the domain requirement, we need to ensure $Q(x) \neq 0$. We want to find $T \in R$ so that $Q(x) \neq 0$ for all $x > T$.

We define $B = |b_{m-1}| + |b_{m-2}| + \dots + |b_0|$.

If $x \geq 1$, then

$$\begin{aligned} |b_{m-1}x^{m-1} + b_{m-2}x^{m-2} + \dots + b_0| &\leq |b_{m-1}x^{m-1}| + |b_{m-2}x^{m-2}| + \dots + |b_0| \\ &\leq |b_{m-1}||x^{m-1}| + |b_{m-2}||x^{m-1}| + \dots + |b_0||x^{m-1}| \\ &\leq B|x^{m-1}| \end{aligned}$$

Thus, we conclude if $x \geq 1$,

$$a - |b| \leq a + b \leq a + |b|$$

\Downarrow

$$b_m x^m - B x^{m-1} \leq Q(x) \leq b_m x^m + B x^{m-1}$$

Next, we take

$$b_m x^m - B x^{m-1} = x^{m-1}(b_m x - B)$$

And let $T = B/|b_m|$.

If $x > T$, then

If $b_m < 0$,

Since $B > 0$, $b_mx < 0$, $b_mx - B < 0$.

Also $x^{m-1} > 0$, thus $x^{m-1}(b_mx - B) < 0$.

$Q(x) \neq 0$.

If $b_m > 0$,

$b_mx - B > 0$.

Also $x^{m-1} > 0$, thus $x^{m-1}(b_mx - B) > 0$.

$Q(x) \neq 0$.

Thus, $T = B/|b_m|$ will work.

Next, let $\epsilon > 0$.

We need to find $R \in \mathbb{R}$, so that for all $x > R$, $|\frac{P(x)}{Q(x)}| < \epsilon$.

Similar to B , we define $C = |a_{n-1}| + |a_{n-2}| + \dots + |a_0|$. Then if $x \geq 1$,

$$|P(x)| \leq Cx^n$$

Next, if we choose $x > 2B/|b_m|$, then

$$x^{m-1}(|b_m|x - 2B) > 0$$

$$|b_m|x^m - 2Bx^{m-1} > 0$$

$$\frac{1}{2}|b_m|x^m > Bx^{m-1}$$

$$-\frac{1}{2}|b_m|x^m < -Bx^{m-1} \Rightarrow \text{add to both sides } |b_m|x^m$$

$$\frac{1}{2}|b_m|x^m < |b_m|x^m - Bx^{m-1} < Q(x)$$

Select $R = \max(1, 2B/|b_m|, 2C/|b_m|\epsilon)$, then:

$$\begin{aligned} \frac{|P(x)|}{|Q(x)|} &\leq \frac{Cx^n}{\frac{1}{2}|b_m|x^m} = \frac{2C}{|b_m|} \cdot \frac{1}{x^{n-m}} \\ &\leq \frac{2C}{|b_m|} \cdot \frac{1}{x} \quad \text{Since } n - m \geq 1, 1/x \geq x \\ &< \frac{2C}{|b_m|} \cdot \frac{1}{R} \\ &< \frac{2C}{|b_m|} \cdot \frac{|b_m|\epsilon}{2C} = \epsilon \end{aligned}$$

Proof of 2: Suppose

$$\left. \begin{aligned} P(x) &= a_nx^n + \dots + a_0 \\ Q(x) &= b_mx^n + \dots + a_0 \end{aligned} \right\} \text{ to some degree}$$

First,

$$\lim_{x \rightarrow \infty} \frac{a_n}{b_n} = \frac{a_n}{b_n}$$

Next,

$$\frac{P(x)}{Q(x)} - \frac{a_n}{b_n} = \frac{b_n P(x) - a_n Q(x)}{b_n Q(x)}$$

when $Q(x) \neq 0$.

$$\begin{aligned} &= \frac{b_n(a_n x^n + a_{n-1} x^{n-1} + \dots + a_0) - a_n(b_n x^n + b_{n-1} x^{n-1} + \dots + b_0)}{b_n Q(x)} \\ &= \frac{b_n a_n x^n - b_n a_n x^n + b_n(a_{n-1} x^{n-1} + \dots + a_0) - a_n(b_{n-1} x^{n-1} + \dots + b_0)}{b_n Q(x)} \\ &= \frac{S(x)}{b_n Q(x)} \text{ Where deg. of } S \text{ is less than deg. of } Q \end{aligned}$$

So

$$\lim_{x \rightarrow \infty} \left(\frac{P(x)}{Q(x)} - \frac{a_n}{b_n} \right) = \lim_{x \rightarrow \infty} \frac{S(x)}{b_n Q(x)} = 0$$

Then

$$\lim_{x \rightarrow \infty} \frac{P(x)}{Q(x)} = \lim_{x \rightarrow \infty} \left(\frac{P(x)}{Q(x)} - \frac{a_n}{b_n} + \frac{a_n}{b_n} \right) = 0 + \frac{a_n}{b_n} = \frac{a_n}{b_n}$$

Proof of 3: To satisfy the domain requirement, we need to ensure $Q(x) \neq 0$. We want to find $T \in \mathbb{R}$ so that $Q(x) \neq 0$ for all $x > T$.

We define $B = |b_{m-1}| + |b_{m-2}| + \dots + |b_0|$.

If $x \geq 1$, then

$$\begin{aligned} |b_{m-1} x^{m-1} + b_{m-2} x^{m-2} + \dots + b_0| &\leq |b_{m-1} x^{m-1}| + |b_{m-2} x^{m-2}| + \dots + |b_0| \\ &\leq |b_{m-1}| |x^{m-1}| + |b_{m-2}| |x^{m-1}| + \dots + |b_0| |x^{m-1}| \\ &\leq B |x^{m-1}| \end{aligned}$$

We can satisfy the domain requirement similar to proof 1.

Next, let $M \in \mathbb{R}$.

We need to find $R \in \mathbb{R}$, so that for all $x > R$, $P(x)/Q(x) > M$.

We can assume $a_n > 0, b_n > 0$, and that $n > m$

Similar to B , we define $A = |a_{n-1}| + |a_{n-2}| + \dots + |a_0|$. We have:

$$a_n x^n - A x^{n-1} \leq P(x) \leq a_n x^n + A x^{n-1}$$

Next, if we choose $x > 2A/a_n$, then

$$\begin{aligned}
x^{n-1}(a_n x - 2A) &> 0 \\
a_n x^n - 2Ax^{n-1} &> 0 \\
\frac{1}{2}a_n x^n &> Ax^{n-1} \\
-\frac{1}{2}a_n x^n &< -Ax^{n-1} \\
\frac{1}{2}a_n x^n &< a_n x^n - Ax^{n-1} < P(x)
\end{aligned}$$

We define $C = B + b_m$. If $x \geq 1$,

$$\begin{aligned}
Q(x) &\leq Bx^m + b_m x^m \\
Q(x) &\leq Cx^m
\end{aligned}$$

We can define $M' = \max(1, M)$, so that $M' \geq M$.
Select $R = \max(1, 2A/a_n, 2Ca_n M')$

Let $x > R$. Since $\frac{1}{2}a_n x^n < P(x)$, $Q(x) \leq Cx^m$

$$\begin{aligned}
\frac{P(x)}{Q(x)} &> \frac{1/2|a_n|x^n}{Cx^m} = \frac{1}{2C|a_n|} \cdot x^{n-m} \\
&\geq \frac{1}{2C|a_n|} \cdot x \quad \text{Since } n - m \geq 1, 1/x \leq x \\
&\geq \frac{1}{2C|a_n|} \cdot M' \\
&> \frac{1}{2C|a_n|} \cdot 2C|a_n|M = M
\end{aligned}$$

For $a_n < 0$ and $b_m < 0$, the proof is similar where $P(x)/Q(x) > M$;

For $a_n > 0$ and $b_m < 0$, or $a_n < 0$ and $b_m > 0$, the proof is different as $P(x)/Q(x) < M$.

2.10 Limit Laws

*Suppose c is a constant and $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exists,

$$\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \quad (1)$$

$$\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x) \quad (2)$$

$$\lim_{x \rightarrow a} [cf(x)] = c \lim_{x \rightarrow a} f(x) \quad (3)$$

$$\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x) \quad (4)$$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} \quad (\lim_{x \rightarrow a} g(x) \neq 0) \quad (5)$$

$$\lim_{x \rightarrow a} [f(x)]^n = [\lim_{x \rightarrow a} f(x)]^n$$

$$\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$$

2.11 Limit Calculations

Example 1. Find $\lim_{h \rightarrow 0} \frac{(3+h)^2 - 9}{h}$

$$\begin{aligned} F(h) &= \frac{(3+h)^2 - 9}{h} \\ &= \frac{h^2 + 6h + 9 - 9}{h} \\ &= h + 6 \end{aligned}$$

By evaluation, $\lim_{h \rightarrow 0} h + 6 = 6$

2.11.1 Direct Substitution Property

If f is a **polynomial** or a **rational function** and a is in the domain of f , then:

$$\lim_{x \rightarrow a} f(x) = f(a)$$

Example 2. Find $\lim_{t \rightarrow 0} \frac{\sqrt{t^2+9}-3}{t^2}$

$$\begin{aligned}
 &= \lim_{t \rightarrow 0} \frac{\sqrt{t^2+9}-3}{t^2} \cdot \frac{\sqrt{t^2+9}+3}{\sqrt{t^2+9}+3} \\
 &= \lim_{t \rightarrow 0} \frac{t^2+9-9}{t^2(\sqrt{t^2+9}+3)} \\
 &= \lim_{t \rightarrow 0} \frac{t^2}{t^2\sqrt{t^2+9}+3t^2} \\
 &= \lim_{t \rightarrow 0} \frac{1}{\sqrt{t^2+9}+3} \\
 &= \frac{1}{\sqrt{\lim_{t \rightarrow 0} t^2+9}+3} \\
 &= \frac{1}{\sqrt{9}+3} = \frac{1}{6}
 \end{aligned}$$

Example 3. Find $\lim_{x \rightarrow 1} \frac{\sqrt[3]{x}-1}{\sqrt{x}-1}$

3 Continuity

Let f be a function, and let $a \in \mathbb{R}$. We say that f is continuous at a , if

$$\lim_{x \rightarrow a} f(x) = f(a)$$

In particular, this implies:

- ① $a \in D(f)$
- ② $f(a) \in \mathbb{R}$
- ③ $\lim_{x \rightarrow a} f(x)$ exists as a real number

Ex.

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{Q} \\ 0, & \text{if } x \notin \mathbb{Q} \end{cases}$$

$$\lim_{x \rightarrow 0} f(x) = 0 = f(0).$$

So f is continuous at 0. It is not continuous at every other point. If f is not continuous at a , we say it is **discontinuous** at a .

If $\lim_{x \rightarrow a^+} f(x) = f(a)$, we say f is right continuous.

If $\lim_{x \rightarrow a^-} f(x) = f(a)$, we say f is left continuous.

Let f be a function, and let (a, b) be an (open) interval. We say f is continuous on (a, b) if it is continuous at every point $x \in \mathbb{R}, x \in (a, b)$.

If f is a function , and $[a, b]$ is a closed interval, We say f is continuous on $[a, b]$ if

- $\lim_{x \rightarrow c} f(x) = f(c)$ for all $c \in (a, b)$.
- $\lim_{x \rightarrow a^+} f(x) = f(a)$
- $\lim_{x \rightarrow b^-} f(x) = f(b)$

Lemma 1:

Let f be a function that is continuous at the point $c \in \mathbb{R}$.

If $f(c) \neq 0$, then there exists $\delta > 0$, so that $f(x)$ is defined, and $f(x) \neq 0$ for all

$$x \in (c - \delta, c + \delta)$$

Ex.

$$f(x) = \begin{cases} 1, & \text{if } x \in \mathbb{Q} \\ x, & \text{if } x \notin \mathbb{Q} \end{cases}$$

Proof of Lemma:

Since f is continuous at c , $\lim_{x \rightarrow c} f(x) = f(c) \neq 0$.

Choose $\epsilon = \frac{|f(c)|}{2}$

Then, there exists $\delta > 0$, so that for every $x \in \mathbb{R}$ with $0 < |x - c| < \delta$, $f(x)$ exists and $|f(x) - f(c)| < \epsilon$.