BIOS668 HW#4

Sara O'Brien

Honor Code: On my honor, I have neither given nor received unauthorized aid on this assignment.

Q1)

1) Note that we observe no outcomes in one arm and no outcomes in the other.

We assume that n = knz.

To determine where to evaluate power, we can use the following

test statistics based on the sample nears \hat{x}_1, \hat{x}_2 . $\hat{x}_1 \sim N(\mu_1, \frac{\sigma^2}{n_1}) = N(\mu_1, \frac{\sigma^2}{\kappa n_2})$ $\hat{x}_2 \sim N(\mu_2, \frac{\sigma^2}{n_2})$

We can then compare these two means:
$$\bar{\chi}_1 - \bar{\chi}_2 \sim N(\mu_1 - \mu_2, \frac{\sigma^2}{kn_2} + \frac{\sigma^2}{n^2})^* N(\mu_1 - \mu_2, \frac{(\kappa+1)\sigma^2}{kn_2})$$

If we denote of kill or and M=M,-Mz we are left with x, - xz ~ N (m, +2)

We now have a standard 1-sample superiority test. 50, We know that
$$N_{z} = \frac{\left(\frac{1}{2} - \frac{\alpha}{2} + \frac{1}{2} - \frac{\beta}{2}\right)^{2}}{\left(\frac{d}{\sigma_{w}}\right)^{2}} = \frac{\sigma_{w}^{2} \left(\frac{1}{2} - \frac{\alpha}{2} + \frac{1}{2} - \frac{\beta}{2}\right)^{2}}{\sigma_{w}^{2} \left(\frac{1}{2} - \frac{\alpha}{2} + \frac{1}{2} - \frac{\beta}{2}\right)^{2}} = \frac{\left(\frac{1}{2} - \frac{\alpha}{2} + \frac{1}{2} - \frac{\beta}{2}\right)^{2} \sigma^{2} \left(\frac{1}{2} + \frac{1}{2} - \frac{\beta}{2}\right)^{2}}{\sigma^{2} \left(\frac{1}{2} - \frac{\alpha}{2}\right)^{2} \sigma^{2} \left(\frac{1}{2} - \frac{\alpha}{2}\right)^{2}}$$

Then, since NI= Knz,

$$N_{1} = KN_{2} > \frac{K(Z_{1-\alpha/2} + Z_{1-\beta})^{2} \sigma^{2}(K+1)}{K\delta^{2}} = \frac{(Z_{1-\alpha/2} + Z_{1-\beta}) \sigma^{2}(K+1)}{\delta^{2}}$$

In this study, we plan to test if treatment strategy A is not too different from treatment strategy B for a specific disease. We are measuring this equivalence using time in days to disease resolution since randomization, where 'not too different' or equivalence is defined as 1.6 days. Hence, our null hypothesis is that the difference between treatments, in either direction, is ≥ 1.6 days. Our alternative hypothesis is that this difference is < 1.6 days. The study is interested in having 80% power and testing at alpha-level 0.05 for this controlled equivalence test. The true mean difference between treatments is 0.35 days and the standard deviation is 7.0 days. 582 participants are randomized into treatment A and 291 participants are randomized into treatment B in two arms to detect the statistically meaningful difference of 1.6 days under the study conditions. There is no interim analysis and since this is a parallel study design, participants only receive the treatment for the group they were randomized into for the duration of the study. Similar to a one sample design, our rejection region containing the critical

values that we use to determine when to reject our null hypothesis is set at $|T| \le c = \sqrt{n}\delta - z_{1-\alpha}$, with adjustment for two-sample design.