

Adjoint Current-Based Approaches to Brachytherapy Optimization

Jeremy A. Roberts

Department of Engineering Physics
University of Wisconsin–Madison

May 19, 2008

Outline

Background

Data and Modeling

Current-displacement: a “Nudge”

Normed CD: a New Greedy Criterion



Introduction

Introduction

- ▶ Brachytherapy a leading treatment modality for many cancers



Introduction

Introduction

- ▶ Brachytherapy a leading treatment modality for many cancers
- ▶ Interstitial, i.e. internal, brachytherapy very useful in treating prostate cancer



Introduction

Introduction

- ▶ Brachytherapy a leading treatment modality for many cancers
- ▶ Interstitial, i.e. internal, brachytherapy very useful in treating prostate cancer
- ▶ Both low dose (permanent) and high dose (temporary) sources used

Introduction

Introduction

- ▶ Brachytherapy a leading treatment modality for many cancers
- ▶ Interstitial, i.e. internal, brachytherapy very useful in treating prostate cancer
- ▶ Both low dose (permanent) and high dose (temporary) sources used
- ▶ Focus here is on *permanent* sources



Introduction

Motivation

- ▶ Optimizing seed placement is extremely difficult!



Introduction

Motivation

- ▶ Optimizing seed placement is extremely difficult!
- ▶ Robust methods applied; **good results, long time**



Introduction

Motivation

- ▶ Optimizing seed placement is extremely difficult!
- ▶ Robust methods applied; **good results, long time**
- ▶ New methods desired for “on the fly” treatment planning

Introduction

Motivation

- ▶ Optimizing seed placement is extremely difficult!
- ▶ Robust methods applied; **good results, long time**
- ▶ New methods desired for “on the fly” treatment planning
- ▶ Work done at UW-Madison to use “adjoint” methods from reactor physics

Introduction

Motivation

- ▶ Optimizing seed placement is extremely difficult!
- ▶ Robust methods applied; **good results, long time**
- ▶ New methods desired for “on the fly” treatment planning
- ▶ Work done at UW-Madison to use “adjoint” methods from reactor physics
- ▶ Near-optimal results...but could they be better?

Introduction

Motivation

- ▶ Optimizing seed placement is extremely difficult!
- ▶ Robust methods applied; **good results, long time**
- ▶ New methods desired for “on the fly” treatment planning
- ▶ Work done at UW-Madison to use “adjoint” methods from reactor physics
- ▶ Near-optimal results...but could they be better?
- ▶ Here, two methods incorporating further aspects of the “adjoint” theory explored



Neutral Particle Transport

The Boltzmann Equation

A time-independent treatment of particle interactions suffices, done via

$$[\hat{\Omega} \cdot \vec{\nabla} + \Sigma(\vec{r}, E)]\psi(\vec{r}, \hat{\Omega}, E) = q_{ex}(\vec{r}, \hat{\Omega}, E) + \int dE' \int d\Omega' \Sigma_s(\vec{r}, E' \rightarrow E, \hat{\Omega}' \cdot \hat{\Omega})\psi(\vec{r}, \hat{\Omega}', E') \quad (1)$$

where \vec{r} denotes position, $\hat{\Omega}$ denotes the direction of travel, and E is the energy of a particle. [3]



Transport Phase-Space




Figure: The position and direction variables describing a particle.



Forward Flux

The solution of the Boltzmann equation is the *angular flux*, $\psi(\vec{r}, \hat{\Omega}, E)$. From this, the *scalar flux* is defined

$$\phi(\vec{r}, E) = \int_{4\pi} d\Omega \psi(\vec{r}, \hat{\Omega}, E). \quad (2)$$

Forward Flux

The solution of the Boltzmann equation is the *angular flux*, $\psi(\vec{r}, \hat{\Omega}, E)$. From this, the *scalar flux* is defined

$$\phi(\vec{r}, E) = \int_{4\pi} d\Omega \psi(\vec{r}, \hat{\Omega}, E). \quad (2)$$

- ▶ $\phi(\vec{r}, E)$ describes the particle population at a given point and at a given energy → *direction of motion not considered*
- ▶ Reaction rates, e.g. dose rates, calculated via

$$R(\vec{r}) = \int_0^\infty \phi(\vec{r}, E) DF(E) dE \quad (3)$$

where $DF(E)$ are flux-to-dose conversion factors.

Neutral Particle Transport

The Adjoint Transport Equation

- ▶ Particles were just *forward transported*, i.e. emitted from a source and absorbed at some detector.

The Adjoint Transport Equation

- ▶ Particles were just *forward transported*, i.e. emitted from a source and absorbed at some detector.
- ▶ Let a source q^+ be the “pseudo particles” representing a desired *detector response*

The Adjoint Transport Equation

- ▶ Particles were just *forward transported*, i.e. emitted from a source and absorbed at some detector.
- ▶ Let a source q^+ be the “pseudo particles” representing a desired *detector response*
- ▶ Set those particles in motion, but apply “reverse physics” ...

The Adjoint Transport Equation

- ▶ Particles were just *forward transported*, i.e. emitted from a source and absorbed at some detector.
- ▶ Let a source q^+ be the “pseudo particles” representing a desired *detector response*
- ▶ Set those particles in motion, but apply “reverse physics” ...
- ▶ ... the adjoint equation:

$$[-\hat{\Omega} \cdot \vec{\nabla} + \Sigma(\vec{r}, E)]\psi^+(\vec{r}, \hat{\Omega}, E) = q^+(\vec{r}, \hat{\Omega}, E) + \int dE' \int d\Omega' \Sigma_s(\vec{r}, E \rightarrow E', \hat{\Omega} \cdot \hat{\Omega}') \psi^+(\vec{r}, \hat{\Omega}', E') \quad (4)$$

The Adjoint Flux

- ▶ Solution of the adjoint equation → *angular flux*, $\psi^+(\vec{r}, \hat{\Omega}, E)$



Neutral Particle Transport

The Adjoint Flux

- ▶ Solution of the adjoint equation → *angular flux*, $\psi^+(\vec{r}, \hat{\Omega}, E)$
- ▶ From this, the *adjoint scalar flux*

$$\phi^+(\vec{r}, E) = \int_{4\pi} d\Omega \psi(\vec{r}, \hat{\Omega}, E). \quad (5)$$



The Adjoint Flux

- ▶ Solution of the adjoint equation → *angular flux*, $\psi^+(\vec{r}, \hat{\Omega}, E)$
- ▶ From this, the *adjoint scalar flux*

$$\phi^+(\vec{r}, E) = \int_{4\pi} d\Omega \psi(\vec{r}, \hat{\Omega}, E). \quad (5)$$

- ▶ Adjoint flux ≈ *relative importance some component of a forward source has on detector response*

Current Vector

- ▶ The *forward current vector* is defined

$$\vec{J}(\vec{r}, E) = \int_{4\pi} d\Omega \hat{\Omega} \psi(\vec{r}, \hat{\Omega}, E), \quad (6)$$

Current Vector

- ▶ The *forward current vector* is defined

$$\vec{J}(\vec{r}, E) = \int_{4\pi} d\Omega \hat{\Omega} \psi(\vec{r}, \hat{\Omega}, E), \quad (6)$$

- ▶ and the *adjoint current vector* is defined

$$\vec{J}^+(\vec{r}, E) = \int_{4\pi} d\Omega \hat{\Omega} \psi^+(\vec{r}, \hat{\Omega}, E). \quad (7)$$

Current Vector

- ▶ The *forward current vector* is defined

$$\vec{J}(\vec{r}, E) = \int_{4\pi} d\Omega \hat{\Omega} \psi(\vec{r}, \hat{\Omega}, E), \quad (6)$$

- ▶ and the *adjoint current vector* is defined

$$\vec{J}^+(\vec{r}, E) = \int_{4\pi} d\Omega \hat{\Omega} \psi^+(\vec{r}, \hat{\Omega}, E). \quad (7)$$

- ▶ Current represents *net flow* of particles at \vec{r} , a *vector quantity*



The Greedy Heuristic

The Adjoint Function

- ▶ For a region of interest (ROI) with adjoint flux $\phi_{ROI}^+(\vec{r})$, the *adjoint function* is defined

$$D_{ROI}^+(\vec{r}) = C_{1S_u} \phi_{ROI}^+(\vec{r}) / V_{ROI} \quad (8)$$

where C_{1S_u} yields units of Gy/S_u and S_u denotes one *source unit* (0.43 mCi for a ¹²⁵I seed) [4]



The Greedy Heuristic

The Adjoint Function

- ▶ For a region of interest (ROI) with adjoint flux $\phi_{ROI}^+(\vec{r})$, the *adjoint function* is defined

$$D_{ROI}^+(\vec{r}) = C_{1S_u} \phi_{ROI}^+(\vec{r}) / V_{ROI} \quad (8)$$

where C_{1S_u} yields units of Gy/S_u and S_u denotes one *source unit* (0.43 mCi for a ¹²⁵I seed) [4]

- ▶ The *adjoint ratio* is defined

$$\rho(\vec{r}) = \frac{w_u D_u^+(\vec{r}) + w_r D_r^+(\vec{r}) + w_n D_n^+(\vec{r})}{D_t^+(\vec{r})} \quad (9)$$

where w_i weight tissue importance; these are set to unity. [4]



The Greedy Heuristic

- ▶ Greedy algorithms: choose at each step the option that best satisfies some *greedy criterion*



The Greedy Heuristic

- ▶ Greedy algorithms: choose at each step the option that best satisfies some *greedy criterion*
- ▶ Hope solutions to each step (i.e. subproblems) lead to “global solution” ...or something like it



The Greedy Heuristic

- ▶ Greedy algorithms: choose at each step the option that best satisfies some *greedy criterion*
- ▶ Hope solutions to each step (i.e. subproblems) lead to “global solution” ...or something like it
- ▶ For brachytherapy, use *adjoint ratio* as greedy criterion



The Greedy Heuristic

The Greedy Heuristic

- ▶ Greedy algorithms: choose at each step the option that best satisfies some *greedy criterion*
- ▶ Hope solutions to each step (i.e. subproblems) lead to “global solution” ...or something like it
- ▶ For brachytherapy, use *adjoint ratio* as greedy criterion
- ▶ ...seed clustering? → *dose-update*

$$S_{dose}(j) = \frac{1}{D_p} \sum_{i=1}^n D(i,j) \quad (10)$$

$$\rho_d(j) = \rho(j) S_{dose}(j) \quad (11)$$

- ▶ $\rho_d(j)$ the *greedy criterion*



DANTSYS

- ▶ Diffusion Accelerated Neutral-particle Transport SYStem

DANTSYS

DANTSYS

- ▶ Diffusion Accelerated Neutral-particle Transport SYStem
- ▶ Discrete-ordinates: discretizes angular variable into n directions...

$$\phi(i, E) = \sum_{k=1}^n w_k \psi_{ik}(E) \quad (12)$$

$$J_x(i, E) = \sum_{k=1}^n \mu_k w_k \psi_{ik}(E) \quad (13)$$

DANTSYS

DANTSYS

- ▶ Diffusion Accelerated Neutral-particle Transport SYStem
- ▶ Discrete-ordinates: discretizes angular variable into n directions...

$$\phi(i, E) = \sum_{k=1}^n w_k \psi_{ik}(E) \quad (12)$$

$$J_x(i, E) = \sum_{k=1}^n \mu_k w_k \psi_{ik}(E) \quad (13)$$

- ▶ Discretizes space and energy variables... $\phi(i, g)$ for a voxel i and energy group g

DANTSYS

DANTSYS

- ▶ Diffusion Accelerated Neutral-particle Transport SYStem
- ▶ Discrete-ordinates: discretizes angular variable into n directions...

$$\phi(i, E) = \sum_{k=1}^n w_k \psi_{ik}(E) \quad (12)$$

$$J_x(i, E) = \sum_{k=1}^n \mu_k w_k \psi_{ik}(E) \quad (13)$$

- ▶ Discretizes space and energy variables... $\phi(i, g)$ for a voxel i and energy group g
- ▶ Lowest three energy groups of the FENDL-42 group photon library used



Models

Models

- ▶ *Prostate model* used based on set of three-dimensional ROI's used by Chaswal [1]



Models

Models

- ▶ *Prostate model* used based on set of three-dimensional ROI's used by Chaswal [1]
- ▶ Model spans $60 \text{ mm} \times 55 \text{ mm}$ in the plane contained in ultrasound images



Models

Models

- ▶ *Prostate model* used based on set of three-dimensional ROI's used by Chaswal [1]
- ▶ Model spans $60 \text{ mm} \times 55 \text{ mm}$ in the plane contained in ultrasound images
- ▶ 14 slices → slices 3, 5, 7, 9, 11 serve as two-dimensional models



Models

Models

- ▶ *Prostate model* used based on set of three-dimensional ROI's used by Chaswal [1]
- ▶ Model spans $60\text{ mm} \times 55\text{ mm}$ in the plane contained in ultrasound images
- ▶ 14 slices → slices 3, 5, 7, 9, 11 serve as two-dimensional models
- ▶ *Seed model* a simple photon source in a water medium → *no other materials modeled*



Models

Models

- ▶ *Prostate model* used based on set of three-dimensional ROI's used by Chaswal [1]
- ▶ Model spans $60\text{ mm} \times 55\text{ mm}$ in the plane contained in ultrasound images
- ▶ 14 slices → slices 3, 5, 7, 9, 11 serve as two-dimensional models
- ▶ *Seed model* a simple photon source in a water medium → *no other materials modeled*
- ▶ ICRP 21 flux-to-dose conversion factors (DF) used [2]



Theory

Introduction

- ▶ Adjoint function approach is not quite optimal



Theory

Introduction

- ▶ Adjoint function approach is not quite optimal
- ▶ Current vector: a more complete description of the physics...



Introduction

- ▶ Adjoint function approach is not quite optimal
- ▶ Current vector: a more complete description of the physics...
- ▶ ...may generate better insight to the physical nature of the problem → *better results*



Summing Adjoint Currents

- ▶ Define the horizontal (and vertical) summed tissue adjoint currents:

$$J_x^{tot}(i) = w_r J_x^r(i) + w_u J_x^u(i) + w_n J_x^n(i) + w_t J_x^t(i) \quad (14)$$

$$J_y^{tot}(i) = w_r J_y^r(i) + w_u J_y^u(i) + w_n J_y^n(i) + w_t J_y^t(i) \quad (15)$$

where $w_i = 1/V_i$ for the sensitive tissues i , but $-1/V_i$ for target

Summing Adjoint Currents

- ▶ Define the horizontal (and vertical) summed tissue adjoint currents:

$$J_x^{tot}(i) = w_r J_x^r(i) + w_u J_x^u(i) + w_n J_x^n(i) + w_t J_x^t(i) \quad (14)$$

$$J_y^{tot}(i) = w_r J_y^r(i) + w_u J_y^u(i) + w_n J_y^n(i) + w_t J_y^t(i) \quad (15)$$

where $w_i = 1/V_i$ for the sensitive tissues i , but $-1/V_i$ for target

- ▶ Write the adjoint current more succinctly:

$$\vec{J}^{tot}(i) = \hat{x} J_x^{tot}(i) + \hat{y} J_y^{tot}(i). \quad (16)$$

Summing Adjoint Currents

- ▶ Define the horizontal (and vertical) summed tissue adjoint currents:

$$J_x^{tot}(i) = w_r J_x^r(i) + w_u J_x^u(i) + w_n J_x^n(i) + w_t J_x^t(i) \quad (14)$$

$$J_y^{tot}(i) = w_r J_y^r(i) + w_u J_y^u(i) + w_n J_y^n(i) + w_t J_y^t(i) \quad (15)$$

where $w_i = 1/V_i$ for the sensitive tissues i , but $-1/V_i$ for target

- ▶ Write the adjoint current more succinctly:

$$\vec{J}^{tot}(i) = \hat{x} J_x^{tot}(i) + \hat{y} J_y^{tot}(i). \quad (16)$$

- ▶ Also have a *forward* current for seeds, \vec{J}^s



Theory

Defining Current-displacement

- ▶ From greedy heuristic, we have a given seed arrangement.

Theory

Defining Current-displacement

- ▶ From greedy heuristic, we have a given seed arrangement.
- ▶ Define the current-displacement vector,

$$\vec{C}(i) = a \left(-\vec{J}^{tot}(i) + b \sum_{\substack{j=1 \\ j \neq i}}^n \vec{J}^s(i,j) \right) \quad (17)$$

where $\vec{J}^s(i,j)$ is the current from a seed at i at a seed location j .

Defining Current-displacement

- ▶ From greedy heuristic, we have a given seed arrangement.
- ▶ Define the current-displacement vector,

$$\vec{C}(i) = a \left(-\vec{J}^{tot}(i) + b \sum_{\substack{j=1 \\ j \neq i}}^n \vec{J}^s(i,j) \right) \quad (17)$$

where $\vec{J}^s(i,j)$ is the current from a seed at i at a seed location j .

- ▶ b relates tissue and seed currents; a defines appropriate displacement

Defining Current-displacement

- ▶ From greedy heuristic, we have a given seed arrangement.
- ▶ Define the current-displacement vector,

$$\vec{C}(i) = a \left(-\vec{J}^{tot}(i) + b \sum_{\substack{j=1 \\ j \neq i}}^n \vec{J}^s(i,j) \right) \quad (17)$$

where $\vec{J}^s(i,j)$ is the current from a seed at i at a seed location j .

- ▶ b relates tissue and seed currents; a defines appropriate displacement
- ▶ This displacement is defined

$$i' = (x, y)' \longleftarrow (x + \vec{C}_x(i), y + \vec{C}_y(i)) = i + \vec{C}(i) \quad (18)$$



Theory

Dynamic Constants

- ▶ A set of static constants not “one size fits all”

Dynamic Constants

- ▶ A set of static constants not “one size fits all”
- ▶ Relating currents:

$$b = \max ||\vec{J}^{tot}||_2 / (k \max ||\vec{J}^s||_2), \quad (19)$$

$$k = 1 / \sqrt{m_1 \sqrt{m_2 / m_1}}. \quad (20)$$

$$m_1 = \text{mean}((J_x^s)^2 + (J_y^s)^2) \quad (21)$$

$$m_2 = \text{mean}((J_x^{tot})^2 + (J_y^{tot})^2). \quad (22)$$

Dynamic Constants

- ▶ A set of static constants not “one size fits all”
- ▶ Relating currents:

$$b = \max ||\vec{J}^{tot}||_2 / (k \max ||\vec{J}^s||_2), \quad (19)$$

$$k = 1 / \sqrt{m_1 \sqrt{m_2 / m_1}}. \quad (20)$$

$$m_1 = \text{mean}((J_x^s)^2 + (J_y^s)^2) \quad (21)$$

$$m_2 = \text{mean}((J_x^{tot})^2 + (J_y^{tot})^2). \quad (22)$$

- ▶ Defining displacement

$$a = k / \max ||\vec{J}^{tot}||_2 \quad (23)$$

Theory

More Details

- ▶ Multiple displacements, i.e. iterations: how many seeds? how many iterations?

More Details

- ▶ Multiple displacements, i.e. iterations: how many seeds? how many iterations?
- ▶ Rank seeds:

$$\|\vec{C}(i)\|_2 = \sqrt{C_x(i)^2 + C_y(i)^2} \quad (24)$$

More Details

- ▶ Multiple displacements, i.e. iterations: how many seeds? how many iterations?
- ▶ Rank seeds:

$$\|\vec{C}(i)\|_2 = \sqrt{C_x(i)^2 + C_y(i)^2} \quad (24)$$

- ▶ Diminish \vec{C} with each iteration:

$$\vec{C}(i) = \text{round}(\vec{C}(i)/it^c). \quad (25)$$

where “round” accounts for discretization

Results and Discussion

		$f = 1$	$f = 2$	$f = 3$	Total	% Δ
TARGET						
$V_{100}(\%)$	CD	98.08	98.87	99.39	98.78	-0.26
	GY	98.42	99.23	99.47	99.04	
$V_{150}(\%)$	CD	79.55	81.50	85.32	82.12	-1.44
	GY	80.20	82.87	86.89	83.32	
CN	CD	0.51	0.49	0.45	0.48	-2.30
	GY	0.50	0.50	0.45	0.49	
DNR	CD	0.81	0.82	0.86	0.83	-1.03
	GY	0.81	0.83	0.87	0.84	
URETHRA						
$V_{125,ur}(\%)$	CD	57.16	69.29	80.42	68.96	-8.40
	GY	67.73	75.81	82.32	75.29	
$V_{360,ur}(\%)$	CD	0.00	0.00	0.00	0.00	0.00
	GY	0.00	0.00	0.00	0.00	
$D_{90,ur}(Gy)$	CD	167.35	177.00	195.86	180.07	-1.81
	GY	170.95	180.90	198.33	183.39	

Table: Average evaluation parameters, by f and total.

Results and Discussion

		$f = 1$	$f = 2$	$f = 3$	Total	% Δ
RECTUM						
$V_{80,re}(\%)$	CD	26.53	38.70	64.86	43.36	-2.65
	GY	28.15	41.45	64.03	44.54	
$V_{90,re}(\%)$	CD	80.87	87.73	95.68	88.09	0.11
	GY	80.78	87.54	95.66	87.99	
$D_{90,re}(Gy)$	CD	86.40	90.07	102.98	93.15	0.13
	GY	86.47	90.41	102.21	93.03	
NORMAL TISSUE						
$D_{90,no}(Gy)$	CD	46.74	50.79	53.37	50.30	1.65
	GY	45.71	49.91	52.82	49.48	
QUALITATIVE						
Seeds	CD	19.60	10.40	7.60	12.53	0.00
	GY	19.60	10.40	7.60	12.53	

Table: Average evaluation parameters, by f and total.

Results and Discussion









Figure: DVH's and placement for slices 3, 5, and 7 with $f = 1$.

Results and Discussion







Figure: DVH's and placement for slices 9 and 11 with $f = 1$.



Results and Discussion

Key Points

- ▶ CD reduces V_{150} and DNR ; raises CN .



Results and Discussion

Key Points

- ▶ CD reduces V_{150} and DNR ; raises CN .
- ▶ V_{100} must be 98%; one case failed → adjust algorithm?

Results and Discussion

Key Points

- ▶ CD reduces V_{150} and DNR ; raises CN .
- ▶ V_{100} must be 98%; one case failed → adjust algorithm?
- ▶ Target irradiation with CD **slightly better**

Results and Discussion

Key Points

- ▶ CD reduces V_{150} and DNR ; raises CN .
- ▶ V_{100} must be 98%; one case failed → adjust algorithm?
- ▶ Target irradiation with CD **slightly better**
- ▶ All urethra parameters improve in *each case perturbed* → CD **improves urethra sparing**

Results and Discussion

Key Points

- ▶ CD reduces V_{150} and DNR ; raises CN .
- ▶ V_{100} must be 98%; one case failed → adjust algorithm?
- ▶ Target irradiation with CD **slightly better**
- ▶ All urethra parameters improve in *each case perturbed* → CD **improves urethra sparing**
- ▶ CD reduces $V_{80,re}$; very slightly increases other parameters → little change

Results and Discussion

Key Points

- ▶ CD reduces V_{150} and DNR ; raises CN .
- ▶ V_{100} must be 98%; one case failed → adjust algorithm?
- ▶ Target irradiation with CD **slightly better**
- ▶ All urethra parameters improve in *each case perturbed* → CD **improves urethra sparing**
- ▶ CD reduces $V_{80,re}$; very slightly increases other parameters → little change
- ▶ $D_{90,no}$ increased slightly by CD → **slightly reduced normal tissue sparing**

Results and Discussion

Key Points

- ▶ CD reduces V_{150} and DNR ; raises CN .
- ▶ V_{100} must be 98%; one case failed → adjust algorithm?
- ▶ Target irradiation with CD **slightly better**
- ▶ All urethra parameters improve in *each case perturbed* → CD **improves urethra sparing**
- ▶ CD reduces $V_{80,re}$; very slightly increases other parameters → little change
- ▶ $D_{90,no}$ increased slightly by CD → **slightly reduced normal tissue sparing**
- ▶ *Applying CD leads to better treatment plans on average*



Theory

Redefining Current-displacement

- ▶ Redefine current-displacement as:

$$\vec{C}(i) = (\hat{x} J_x^{tot}(i) + \hat{y} J_y^{tot}(i)) . \quad (26)$$

Redefining Current-displacement

- ▶ Redefine current-displacement as:

$$\vec{C}(i) = (\hat{x} J_x^{tot}(i) + \hat{y} J_y^{tot}(i)). \quad (26)$$

- ▶ Summed current from tissues at a voxel i is redefined

$$\vec{J}^{tot}(i) = w_r |\vec{J}^r(i)| + w_u |\vec{J}^u(i)| + w_n |\vec{J}^n(i)| + w_t |\vec{J}^t(i)|. \quad (27)$$

Redefining Current-displacement

- ▶ Redefine current-displacement as:

$$\vec{C}(i) = (\hat{x} J_x^{tot}(i) + \hat{y} J_y^{tot}(i)). \quad (26)$$

- ▶ Summed current from tissues at a voxel i is redefined

$$\vec{J}^{tot}(i) = w_r |\vec{J}^r(i)| + w_u |\vec{J}^u(i)| + w_n |\vec{J}^n(i)| + w_t |\vec{J}^t(i)|. \quad (27)$$

- ▶ a *new greedy criterion*

$$\|\vec{C}(i)\|_2 = \sqrt{C_x(i)^2 + C_y(i)^2} \quad (28)$$



Theory

Redefining Current-displacement cont.

- ▶ Need dynamic weights...

Redefining Current-displacement cont.

- ▶ Need dynamic weights...
- ▶ Define for a tissue i the parameter:

$$rms_i = \sqrt{\text{mean}\left((J_x^i)^2 + (J_y^i)^2\right)} \quad (29)$$

Redefining Current-displacement cont.

- ▶ Need dynamic weights...
- ▶ Define for a tissue i the parameter:

$$rms_i = \sqrt{\text{mean}\left((J_x^i)^2 + (J_y^i)^2\right)} \quad (29)$$

- ▶ Define weights:

$$w_i = 1/rms_i \div \min rms_i, i \in (u, r, n). \quad (30)$$

Redefining Current-displacement cont.

- ▶ Need dynamic weights...
- ▶ Define for a tissue i the parameter:

$$rms_i = \sqrt{\text{mean}\left((J_x^i)^2 + (J_y^i)^2\right)} \quad (29)$$

- ▶ Define weights:

$$w_i = 1/rms_i \div \min rms_i, i \in (u, r, n). \quad (30)$$

- ▶ Target not included!

Results and Discussion

		With Grid	Without Grid	Total	% Δ
TARGET					
$V_{100}(\%)$	<i>CD</i>	99.50	99.11	99.30	0.20
	<i>GY</i>	99.39	98.82	99.11	
$V_{150}(\%)$	<i>CD</i>	79.64	79.87	79.76	-1.68
	<i>GY</i>	80.70	81.54	81.12	
CN	<i>CD</i>	0.59	0.54	0.56	5.04
	<i>GY</i>	0.56	0.52	0.54	
DNR	<i>CD</i>	0.80	0.81	0.80	-1.77
	<i>GY</i>	0.81	0.82	0.82	
URETHRA					
$V_{125,ur}(\%)$	<i>CD</i>	72.35	57.03	64.69	-16.48
	<i>GY</i>	83.13	71.77	77.45	
$V_{360,ur}(\%)$	<i>CD</i>	0.00	0.00	0.00	0.00
	<i>GY</i>	0.00	0.00	0.00	
$D_{90,ur}(Gy)$	<i>CD</i>	178.86	171.95	175.41	-1.89
	<i>GY</i>	181.63	175.92	178.78	

Table: Average evaluation parameters for grid, no grid, and overall.

Results and Discussion

		With Grid	Without Grid	Total	% Δ
RECTUM					
$V_{80,re}(\%)$	<i>CD</i>	42.69	42.41	42.55	10.85
	<i>GY</i>	41.97	34.80	38.38	
$V_{90,re}(\%)$	<i>CD</i>	84.95	88.00	86.48	2.68
	<i>GY</i>	84.28	84.16	84.22	
$D_{90,re}(Gy)$	<i>CD</i>	90.73	91.73	91.23	2.10
	<i>GY</i>	90.27	88.44	89.35	
NORMAL TISSUE					
$D_{90,no}(Gy)$	<i>CD</i>	38.74	40.48	39.66	-14.12
	<i>GY</i>	44.55	47.81	46.18	
QUALITATIVE					
<i>Seeds</i>	<i>CD</i>	13.20	13.90	13.55	-7.19
	<i>GY</i>	14.20	15.00	14.60	

Table: Average evaluation parameters for grid, no grid, and overall.

Results and Discussion







Figure: DVH's and placement for slices 3 and 5 with $f = 1$ and grid.

Results and Discussion







Figure: DVH's and placement for slices 7 and 9 with $f = 1$ and grid.

Results and Discussion









Figure: DVH's and placement for slices 3, 5, and 7 with $f = 1$ and no grid.

Results and Discussion







Figure: DVH's and placement for slices 9 and 11 with $f = 1$ and no grid.

Results and Discussion

Key Points

- ▶ CD reduces both V_{150} and DNR and raises CN (13/20 cases, 2/20 unchanged)

Results and Discussion

Key Points

- ▶ CD reduces both V_{150} and DNR and raises CN (13/20 cases, 2/20 unchanged)
- ▶ V_{100} negligibly higher for CD

Results and Discussion

Key Points

- ▶ CD reduces both V_{150} and DNR and raises CN (13/20 cases, 2/20 unchanged)
- ▶ V_{100} negligibly higher for CD
- ▶ Target irradiation with CD **significantly better**

Results and Discussion

Key Points

- ▶ CD reduces both V_{150} and DNR and raises CN (13/20 cases, 2/20 unchanged)
- ▶ V_{100} negligibly higher for CD
- ▶ Target irradiation with CD significantly better
- ▶ Urethra parameters improve, more so without the grid → CD improves urethra sparing

Results and Discussion

Key Points

- ▶ CD reduces both V_{150} and DNR and raises CN (13/20 cases, 2/20 unchanged)
- ▶ V_{100} negligibly higher for CD
- ▶ Target irradiation with CD **significantly better**
- ▶ Urethra parameters improve, more so without the grid → CD **improves urethra sparing**
- ▶ Rectum parameters worsen, more so without the grid → CD yields **less rectum sparing**

Results and Discussion

Key Points

- ▶ CD reduces both V_{150} and DNR and raises CN (13/20 cases, 2/20 unchanged)
- ▶ V_{100} negligibly higher for CD
- ▶ Target irradiation with CD significantly better
- ▶ Urethra parameters improve, more so without the grid → CD improves urethra sparing
- ▶ Rectum parameters worsen, more so without the grid → CD yields less rectum sparing
- ▶ $D_{90,no}$ significantly decreased → CD produces better normal tissue sparing

Results and Discussion

Key Points

- ▶ CD reduces both V_{150} and DNR and raises CN (13/20 cases, 2/20 unchanged)
- ▶ V_{100} negligibly higher for CD
- ▶ Target irradiation with CD significantly better
- ▶ Urethra parameters improve, more so without the grid → CD improves urethra sparing
- ▶ Rectum parameters worsen, more so without the grid → CD yields less rectum sparing
- ▶ $D_{90,no}$ significantly decreased → CD produces better normal tissue sparing
- ▶ *CD as the greedy criterion leads to better treatment plans on average*

Concluding Remarks

- ▶ Two current-based approaches for brachytherapy optimization developed

Concluding Remarks

- ▶ Two current-based approaches for brachytherapy optimization developed
- ▶ CD as “nudge” → some better urethra, rectal sparing; less normal sparing

Concluding Remarks

- ▶ Two current-based approaches for brachytherapy optimization developed
- ▶ CD as “nudge” → some better urethra, rectal sparing; less normal sparing
- ▶ CD as the greedy criterion → substantial improvement in target coverage and urethra and normal tissue sparing

Concluding Remarks

- ▶ Two current-based approaches for brachytherapy optimization developed
- ▶ CD as “nudge” → some better urethra, rectal sparing; less normal sparing
- ▶ CD as the greedy criterion → substantial improvement in target coverage and urethra and normal tissue sparing
- ▶ *limits to 2-D treatment:* go to 3-D!

Concluding Remarks


- ▶ Two current-based approaches for brachytherapy optimization developed
- ▶ CD as “nudge” → some better urethra, rectal sparing; less normal sparing
- ▶ CD as the greedy criterion → substantial improvement in target coverage and urethra and normal tissue sparing
- ▶ *limits to 2-D treatment:* go to 3-D!
- ▶ **Questions?** *Jeremy Roberts, jaroberts2@wisc.edu*

Thanks!


References

-  V Chaswal.
Adjoint Based Treatment Planning for Brachytherapy: Novel Techniques and Further Developments.
PhD thesis, University of Wisconsin - Madison, 2008.
-  ICRP Committee 3 Task Group, P Grande, and M. C. O'riordant, chairman.
Data for Protection Against Ionizing Radiation from External Sources: Supplement to ICRP Publication 15.
Technical report, International Commission of Radiological Protection, APR 1971.
-  E. E. Lewis and W. F. Miller.
Computational Methods of Neutron Transport.
American Nuclear Society, 1984.
-  Sua Yoo, Michael E. Kowalok, Bruce R. Thomadsen, and Douglass L. Henderson.
Treatment planning for prostate brachytherapy using region of interest adjoint functions and a greedy heuristic.
Physics in Medicine and Biology, 48(24):4077–4090, DEC 21 2003.


The Adjoint Function - Plots




(a) Target adj. function.




(b) Urethra adj. func-



(c) Rectum adj. func-




(d) Normal tissue adj.




(e) Adjoint ratio.


Adjoint Current - Plots




(a) Target adjoint cur-
rent.



(b) Urethra adjoint cur-
rent.



(c) Rectum adjoint cur-
rent.



(d) Normal adjoint cur-