

You are given the following frequency distribution for 25 data points:

Class	Class intervals	Frequencies	Relative freq.	Cumulative freq.
1	[100; 200]	3		
2	(200; 300]	5		
3	(300; 400]	8		
4	(400; 500]	6		
5	(500; 600]	1		
6	(600; 700]	2		

- Suppose we are interested in constructing a **relative frequency polygon**. Provide the necessary coordinates.

x							
y							

- Give the coordinates for constructing an **ogive**.

x							
y							

You are given the following frequency distribution for 25 data points:

Class	Class intervals	Frequencies	Relative freq.	Cumulative freq.
1	[100; 200]	3	0.12	0.12
2	(200; 300]	5	0.20	0.32
3	(300; 400]	8	0.32	0.64
4	(400; 500]	6	0.24	0.88
5	(500; 600]	1	0.04	0.92
6	(600; 700]	2	0.08	1


1. Suppose we are interested in constructing a **relative frequency polygon**. Provide the necessary coordinates.

x	50	150	250	350	450	550	650
y	0	0.12	0.20	0.32	0.24	0.04	0.08

2. Give the coordinates for constructing an **ogive**.

x	100	200	300	400	500	600	700
y	0	0.12	0.32	0.64	0.88	0.92	1

Relative frequency polygon



Ogive

