

# Equilibrium states on the Toeplitz-Cuntz-Krieger algebras of finite graphs

Astrid an Huef

University of Otago

Operator Algebra Day  
Institute of Mathematics, University of Aberdeen  
13 May 2013

This talk contains some results obtained in



A. an Huef, M. Laca, I. Raeburn and A. Sims, KMS states  
on  $C^*$ -algebras of finite graphs, *J. Math. Anal. Appl.*, 2013.

Let  $\alpha : \mathbb{R} \rightarrow \text{Aut } A$  be an action of the real line  $\mathbb{R}$  on a  $C^*$ -algebra  $A$ . (Today  $A$  always has an identity.)

In physical models, observables of the system are represented by self-adjoint elements of  $A$ , and states of the system by positive functionals of norm 1 on  $A$ :  $\phi(a)$  is the expected value of the observable  $a$  in the state  $\phi$  (which is real because  $a = a^*$  and  $\phi \geq 0$ ).

The action  $\alpha$  represents the time evolution of the system: the observable  $a$  at time 0 moves to  $\alpha_t(a)$  at time  $t$ , or the state  $\phi$  at time 0 moves to  $\phi \circ \alpha_t$ .

In statistical physics, an important role is played by *equilibrium states*, which are in particular invariant under the time evolution. In  $C^*$ -algebraic models equilibrium states are called *KMS states*, after Kubo, Martin and Schwinger.

Let  $\alpha : \mathbb{R} \rightarrow \text{Aut } A$  be an action. Then  $a \in A$  is an *analytic element* if the function  $t \mapsto \alpha_t(a)$  from  $\mathbb{R}$  to  $A$  has an extension to an entire function on  $\mathbb{C}$ . The set of analytic elements is always a dense subalgebra of  $A$ . A state  $\phi$  on  $A$  is a *KMS state at inverse temperature  $\beta$*  if

$$\phi(ab) = \phi(b\alpha_{i\beta}(a)) \text{ for all analytic } a, b.$$

- ▶ KMS states are  $\alpha$ -invariant.
- ▶ It suffices to check the  $\text{KMS}_\beta$  condition on a set of analytic elements which span a dense subspace of  $A$ .
- ▶ The  $\text{KMS}_\beta$  states always form a simplex.
- ▶ In a physical model we expect KMS states for most  $\beta$ .

Suppose that  $E = (E^0, E^1, r, s)$  is a directed graph. Today it is always finite. A *Cuntz-Krieger  $E$ -family* consists of mutually orthogonal projections  $\{P_v : v \in E^0\}$  and partial isometries  $\{S_e : e \in E^1\}$  satisfying  $S_e^* S_e = P_{s(e)}$  and

$$P_v = \sum_{r(e)=v} S_e S_e^* \quad \text{whenever } v \text{ is not a source.}$$

The *graph algebra* is a  $C^*$ -algebra  $C^*(E)$  which is generated by a universal Cuntz-Krieger family  $\{p, s\}$ .

Each graph algebra carries a *gauge action*  $\gamma : \mathbb{T} \rightarrow \text{Aut } C^*(E)$  characterised by  $\gamma_z(s_e) = z s_e$  and  $\gamma_z(p_v) = p_v$ . This lifts to an action of  $\mathbb{R}$  such that  $\alpha_t(s_e) = e^{it} s_e$  and  $\alpha_t(p_v) = p_v$ .

We are going to compute the KMS states of  $(C^*(E), \alpha)$ .

The Cuntz-Krieger relation  $p_\nu = \sum_{r(e)=\nu} s_e s_e^*$  and the orthogonality of the  $p_\nu$  imply that the projections  $\{s_e s_e^* : e \in E^1\}$  are mutually orthogonal: algebraically,  $s_e^* s_f = 0$  whenever  $e \neq f$ .

Thus  $C^*(E) = C^*(s, p)$  is spanned by the elements

$$s_\mu s_\nu^* := s_{\mu_1} s_{\mu_2} \cdots s_{\mu_{|\mu|}} s_{\nu_{|\nu|}}^* \cdots s_{\nu_1}^*$$

for paths  $\mu$  and  $\nu$  in  $E$ : we write  $\mu, \nu \in E^*$ . Crucial for us is that

$$t \mapsto \alpha_t(s_\mu s_\nu^*) = e^{it(|\mu| - |\nu|)} s_\mu s_\nu^*$$

extends to an analytic function (just replace  $t$  by  $z$ ).

Let  $\phi$  be a  $\text{KMS}_\beta$  state on  $(C^*(E), \alpha)$ . We have

$$\begin{aligned}\phi(s_\mu s_\nu^*) &= \phi(s_\nu^* \alpha_{i\beta}(s_\mu)) = e^{-\beta|\mu|} \phi(s_\nu^* s_\mu) \\ &= e^{-\beta(|\mu| - |\nu|)} \phi(s_\mu s_\nu^*).\end{aligned}$$

Thus  $\phi(s_\mu s_\nu^*) = 0$  unless  $|\mu| = |\nu|$ , and then  $s_\nu^* s_\mu = 0$  unless  $\mu = \nu$ .

**Lemma.** A state  $\phi$  on  $C^*(E)$  is  $\text{KMS}_\beta$  for  $\alpha$  if and only if

$$\phi(s_\mu s_\nu^*) = \begin{cases} 0 & \text{when } \nu \neq \mu \\ e^{-\beta|\mu|} \phi(p_{s(\mu)}) & \text{when } \nu = \mu. \end{cases}$$

We haven't used the Cuntz-Krieger relation yet:

Let  $\phi$  be a  $\text{KMS}_\beta$  state on  $(C^*(E), \alpha)$ . Suppose  $v \in E^0$  is not a source. Then

$$\phi(p_v) = \sum_{r(e)=v} \phi(s_e s_e^*) = \sum_{r(e)=v} e^{-\beta} \phi(p_{s(e)}).$$

The **vertex matrix** of  $E$  is the  $E^0 \times E^0$  integer matrix  $A$  with entries  $A(v, w) = |r^{-1}(v) \cap s^{-1}(w)|$ . We can rearrange the above sum as

$$e^\beta \phi(p_v) = \sum_{w \in E^0} \sum_{r(e)=v, s(e)=w} \phi(p_w) = \sum_{w \in E^0} A(v, w) \phi(p_w). \quad (1)$$

So if  $E$  has no sources, then the vector  $m = (m_v) := (\phi(p_v)) \in [0, \infty)^{E^0}$  satisfies  $Am = e^\beta m$ .

If  $E$  is strongly connected, then  $A$  is irreducible, and  $e^\beta$  has to be the **Perron-Frobenius** eigenvalue of  $A$ . Since

$1 = \phi(1) = \sum_v \phi(p_v) = \sum_v m_v$ , the vector  $m$  is the unique PF eigenvector with  $\|m\|_1 = 1$ .

**Theorem (Enomoto-Fujii-Watatani 1984).** Let  $E$  be a strongly connected finite graph with vertex matrix  $A$ . Then  $(C^*(E), \alpha)$  has a unique KMS state. This state has inverse temperature  $\beta = \ln \rho(A)$ , where  $\rho(A)$  is the spectral radius of  $A$ .

We have shown there is at most one, so we need to show existence. The easiest way to construct the state is to use the idea from Exel-Laca (2003), Laca-Neshveyev (2004): the Toeplitz algebra  $\mathcal{TC}^*(E)$  of  $E$  has a much richer supply of KMS states.

$\mathcal{TC}^*(E)$  is generated by a universal Toeplitz-Cuntz-Krieger family of mutually orthogonal projections  $\{q_v\}$  and partial isometries  $\{t_e\}$  such that  $t_e^* t_e = q_{s(e)}$  and, if  $v$  is not a source,

$$q_v \geq \sum_{e \in F} t_e t_e^* \quad \text{for } F \subset r^{-1}(v).$$

Again  $\mathcal{TC}^*(E) = \overline{\text{span}}\{t_\mu t_\nu^* : \mu, \nu \in E^*, s(\mu) = s(\nu)\}$  and there is a gauge action  $\gamma : \mathbb{T} \rightarrow \text{Aut}(\mathcal{TC}^*(E))$  satisfying  $\gamma_z(t_e) = z t_e$  and  $\gamma_z(q_v) = q_v$ , which we can lift to  $\mathbb{R}$ .



Let  $I$  be the ideal of  $\mathcal{TC}^*(E)$  generated by

$$\{q_v - \sum_{r(e)=v} t_e t_e^* : v \text{ is not a source}\}.$$

Then  $C^*(E) \cong \mathcal{TC}^*(E)/I$ .

It is again easy to recognise a  $\text{KMS}_\beta$  state  $\phi$  on  $(\mathcal{TC}^*(E), \alpha)$ , and the formulas look the same.

But now  $m = (m_v) := (\phi(q_v))$  is a unit vector in  $\ell^1(E^0)$  satisfying  $Am \leq e^\beta m$ .

PF theory says for  $A$  irreducible:

- ▶  $Am = e^\beta m \implies e^\beta = \rho(A) \implies \beta = \ln \rho(A)$ ;
- ▶  $Am \leq e^\beta m$  and  $\beta = \ln \rho(A) \implies m$  is the PF eigenvector;
- ▶  $Am \leq e^\beta m$  and  $Am \neq e^\beta m \implies \beta > \ln \rho(A)$ .

Note  $Am \leq e^\beta m \iff (I - e^{-\beta}A)m \geq 0$ , and consider  $\beta > \ln \rho(A)$ . Then  $\sum_{n=0}^{\infty} e^{-\beta n} A^n$  converges to  $(1 - e^{-\beta}A)^{-1}$ .

Take  $\epsilon := (1 - e^{-\beta}A)m$ . Which  $\epsilon \in [0, \infty]^{E^0}$  arise?

For  $v \in E^0$ , set

$$y_v := \sum_{n=0}^{\infty} \sum_{w \in E^0} e^{-\beta n} A^n(w, v)$$

(which again converges because  $e^\beta > \rho(A)$ ), and take  $y = (y_v)$ . Then:

**Lemma.** Let  $\beta > \ln \rho(A)$ . Then  $m := (1 - e^{-\beta}A)^{-1}\epsilon$  is a unit vector in  $\ell^1(E^0)$  satisfying  $Am \leq e^\beta m$  if and only if  $\epsilon \cdot y = 1$ .

We now know that the KMS condition on a state  $\phi$  places restraints on  $m := (\phi(p_v))$ . We still need to construct KMS states. We need a concrete representation of  $\mathcal{TC}^*(E)$ :

**Example.** Consider the usual orthonormal basis  $\{h_\mu : \mu \in E^*\}$  for  $\ell^2(E^*)$  (by convention  $E^0 \subset E^*$ ). There are projections  $Q_v$  and partial isometries  $T_e$  on  $\ell^2(E^*)$  such that

$$Q_v h_\mu = \begin{cases} 0 & \text{unless } r(\mu) = v \\ h_\mu & \text{if } r(\mu) = v, \text{ and} \end{cases}$$

$$T_e h_\mu = \begin{cases} 0 & \text{unless } r(\mu) = s(e) \\ h_{e\mu} & \text{if } r(\mu) = s(e). \end{cases}$$

Then  $(Q, T)$  is a Toeplitz-CK family, and we have a representation  $\pi_{Q,T}$  of  $\mathcal{TC}^*(E)$  on  $\ell^2(E^*)$  (in fact injective).

**Theorem (an Huef-Laca-Raeburn-Sims, 2013).** Suppose  $E$  is a finite graph with vertex matrix  $A$ , and  $\beta > \ln \rho(A)$ . Take  $y = (y_v) \in [1, \infty)^{E^0}$  as above, and suppose  $\epsilon \cdot y = 1$ . Then there is a  $\text{KMS}_\beta$  state  $\phi_\epsilon$  of  $\mathcal{TC}^*(E)$  such that

$$\phi_\epsilon(a) = \sum_{\mu \in E^*} e^{-\beta|\mu|} \epsilon_{s(\mu)} (\pi_{Q,T}(a) h_\mu \mid h_\mu).$$

The map  $\epsilon \mapsto \phi_\epsilon$  is an affine isomorphism of  $\Delta_\beta = \{\epsilon \in [0, 1]^{E^0} : \epsilon \cdot y = 1\}$  onto the simplex of  $\text{KMS}_\beta$  states.

Notice there is no hypothesis on  $E$ , hence no irreducibility assumption on  $A$ . So what happens at  $\beta = \ln \rho(A)$ ? When  $A$  is irreducible, the series defining  $y$  diverges, so the simplex  $\Delta_\beta$  contracts to  $\{0\}$  as  $\beta \rightarrow \ln \rho(A)+$ .

**Corollary (Enomoto-Fujii-Watatani).** If  $E$  is strongly connected, then  $(C^*(E), \alpha)$  has a  $\text{KMS}_{\ln \rho(A)}$  state.

**Proof.** Choose  $\beta_n$  decreasing to  $\ln \rho(A)$ , and  $\text{KMS}_{\beta_n}$  states  $\phi_n$  of  $\mathcal{TC}^*(E)$ . By passing to a subsequence,  $\phi_n \rightarrow \phi$ , and  $\phi$  is a  $\text{KMS}_{\ln \rho(A)}$  state of  $\mathcal{TC}^*(E)$ . Then  $m := (\phi(q_v))$  satisfies  $Am \leq \rho(A)m$ . PF implies  $Am = \rho(A)m$ . Thus

$$\begin{aligned} \rho(A)\phi(q_v) &= \rho(A)m_v = (Am)_v = \sum_{w \in E^0} A(v, w)\phi(q_w) \\ &= \sum_{r(e)=v} \phi(q_{s(e)}) = \sum_{r(e)=v} \rho(A)\phi(t_e t_e^*) \\ &= \rho(A)\phi\left(\sum_{r(e)=v} t_e t_e^*\right). \end{aligned}$$









So for all  $v \in E^0$  which are not sources,

$$\phi\left(q_v - \sum_{r(e)=v} t_e t_e^*\right) = 0.$$

Now a technical lemma implies that  $\phi$  factors through  $C^*(E) = \mathcal{TC}^*(E)/I$ .

This completes the proof of:

**Theorem (Enomoto-Fujii-Watatani 1984).** Let  $E$  be a strongly connected finite graph with vertex matrix  $A$ . Then  $(C^*(E), \alpha)$  has a unique KMS state. This state has inverse temperature  $\beta = \ln \rho(A)$ , where  $\rho(A)$  is the spectral radius of  $A$ .

-  T. Carlsen and N. Larsen, work in progress.
-  M. Enomoto, M. Fujii and Y. Watatani, KMS states for gauge action on  $\mathcal{O}_A$ , *Math. Japon.* **29** (1984).
-  R. Exel and M. Laca, Partial dynamical systems and the KMS condition, *Comm. Math. Phys.* **232** (2003).
-  N.J. Fowler and I. Raeburn, The Toeplitz algebra of a Hilbert bimodule, *Indiana Univ. Math. J.* **48** (1999).
-  A. an Huef, M. Laca, I. Raeburn and A. Sims, KMS states on  $C^*$ -algebras of finite graphs, *J. Math. Anal. Appl.*
-  A. an Huef, M. Laca, I. Raeburn and A. Sims, KMS states on the  $C^*$ -algebras associated to higher-rank graphs, arXiv.
-  T. Kajiwara and Y. Watatani, KMS states on finite-graph  $C^*$ -algebras, arXiv.
-  M. Laca and S. Neshveyev, KMS states of quasi-free dynamics on Pimsner algebras, *J. Funct. Anal.* **211** (2004), 457–482.