

Topological dimension of C^* -algebras

Aaron Tikuisis

`a.tikuisis@abdn.ac.uk`

University of Aberdeen

The prototypical C^* -algebra:

$$\mathcal{B}(\mathcal{H}) := \{\text{continuous (=operator norm-bounded) linear operators on } \mathcal{H}\},$$

where \mathcal{H} is a complex Hilbert space.

Definition

A C^* -algebra is a norm-closed $*$ -subalgebra of $\mathcal{B}(\mathcal{H})$.

(C^* -algebras can also be defined abstractly, as Banach $*$ -algebras satisfying $\|a^*a\| = \|a\|^2$ for all a .)

The prototypical C^* -algebra:

$$\mathcal{B}(\mathcal{H}) := \{\text{continuous (=operator norm-bounded) linear operators on } \mathcal{H}\},$$

where \mathcal{H} is a complex Hilbert space.

Definition

A C^* -algebra is a norm-closed $*$ -subalgebra of $\mathcal{B}(\mathcal{H})$.

(C^* -algebras can also be defined abstractly, as Banach $*$ -algebras satisfying $\|a^*a\| = \|a\|^2$ for all a .)

The prototypical C^* -algebra:

$$\mathcal{B}(\mathcal{H}) := \{\text{continuous (=operator norm-bounded) linear operators on } \mathcal{H}\},$$

where \mathcal{H} is a complex Hilbert space.

Definition

A C^* -algebra is a norm-closed $*$ -subalgebra of $\mathcal{B}(\mathcal{H})$.

(C^* -algebras can also be defined abstractly, as Banach $*$ -algebras satisfying $\|a^*a\| = \|a\|^2$ for all a .)

The prototypical C^* -algebra:

$$\mathcal{B}(\mathcal{H}) := \{\text{continuous (=operator norm-bounded) linear operators on } \mathcal{H}\},$$

where \mathcal{H} is a complex Hilbert space.

Definition

A C^* -algebra is a norm-closed $*$ -subalgebra of $\mathcal{B}(\mathcal{H})$.

(C^* -algebras can also be defined abstractly, as Banach $*$ -algebras satisfying $\|a^*a\| = \|a\|^2$ for all a .)

Examples of C^* -algebras

$$M_n(\mathbb{C}) = \mathcal{B}(\mathcal{H}) \text{ for } \mathcal{H} = \mathbb{C}^n$$

$C_0(X, \mathbb{C})$ where X is a locally compact Hausdorff topological space

$$C_0(X, M_n(\mathbb{C}))$$

Examples of C^* -algebras

$$M_n(\mathbb{C}) = \mathcal{B}(\mathcal{H}) \text{ for } \mathcal{H} = \mathbb{C}^n$$

$C_0(X, \mathbb{C})$ where X is a locally compact Hausdorff topological space

$$C_0(X, M_n(\mathbb{C}))$$

Examples of C^* -algebras

$$M_n(\mathbb{C}) = \mathcal{B}(\mathcal{H}) \text{ for } \mathcal{H} = \mathbb{C}^n$$

$C_0(X, \mathbb{C})$ where X is a locally compact Hausdorff topological space

$$C_0(X, M_n(\mathbb{C}))$$

Examples of C^* -algebras

$$M_n(\mathbb{C}) = \mathcal{B}(\mathcal{H}) \text{ for } \mathcal{H} = \mathbb{C}^n$$

$C_0(X, \mathbb{C})$ where X is a locally compact Hausdorff topological space

$$C_0(X, M_n(\mathbb{C}))$$

Constructions of C^* -algebras

C^* -algebras can be constructed from:

groups (via unitary representations)

coarse metric spaces

rings

graphs

topological dynamical systems (encoding orbit equivalence)

Constructions of C^* -algebras

C^* -algebras can be constructed from:

groups (via unitary representations)

coarse metric spaces

rings

graphs

topological dynamical systems (encoding orbit equivalence)

Constructions of C^* -algebras

C^* -algebras can be constructed from:

groups (via unitary representations)

coarse metric spaces

rings

graphs

topological dynamical systems (encoding orbit equivalence)

Constructions of C^* -algebras

C^* -algebras can be constructed from:

groups (via unitary representations)

coarse metric spaces

rings

graphs

topological dynamical systems (encoding orbit equivalence)

Constructions of C^* -algebras

C^* -algebras can be constructed from:

groups (via unitary representations)

coarse metric spaces

rings

graphs

topological dynamical systems (encoding orbit equivalence)

Constructions of C^* -algebras

C^* -algebras can be constructed from:

groups (via unitary representations)

coarse metric spaces

rings

graphs

topological dynamical systems (encoding orbit equivalence)

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:

..

Orientation-preserving local isometries of the Cantor set

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:



Orientation-preserving local isometries of the Cantor set

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:



Orientation-preserving local isometries of the Cantor set

Constructions of C^* -algebras


Idea: encode input data as continuous operators. Example:



Orientation-preserving local isometries of the Cantor set


Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:



Constructions of C^* -algebras


Idea: encode input data as continuous operators. Example:


$$\left(\begin{array}{c|c} \text{blue circle} & \\ \hline & \text{red circle} \end{array} \right) = \left(\begin{array}{c|c} \mathbb{C} & \mathbb{C} \\ \hline \mathbb{C} & \mathbb{C} \end{array} \right)$$

$M_2(\mathbb{C})$


Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:



Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:



$$M_4(\mathbb{C})$$

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:

$$\begin{pmatrix}
 \begin{array}{|c|c|} \hline \text{CNOT} & \text{SWAP} \\ \hline \end{array} & \begin{array}{|c|c|} \hline \text{CNOT} & \text{SWAP} \\ \hline \end{array} \\ \hline
 \begin{array}{|c|c|} \hline \text{CNOT} & \text{SWAP} \\ \hline \end{array} & \begin{array}{|c|c|} \hline \text{CNOT} & \text{SWAP} \\ \hline \end{array}
 \end{pmatrix} = \begin{pmatrix}
 \begin{array}{|c|c|} \hline \text{CNOT} \otimes \text{CNOT} & \text{CNOT} \otimes \text{CNOT} \\ \hline \end{array} & \begin{array}{|c|c|} \hline \text{CNOT} \otimes \text{CNOT} & \text{CNOT} \otimes \text{CNOT} \\ \hline \end{array} \\ \hline
 \begin{array}{|c|c|} \hline \text{CNOT} \otimes \text{CNOT} & \text{CNOT} \otimes \text{CNOT} \\ \hline \end{array} & \begin{array}{|c|c|} \hline \text{CNOT} \otimes \text{CNOT} & \text{CNOT} \otimes \text{CNOT} \\ \hline \end{array}
 \end{pmatrix}$$

$$M_8(\mathbb{C})$$

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:

$$\begin{array}{c} \dots \dots \dots \dots \end{array} \left(\begin{array}{c|c} \begin{array}{c|c} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \hline \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} \right) = \overline{\begin{array}{c} \dots \dots \dots \dots \end{array} \left(\begin{array}{c|c} \begin{array}{c|c} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \hline \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} \right)} \quad || \cdot ||$$

M_{2^∞}

“Uniformly hyperfinite” (UHF) algebra.

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:

$$\begin{array}{c} \dots \dots \end{array} \left(\begin{array}{c|c} \begin{array}{c|c} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \hline \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} \right) = \overline{\begin{array}{c} \dots \dots \end{array} \left(\begin{array}{c|c} \begin{array}{c|c} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \hline \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} \right)} \quad || \cdot ||$$

$$M_{2\infty}$$

“Uniformly hyperfinite” (UHF) algebra.
Can likewise define $M_{p\infty}$ for any $p \in \mathbb{N}$.

Constructions of C^* -algebras

Idea: encode input data as continuous operators. Example:

$$\begin{array}{c} \dots \dots \dots \dots \end{array} \left(\begin{array}{c|c} \begin{array}{c|c} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \hline \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} \right) = \overline{\left(\begin{array}{c|c} \begin{array}{c|c} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \hline \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} & \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \end{array} \right)} \quad || \cdot ||$$

“Uniformly hyperfinite” (UHF) algebra.

Can likewise define M_{p^∞} for any $p \in \mathbb{N}$.

Note: closure in a weaker topology gives a famous von Neumann algebra \mathcal{R} , the “hyperfinite II_1 -factor.”

There is only one Cantor set, but:

Glimm (1959)

For p, q prime, $M_{p^\infty} \cong M_{q^\infty}$ if and only if $p = q$.

Note for later: $M_{p^\infty} \otimes M_{p^\infty} \cong M_{p^\infty}$.

There is only one Cantor set, but:

Glimm (1959)

For p, q prime, $M_{p^\infty} \cong M_{q^\infty}$ if and only if $p = q$.


Note for later: $M_{p^\infty} \otimes M_{p^\infty} \cong M_{p^\infty}$.

Amenability

“Amenability” for C^* -algebras is characterised by a finite approximation property:

Definition

A C^* -algebra A is **amenable** if there is a sequence (or net) of **approximately** $^\Delta$ commuting diagrams



where F_n is **finite dimensional**.

Requiring $*$ -homomorphisms would be too strong. Instead, we require ϕ_n, ψ_n to be “completely positive contractions” (maps that order structure).


$^\Delta$: approximately means $\|\psi_n(\phi_n(a)) - a\| \xrightarrow{n \rightarrow \infty} 0 \forall a \in A$.

Amenability

“Amenability” for C^* -algebras is characterised by a finite approximation property:

Definition

A C^* -algebra A is **amenable** if there is a sequence (or net) of **approximately** $^\Delta$ commuting diagrams



where F_n is **finite dimensional**.

Requiring $*$ -homomorphisms would be too strong. Instead, we require ϕ_n, ψ_n to be “completely positive contractions” (maps that order structure).


$^\Delta$: approximately means $\|\psi_n(\phi_n(a)) - a\| \xrightarrow{n \rightarrow \infty} 0 \forall a \in A$.

Amenability

“Amenability” for C^* -algebras is characterised by a finite approximation property:

Definition

A C^* -algebra A is **amenable** if there is a sequence (or net) of **approximately** $^\Delta$ commuting diagrams



where F_n is **finite dimensional**.

Requiring $*$ -homomorphisms would be too strong.

Instead, we require ϕ_n, ψ_n to be “completely positive contractions” (maps that order structure).


$^\Delta$: approximately means $\|\psi_n(\phi_n(a)) - a\| \xrightarrow{n \rightarrow \infty} 0 \ \forall a \in A$.

Amenability

“Amenability” for C^* -algebras is characterised by a finite approximation property:

Definition

A C^* -algebra A is **amenable** if there is a sequence (or net) of **approximately** $^\Delta$ commuting diagrams



where F_n is **finite dimensional**.

Requiring $*$ -homomorphisms would be too strong. Instead, we require ϕ_n, ψ_n to be “completely positive contractions” (maps that order structure).


$^\Delta$: approximately means $\|\psi_n(\phi_n(a)) - a\| \xrightarrow{n \rightarrow \infty} 0 \forall a \in A$.

Amenability

“Amenability” for C^* -algebras is characterised by a finite approximation property:

Definition

A C^* -algebra A is **amenable** if there is a sequence (or net) of **approximately** $^\triangle$ commuting diagrams



where F_n is **finite dimensional**.

Requiring $*$ -homomorphisms would be too strong. Instead, we require ϕ_n, ψ_n to be “completely positive contractions” (maps that order structure).

$^\triangle$: approximately means $\|\psi_n(\phi_n(a)) - a\| \xrightarrow{n \rightarrow \infty} 0 \ \forall a \in A$.

Covering dimension

From Lebesgue, we have the following fruitful notion.

Definition

The **covering dimension** of a normal topological space X is the least number d such that, every finite open cover \mathcal{U} of X has finite refinement \mathcal{V} that can be coloured with $(d + 1)$ colours.

Covering dimension

From Lebesgue, we have the following fruitful notion.

Definition


The **covering dimension** of a normal topological space X is the least number d such that, every finite open cover \mathcal{U} of X has finite refinement \mathcal{V} that can be coloured with $(d + 1)$ colours.

Covering dimension

From Lebesgue, we have the following fruitful notion.

Definition

The **covering dimension** of a normal topological space X is the least number d such that, every finite open cover \mathcal{U} of X has finite refinement \mathcal{V} that can be coloured with $(d + 1)$ colours.



Covering dimension

From Lebesgue, we have the following fruitful notion.

Definition

The **covering dimension** of a normal topological space X is the least number d such that, every finite open cover \mathcal{U} of X has finite refinement \mathcal{V} that can be coloured with $(d + 1)$ colours.

(Warning: this is a minor, but equivalent, variation on the standard definition.)

Covering dimension

From Lebesgue, we have the following fruitful notion.

Definition

The **covering dimension** of a normal topological space X is the least number d such that, every finite open cover \mathcal{U} of X has finite refinement \mathcal{V} that can be coloured with $(d + 1)$ colours.

(Warning: this is a minor, but equivalent, variation on the standard definition.)


Highly desirable to extend dimension to all C^* -algebras (from the class $\{C_0(X, \mathbb{C})\}$ of abelian C^* -algebras).

Nuclear dimension

Nuclear dimension marries the two previous concepts: amenability and covering dimension.

Definition (Kirchberg–Winter '04, Winter–Zacharias '10)

The **nuclear dimension** of a C^* -algebra A is the least integer d such that there is a sequence (or net) of approximately commuting diagrams



such that ψ_n is a completely positive contraction **and** (F_n, ϕ_n) **can be coloured with $d + 1$ colours.**


$d + 1$ -colourable: F_n decomposes as a direct sum $F_n^{(0)} \oplus \cdots \oplus F_n^{(d)}$ and $\phi_n|_{F_n^{(d)}}$ preserves orthogonality.

Nuclear dimension

Nuclear dimension marries the two previous concepts: amenability and covering dimension.

Definition (Kirchberg–Winter '04, Winter–Zacharias '10)

The **nuclear dimension** of a C^* -algebra A is the least integer d such that there is a sequence (or net) of approximately commuting diagrams



such that ψ_n is a completely positive contraction **and** (F_n, ϕ_n) can be coloured with $d + 1$ colours.


$d + 1$ -colourable: F_n decomposes as a direct sum $F_n^{(0)} \oplus \cdots \oplus F_n^{(d)}$ and $\phi_n|_{F_n^{(d)}}$ preserves orthogonality.

Nuclear dimension

Nuclear dimension marries the two previous concepts: amenability and covering dimension.

Definition (Kirchberg–Winter '04, Winter–Zacharias '10)

The **nuclear dimension** of a C^* -algebra A is the least integer d such that there is a sequence (or net) of approximately commuting diagrams



such that ψ_n is a completely positive contraction **and** (F_n, ϕ_n) **can be coloured with $d + 1$ colours.**


$d + 1$ -colourable: F_n decomposes as a direct sum $F_n^{(0)} \oplus \cdots \oplus F_n^{(d)}$ and $\phi_n|_{F_n^{(d)}}$ preserves orthogonality.

Nuclear dimension

Nuclear dimension marries the two previous concepts: amenability and covering dimension.

Definition (Kirchberg–Winter '04, Winter–Zacharias '10)

The **nuclear dimension** of a C^* -algebra A is the least integer d such that there is a sequence (or net) of approximately commuting diagrams



such that ψ_n is a completely positive contraction **and** (F_n, ϕ_n) **can be coloured with $d + 1$ colours.**


$d + 1$ -colourable: F_n decomposes as a direct sum $F_n^{(0)} \oplus \cdots \oplus F_n^{(d)}$ and $\phi_n|_{F_n^{(d)}}$ preserves orthogonality.

Nuclear dimension

Nuclear dimension marries the two previous concepts: amenability and covering dimension.

Definition (Kirchberg–Winter '04, Winter–Zacharias '10)

The **nuclear dimension** of a C^* -algebra A is the least integer d such that there is a sequence (or net) of approximately commuting diagrams




such that ψ_n is a completely positive contraction **and** (F_n, ϕ_n) **can be coloured with $d + 1$ colours.**

$d + 1$ -colourable: F_n decomposes as a direct sum $F_n^{(0)} \oplus \cdots \oplus F_n^{(d)}$ and $\phi_n|_{F_n^{(d)}}$ preserves orthogonality.

Nuclear dimension: properties

Definition

$\dim_{\text{nuc}} A \leq d$ if there exist approximately commuting diagrams



such that ψ_n is a completely positive contraction and (F_n, ϕ_n) can be coloured with $(d + 1)$ colours.

For a compact metrisable space X ,


$$\dim_{\text{nuc}} C(X, \mathbb{C}) = \dim X = \dim_{\text{nuc}} C(X, M_p(\mathbb{C}))$$

If A is simple and $\dim_{\text{nuc}} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

Nuclear dimension: properties

Definition

$\dim_{\text{nuc}} A \leq d$ if there exist approximately commuting diagrams



such that ψ_n is a completely positive contraction and (F_n, ϕ_n) can be coloured with $(d + 1)$ colours.

For a compact metrisable space X ,


$$\dim_{\text{nuc}} C(X, \mathbb{C}) = \dim X = \dim_{\text{nuc}} C(X, M_p(\mathbb{C}))$$

If A is simple and $\dim_{\text{nuc}} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

Nuclear dimension: properties

Definition

$\dim_{\text{nuc}} A \leq d$ if there exist approximately commuting diagrams



such that ψ_n is a completely positive contraction and (F_n, ϕ_n) can be coloured with $(d + 1)$ colours.

For a compact metrisable space X ,


$$\dim_{\text{nuc}} C(X, \mathbb{C}) = \dim X = \dim_{\text{nuc}} C(X, M_p(\mathbb{C}))$$

If A is simple and $\dim_{\text{nuc}} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

Nuclear dimension: properties

Definition

$\dim_{\text{nuc}} A \leq d$ if there exist approximately commuting diagrams



such that ψ_n is a completely positive contraction and (F_n, ϕ_n) can be coloured with $(d + 1)$ colours.

For a compact metrisable space X ,


$$\dim_{\text{nuc}} C(X, \mathbb{C}) = \dim X = \dim_{\text{nuc}} C(X, M_p(\mathbb{C}))$$

If A is simple and $\dim_{\text{nuc}} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

Nuclear dimension: properties

Definition

$\dim_{nuc} A \leq d$ if there exist approximately commuting diagrams



such that ψ_n is a completely positive contraction and (F_n, ϕ_n) can be coloured with $(d + 1)$ colours.

For a compact metrisable space X ,


$$\dim_{nuc} C(X, \mathbb{C}) = \dim X = \dim_{nuc} C(X, M_p(\mathbb{C}))$$

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

Nuclear dimension: properties

Definition

$\dim_{\text{nuc}} A \leq d$ if there exist approximately commuting diagrams



such that ψ_n is a completely positive contraction and (F_n, ϕ_n) can be coloured with $(d + 1)$ colours.

For a compact metrisable space X ,

$$\dim_{\text{nuc}} C(X, \mathbb{C}) = \dim X = \dim_{\text{nuc}} C(X, M_p(\mathbb{C}))$$

If A is simple and $\dim_{\text{nuc}} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

The Jiang-Su algebra

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

What is \mathcal{Z} ?

It is a lot like M_{p^∞} except no nontrivial projections; in fact, it is more like an interpolation between M_{p^∞} and M_{q^∞} where p, q are coprime.

$$\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z};$$

$$\mathcal{M}_{p^\infty} \cong \mathcal{Z} \otimes \mathcal{Z}.$$

The Jiang-Su algebra

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

What is \mathcal{Z} ?

It is a lot like M_{p^∞} except no nontrivial projections; in fact, it is more like an interpolation between M_{p^∞} and M_{q^∞} where p, q are coprime.

$$\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z};$$

$$\mathcal{M}_{p^\infty} \cong \mathcal{Z} \otimes \mathcal{Z}.$$

The Jiang-Su algebra

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

What is \mathcal{Z} ?

It is a lot like M_{p^∞} except no nontrivial projections; in fact, it is more like an interpolation between M_{p^∞} and M_{q^∞} where p, q are coprime.

$$\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z};$$

$$\mathcal{M}_{p^\infty} \cong \mathcal{Z} \otimes \mathcal{Z}.$$

The Jiang-Su algebra

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

What is \mathcal{Z} ?

It is a lot like M_{p^∞} except no nontrivial projections; in fact, is more like an interpolation between M_{p^∞} and M_{q^∞} where p, q are coprime.

$$\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z};$$

$$\mathcal{M}_{p^\infty} \cong \mathcal{Z} \otimes \mathcal{Z}.$$

The Jiang-Su algebra

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

What is \mathcal{Z} ?

It is a lot like M_{p^∞} except no nontrivial projections; in fact, it is more like an interpolation between M_{p^∞} and M_{q^∞} where p, q are coprime.

$$\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z};$$

$$\mathcal{M}_{p^\infty} \cong \mathcal{Z} \otimes \mathcal{Z}.$$

The Jiang-Su algebra

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

What is \mathcal{Z} ?

It is a lot like M_{p^∞} except no nontrivial projections; in fact, it is more like an interpolation between M_{p^∞} and M_{q^∞} where p, q are coprime.

$$\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z};$$

$$\mathcal{M}_{p^\infty} \cong \mathcal{Z} \otimes \mathcal{Z}.$$

The Jiang-Su algebra

If A is simple and $\dim_{nuc} A < \infty$ then either $A = M_k(\mathbb{C})$ (some k) or $A \cong A \otimes \mathcal{Z}$.

What is \mathcal{Z} ?

It is a lot like M_{p^∞} except no nontrivial projections; in fact, it is more like an interpolation between M_{p^∞} and M_{q^∞} where p, q are coprime.

$$\mathcal{Z} \cong \mathcal{Z} \otimes \mathcal{Z};$$

$$\mathcal{M}_{p^\infty} \cong \mathcal{Z} \otimes \mathcal{Z}.$$

Dimension reduction

What is $\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty}$?

There is tension between the constant dimension of $C(X, M_{p^n}) = C(X, \mathbb{C}) \otimes M_{p^n}$ and the extra space afforded by the regularity of M_{p^∞} .

Theorem (T-Winter '14)

$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{p^\infty}) \leq 2$ for any compact space X .
(Also, $\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes \mathcal{Z}) \leq 2$.)

This is a bit surprising, since

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes C(Y, \mathbb{C})) = \dim(X \times Y) \geq \dim X = \dim_{\text{nuc}} C(X, \mathbb{C}),$$

so dimension reduction, where

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{2^\infty}) < \dim_{\text{nuc}} C(X, \mathbb{C})$$

is a purely noncommutative phenomenon.



Dimension reduction

What is $\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty}$?

There is tension between the constant dimension of $C(X, M_{p^n}) = C(X, \mathbb{C}) \otimes M_{p^n}$ and the extra space afforded by the regularity of M_{p^∞} .

Theorem (T-Winter '14)

$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{p^\infty}) \leq 2$ for any compact space X .
(Also, $\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes \mathcal{Z}) \leq 2$.)

This is a bit surprising, since

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes C(Y, \mathbb{C})) = \dim(X \times Y) \geq \dim X = \dim_{\text{nuc}} C(X, \mathbb{C}),$$

so dimension reduction, where

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{2^\infty}) < \dim_{\text{nuc}} C(X, \mathbb{C})$$

is a purely noncommutative phenomenon.



Dimension reduction

What is $\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty}$?

There is tension between the constant dimension of $C(X, M_{p^n}) = C(X, \mathbb{C}) \otimes M_{p^n}$ and the extra space afforded by the regularity of M_{p^∞} .

Theorem (T-Winter '14)

$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{p^\infty}) \leq 2$ for any compact space X .
(Also, $\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes \mathcal{Z}) \leq 2$.)

This is a bit surprising, since

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes C(Y, \mathbb{C})) = \dim(X \times Y) \geq \dim X = \dim_{\text{nuc}} C(X, \mathbb{C}),$$

so dimension reduction, where

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{2^\infty}) < \dim_{\text{nuc}} C(X, \mathbb{C})$$

is a purely noncommutative phenomenon.



Dimension reduction

What is $\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty}$?

There is tension between the constant dimension of $C(X, M_{p^n}) = C(X, \mathbb{C}) \otimes M_{p^n}$ and the extra space afforded by the regularity of M_{p^∞} .

Theorem (T-Winter '14)

$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{p^\infty}) \leq 2$ for any compact space X .
(Also, $\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes \mathcal{Z}) \leq 2$.)

This is a bit surprising, since

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes C(Y, \mathbb{C})) = \dim(X \times Y) \geq \dim X = \dim_{\text{nuc}} C(X, \mathbb{C}),$$

so dimension reduction, where

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{2^\infty}) < \dim_{\text{nuc}} C(X, \mathbb{C})$$

is a purely noncommutative phenomenon.



Dimension reduction

What is $\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty}$?

There is tension between the constant dimension of $C(X, M_{p^n}) = C(X, \mathbb{C}) \otimes M_{p^n}$ and the extra space afforded by the regularity of M_{p^∞} .

Theorem (T-Winter '14)

$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{p^\infty}) \leq 2$ for any compact space X .
(Also, $\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes \mathcal{Z}) \leq 2$.)

This is a bit surprising, since

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes C(Y, \mathbb{C})) = \dim(X \times Y) \geq \dim X = \dim_{\text{nuc}} C(X, \mathbb{C}),$$

so dimension reduction, where

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{2^\infty}) < \dim_{\text{nuc}} C(X, \mathbb{C})$$

is a purely noncommutative phenomenon.



Dimension reduction

What is $\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty}$?

There is tension between the constant dimension of $C(X, M_{p^n}) = C(X, \mathbb{C}) \otimes M_{p^n}$ and the extra space afforded by the regularity of M_{p^∞} .

Theorem (T-Winter '14)

$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{p^\infty}) \leq 2$ for any compact space X .
(Also, $\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes \mathcal{Z}) \leq 2$.)

This is a bit surprising, since

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes C(Y, \mathbb{C})) = \dim(X \times Y) \geq \dim X = \dim_{\text{nuc}} C(X, \mathbb{C}),$$

so dimension reduction, where

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{2^\infty}) < \dim_{\text{nuc}} C(X, \mathbb{C})$$

is a purely noncommutative phenomenon.



Dimension reduction

What is $\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty}$?

There is tension between the constant dimension of $C(X, M_{p^n}) = C(X, \mathbb{C}) \otimes M_{p^n}$ and the extra space afforded by the regularity of M_{p^∞} .

Theorem (T-Winter '14)

$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{p^\infty}) \leq 2$ for any compact space X .
(Also, $\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes \mathcal{Z}) \leq 2$.)

This is a bit surprising, since

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes C(Y, \mathbb{C})) = \dim(X \times Y) \geq \dim X = \dim_{\text{nuc}} C(X, \mathbb{C}),$$

so dimension reduction, where

$$\dim_{\text{nuc}} (C(X, \mathbb{C}) \otimes M_{2^\infty}) < \dim_{\text{nuc}} C(X, \mathbb{C})$$

is a purely noncommutative phenomenon.

Dimension reduction

Theorem (T-Winter '12)

$\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty} \leq 2$ for any compact space X .

More generally, inspired by what is known (and conjectured) for classifiable C^* -algebras:

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

(Equivalently, $A \otimes \mathcal{Z}$ has finite nuclear dimension.)

Dimension reduction

Theorem (T-Winter '12)

$\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty} \leq 2$ for any compact space X .

More generally, inspired by what is known (and conjectured) for classifiable C^* -algebras:

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

(Equivalently, $A \otimes \mathcal{Z}$ has finite nuclear dimension.)

Dimension reduction

Theorem (T-Winter '12)

$\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty} \leq 2$ for any compact space X .

More generally, inspired by what is known (and conjectured) for classifiable C^* -algebras:

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

(Equivalently, $A \otimes \mathcal{Z}$ has finite nuclear dimension.)

Theorem (T-Winter '12)

$\dim_{\text{nuc}} C(X, \mathbb{C}) \otimes M_{p^\infty} \leq 2$ for any compact space X .

More generally, inspired by what is known (and conjectured) for classifiable C^* -algebras:

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

(Equivalently, $A \otimes \mathcal{Z}$ has finite nuclear dimension.)

Dimension reduction

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

Theorem (Elliott-Niu-Santiago-T-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 2$ if A is subhomogeneous, i.e., a subalgebra of $C(X, M_k)$.

Theorem (Matui-Sato '13, Sato-White-Winter '14, Bosa-Brown-Sato-T-White-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 1$ if A is simple, unital, amenable, and not too many extreme traces. (Specifically, if $\partial_e T(A)$ is compact.)

Dimension reduction

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

Theorem (Elliott-Niu-Santiago-T-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 2$ if A is subhomogeneous, i.e., a subalgebra of $C(X, M_k)$.

Theorem (Matui-Sato '13, Sato-White-Winter '14, Bosa-Brown-Sato-T-White-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 1$ if A is simple, unital, amenable, and not too many extreme traces. (Specifically, if $\partial_e T(A)$ is compact.)

Dimension reduction

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

Theorem (Elliott-Niu-Santiago-T-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 2$ if A is subhomogeneous, i.e., a subalgebra of $C(X, M_k)$.

Theorem (Matui-Sato '13, Sato-White-Winter '14, Bosa-Brown-Sato-T-White-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 1$ if A is simple, unital, amenable, and not too many extreme traces. (Specifically, if $\partial_e T(A)$ is compact.)

Dimension reduction

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

Theorem (Elliott-Niu-Santiago-T-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 2$ if A is subhomogeneous, i.e., a subalgebra of $C(X, M_k)$.

Theorem (Matui-Sato '13, Sato-White-Winter '14, Bosa-Brown-Sato-T-White-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 1$ if A is simple, unital, amenable, and not too many extreme traces. (Specifically, if $\partial_e T(A)$ is compact.)

Dimension reduction

Conjecture

If A is simple and nuclear then $A \otimes M_{p^\infty}$ has finite nuclear dimension (for any p).

Theorem (Elliott-Niu-Santiago-T-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 2$ if A is subhomogeneous, i.e., a subalgebra of $C(X, M_k)$.

Theorem (Matui-Sato '13, Sato-White-Winter '14, Bosa-Brown-Sato-T-White-Winter '14)

$\dim_{nuc}(A \otimes M_{p^\infty}) \leq 1$ if A is simple, unital, amenable, and not too many extreme traces. (Specifically, if $\partial_e T(A)$ is compact.)