

# Convex structure of unital quantum channels, factorizability and traces on the universal free product of matrix algebras

Magdalena Musat  
University of Copenhagen

Canadian Operator Algebras Symposium  
University of Ottawa  
May 31, 2022

**Central theme:** A certain class of completely positive maps (**factorizable maps**), introduced by C. Anantharaman-Delaroche '05.

Their study **led to** investigating the **convex structure** of the set of **unital quantum channels**, interesting applications in the **analysis of QIT** (e.g., settling in the negative the Asymptotic Quantum Birkhoff Conjecture) and revealed **infinite dim phenomena** therein, connections to/reformulations of the **Connes Embedding Problem**, and recently, through a **new** view-point, some interesting problems in operator algebras.

For  $n \geq 2$ , consider following sets of maps  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ :

$$\mathcal{FM}(n) \subseteq \text{UCPT}(n) \subseteq \begin{matrix} \text{CPT}(n) \\ \text{UCP}(n) \end{matrix}$$

all **compact** and **convex** subsets of  $\mathbb{C}^{n^4}$ .

**Central theme:** A certain class of completely positive maps (**factorizable maps**), introduced by C. Anantharaman-Delaroche '05.

Their study **led to** investigating the **convex structure** of the set of **unital quantum channels**, interesting applications in the **analysis of QIT** (e.g., settling in the negative the Asymptotic Quantum Birkhoff Conjecture) and revealed **infinite dim phenomena** therein, connections to/reformulations of the **Connes Embedding Problem**, and recently, through a **new** view-point, some interesting problems in operator algebras.

For  $n \geq 2$ , consider following sets of maps  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ :

$$\mathcal{FM}(n) \subseteq \text{UCPT}(n) \subseteq \begin{matrix} \text{CPT}(n) \\ \text{UCP}(n) \end{matrix}$$

all **compact** and **convex** subsets of  $\mathbb{C}^{n^4}$ .

► (Choi '73): Let  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  linear. Then

$$T \in \text{CPT}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j A_j^* = I_n.$$

$$T \in \text{UCP}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j^* A_j = I_n.$$

When  $\{A_1, \dots, A_d\}$  **lin independent**,  $d$  is called the **Choi-rank** of  $T$ .

**Thm** (Choi '75):  $T \in \partial_e(\text{CPT}(n)) \iff \{A_i A_j^*\}_{i,j=1}^d$  lin independent.  
 Respectively,  $T \in \partial_e(\text{UCP}(n)) \iff \{A_i^* A_j\}_{i,j=1}^d$  lin independent.

**Thm** (Landau-Streater '93):  $T \in \partial_e(\text{UCPT}(n)) \iff \{A_i^* A_j \oplus A_j A_i^*\}_{i,j}$  lin independent.

Hence  $\partial_e(\text{UCPT}(n)) \supseteq (\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n)$ .

**Question:** Are these sets equal?

► (Choi '73): Let  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  linear. Then

$$T \in \text{CPT}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j A_j^* = I_n.$$

$$T \in \text{UCP}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j^* A_j = I_n.$$

When  $\{A_1, \dots, A_d\}$  **lin independent**,  $d$  is called the **Choi-rank** of  $T$ .

**Thm** (Choi '75):  $T \in \partial_e(\text{CPT}(n)) \iff \{A_i A_j^*\}_{i,j=1}^d$  lin independent.  
 Respectively,  $T \in \partial_e(\text{UCP}(n)) \iff \{A_i^* A_j\}_{i,j=1}^d$  lin independent.

**Thm** (Landau-Streater '93):  $T \in \partial_e(\text{UCPT}(n)) \iff \{A_i^* A_j \oplus A_j A_i^*\}_{i,j}$  lin independent.

Hence  $\partial_e(\text{UCPT}(n)) \supseteq (\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n)$ .

**Question:** Are these sets equal?

► (Choi '73): Let  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  linear. Then

$$T \in \text{CPT}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j A_j^* = I_n.$$

$$T \in \text{UCP}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j^* A_j = I_n.$$

When  $\{A_1, \dots, A_d\}$  **lin independent**,  $d$  is called the **Choi-rank** of  $T$ .

**Thm** (Choi '75):  $T \in \partial_e(\text{CPT}(n)) \iff \{A_i A_j^*\}_{i,j=1}^d$  lin independent.

Respectively,  $T \in \partial_e(\text{UCP}(n)) \iff \{A_i^* A_j\}_{i,j=1}^d$  lin independent.

**Thm** (Landau-Streater '93):  $T \in \partial_e(\text{UCPT}(n)) \iff \{A_i^* A_j \oplus A_j A_i^*\}_{i,j}$  lin independent.

Hence  $\partial_e(\text{UCPT}(n)) \supseteq (\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n)$ .

**Question:** Are these sets equal?

► (Choi '73): Let  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  linear. Then

$$T \in \text{CPT}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j A_j^* = I_n.$$

$$T \in \text{UCP}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j^* A_j = I_n.$$

When  $\{A_1, \dots, A_d\}$  **lin independent**,  $d$  is called the **Choi-rank** of  $T$ .

**Thm** (Choi '75):  $T \in \partial_e(\text{CPT}(n)) \iff \{A_i A_j^*\}_{i,j=1}^d$  lin independent.

Respectively,  $T \in \partial_e(\text{UCP}(n)) \iff \{A_i^* A_j\}_{i,j=1}^d$  lin independent.

**Thm** (Landau-Streater '93):  $T \in \partial_e(\text{UCPT}(n)) \iff \{A_i^* A_j \oplus A_j A_i^*\}_{i,j}$  lin independent.

Hence  $\partial_e(\text{UCPT}(n)) \supseteq (\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n)$ .

**Question:** Are these sets equal?

► (Choi '73): Let  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  linear. Then

$$T \in \text{CPT}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j A_j^* = I_n.$$

$$T \in \text{UCP}(n) \iff \exists A_1, \dots, A_d \in M_n(\mathbb{C}) : Tx = \sum_{j=1}^d A_j^* x A_j, \quad \sum_{j=1}^d A_j^* A_j = I_n.$$

When  $\{A_1, \dots, A_d\}$  **lin independent**,  $d$  is called the **Choi-rank** of  $T$ .

**Thm** (Choi '75):  $T \in \partial_e(\text{CPT}(n)) \iff \{A_i A_j^*\}_{i,j=1}^d$  lin independent.

Respectively,  $T \in \partial_e(\text{UCP}(n)) \iff \{A_i^* A_j\}_{i,j=1}^d$  lin independent.


**Thm** (Landau-Streater '93):  $T \in \partial_e(\text{UCPT}(n)) \iff \{A_i^* A_j \oplus A_j A_i^*\}_{i,j}$  lin independent.

Hence  $\partial_e(\text{UCPT}(n)) \supseteq (\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n)$ .

**Question:** Are these sets equal?



**Definition** (Anantharaman-Delaroche '05): A unital quantum channel  $T: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  is called **factorizable** if  $\exists$  vN alg  $(N, \psi)$  with n.f. tracial state and unital  $*$ -homs  $\alpha, \beta: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C}) \otimes N$ :  $T = \beta^* \circ \alpha$ .




►  $\alpha, \beta$  are injective (thus embeddings) and trace-preserving. Since unital embeddings of  $M_n(\mathbb{C})$  into a vN alg are **unitarily equiv**, can take

$$\beta(x) = x \otimes 1_N, \quad \alpha(x) = u^* \beta(x) u, \quad x \in M_n(\mathbb{C}),$$

for some  $u \in M_n(\mathbb{C}) \otimes N$  unitary;  $N$  can be taken  $\text{II}_1$ -vN alg (even factor).

**Definition** (Anantharaman-Delaroche '05): A unital quantum channel  $T: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  is called **factorizable** if  $\exists$  vN alg  $(N, \psi)$  with n.f. tracial state and unital  $*$ -homs  $\alpha, \beta: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C}) \otimes N$ :  $T = \beta^* \circ \alpha$ .



►  $\alpha, \beta$  are injective (thus embeddings) and trace-preserving. Since unital embeddings of  $M_n(\mathbb{C})$  into a vN alg are **unitarily equiv**, can take

$$\beta(x) = x \otimes 1_N, \quad \alpha(x) = u^* \beta(x) u, \quad x \in M_n(\mathbb{C}),$$

for some  $u \in M_n(\mathbb{C}) \otimes N$  **unitary**;  $N$  can be taken  $\text{II}_1$ -vN alg (even factor).

**Theorem** (Haagerup-M '11):  $T: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  is a **factorizable** quantum channel **iff**  $\exists (N, \tau_N)$  finite vN algebra (called **ancilla**) and a unitary  $u \in M_n(\mathbb{C}) \otimes N$ :  $Tx = (\text{id}_{M_n(\mathbb{C})} \otimes \tau_N)(u^*(x \otimes 1_N)u)$ ,  $x \in M_n(\mathbb{C})$ .

**Def/Thm** (Haagerup-M '11):  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  is a **factorizable** quantum channel **iff**  $\exists (N, \tau_N)$  finite vN algebra (called **ancilla**) and a unitary  $u \in M_n(\mathbb{C}) \otimes N$ :  $Tx = (\text{id}_{M_n(\mathbb{C})} \otimes \tau_N)(u^*(x \otimes 1_N)u)$ ,  $x \in M_n(\mathbb{C})$ .

► (R. Werner): **Factorizable channels** are obtained by coupling the input system to a **maximally mixed** ancillary one, executing a **unitary rotation** on the combined system, and **tracing out** the ancilla.

► Automorphisms of  $M_n(\mathbb{C})$  are **factorizable**.

Let  $\mathcal{FM}(n)$  denote all factorizable quantum channels on  $M_n(\mathbb{C})$ ,  $n \geq 2$ . Then  $\mathcal{FM}(n)$  is **convex** and **closed**.

**Proposition** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ , with **canonical form**

$$Tx = \sum_{i=1}^d A_i^* x A_i, \quad x \in M_n(\mathbb{C}).$$

If  $d := \text{Choi-rank}(T) \geq 2$  and  $\{A_i^* A_j\}_{1 \leq i,j}^d$  **lin indep**, then  $T \notin \mathcal{FM}(n)$ .

**Def/Thm** (Haagerup-M '11):  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  is a **factorizable** quantum channel **iff**  $\exists (N, \tau_N)$  finite vN algebra (called **ancilla**) and a unitary  $u \in M_n(\mathbb{C}) \otimes N$ :  $Tx = (\text{id}_{M_n(\mathbb{C})} \otimes \tau_N)(u^*(x \otimes 1_N)u)$ ,  $x \in M_n(\mathbb{C})$ .

► (R. Werner): **Factorizable channels** are obtained by coupling the input system to a **maximally mixed** ancillary one, executing a **unitary rotation** on the combined system, and **tracing out** the ancilla.

► Automorphisms of  $M_n(\mathbb{C})$  are **factorizable**.

Let  $\mathcal{FM}(n)$  denote all factorizable quantum channels on  $M_n(\mathbb{C})$ ,  $n \geq 2$ . Then  $\mathcal{FM}(n)$  is **convex** and **closed**.

**Proposition** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ , with **canonical form**

$$Tx = \sum_{i=1}^d A_i^* x A_i, \quad x \in M_n(\mathbb{C}).$$

If  $d := \text{Choi-rank}(T) \geq 2$  and  $\{A_i^* A_j\}_{1 \leq i,j}^d$  **lin indep**, then  $T \notin \mathcal{FM}(n)$ .

**Def/Thm** (Haagerup-M '11):  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  is a **factorizable** quantum channel **iff**  $\exists (N, \tau_N)$  finite vN algebra (called **ancilla**) and a unitary  $u \in M_n(\mathbb{C}) \otimes N$ :  $Tx = (\text{id}_{M_n(\mathbb{C})} \otimes \tau_N)(u^*(x \otimes 1_N)u)$ ,  $x \in M_n(\mathbb{C})$ .

► (R. Werner): **Factorizable channels** are obtained by coupling the input system to a **maximally mixed** ancillary one, executing a **unitary rotation** on the combined system, and **tracing out** the ancilla.

► Automorphisms of  $M_n(\mathbb{C})$  are **factorizable**.

Let  $\mathcal{FM}(n)$  denote all factorizable quantum channels on  $M_n(\mathbb{C})$ ,  $n \geq 2$ . Then  $\mathcal{FM}(n)$  is **convex** and **closed**.

**Proposition** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ , with **canonical form**

$$Tx = \sum_{i=1}^d A_i^* x A_i, \quad x \in M_n(\mathbb{C}).$$

If  $d := \text{Choi-rank}(T) \geq 2$  and  $\{A_i^* A_j\}_{1 \leq i, j}^d$  **lin indep**, then  $T \notin \mathcal{FM}(n)$ .

**Def/Thm** (Haagerup-M '11):  $T : M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$  is a **factorizable** quantum channel **iff**  $\exists (N, \tau_N)$  finite vN algebra (called **ancilla**) and a unitary  $u \in M_n(\mathbb{C}) \otimes N$ :  $Tx = (\text{id}_{M_n(\mathbb{C})} \otimes \tau_N)(u^*(x \otimes 1_N)u)$ ,  $x \in M_n(\mathbb{C})$ .

► (R. Werner): **Factorizable channels** are obtained by coupling the input system to a **maximally mixed** ancillary one, executing a **unitary rotation** on the combined system, and **tracing out** the ancilla.

► Automorphisms of  $M_n(\mathbb{C})$  are **factorizable**.

Let  $\mathcal{FM}(n)$  denote all factorizable quantum channels on  $M_n(\mathbb{C})$ ,  $n \geq 2$ . Then  $\mathcal{FM}(n)$  is **convex** and **closed**.

**Proposition** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ , with **canonical form**

$$Tx = \sum_{i=1}^d A_i^* x A_i, \quad x \in M_n(\mathbb{C}).$$

If  $d := \text{Choi-rank}(T) \geq 2$  and  $\{A_i^* A_j\}_{1 \leq i, j}^d$  **lin indep**, then  $T \notin \mathcal{FM}(n)$ .

► (Kümmerer '83):  $\text{UCPT}(2) = \text{conv}(\text{Aut}(M_2(\mathbb{C})))$ . However,

$$\text{UCPT}(n) \not\supseteq \text{conv}(\text{Aut}(M_n(\mathbb{C}))), \quad n \geq 3,$$

cf. (Kümmerer '86, Landau-Streater '93, Kümmerer-Maasen '87).

► **Asymptotic Quantum Birkhoff Conj** (Smolin-Verstraete-Winter '05):

Any  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ , satisfies

$$\lim_{k \rightarrow \infty} d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \text{conv}(\text{Aut}(\bigotimes_{i=1}^k M_n(\mathbb{C}))) \right) = 0.$$

**Theorem** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ . For  $k \geq 1$ ,

$$d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \mathcal{F} \left( \bigotimes_{i=1}^k M_n(\mathbb{C}) \right) \right) \geq d_{\text{cb}}(T, \mathcal{FM}(n)).$$

► If  $T \notin \mathcal{FM}(n)$ , then  $d_{\text{cb}}(T, \mathcal{FM}(n)) > 0$ .

As  $\text{conv}(\text{Aut}(M_n(\mathbb{C}))) \subset \mathcal{FM}(n)$ , any non-factoriz channel fails **AQBC**.

► (Kümmerer '83):  $\text{UCPT}(2) = \text{conv}(\text{Aut}(M_2(\mathbb{C})))$ . However,

$$\text{UCPT}(n) \not\supseteq \text{conv}(\text{Aut}(M_n(\mathbb{C}))), \quad n \geq 3,$$

cf. (Kümmerer '86, Landau-Streater '93, Kümmerer-Maasen '87).

► **Asymptotic Quantum Birkhoff Conj** (Smolin-Verstraete-Winter '05):

Any  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ , satisfies

$$\lim_{k \rightarrow \infty} d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \text{conv}(\text{Aut}(\bigotimes_{i=1}^k M_n(\mathbb{C}))) \right) = 0.$$

**Theorem** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ . For  $k \geq 1$ ,

$$d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \mathcal{F} \left( \bigotimes_{i=1}^k M_n(\mathbb{C}) \right) \right) \geq d_{\text{cb}}(T, \mathcal{FM}(n)).$$

► If  $T \notin \mathcal{FM}(n)$ , then  $d_{\text{cb}}(T, \mathcal{FM}(n)) > 0$ .

As  $\text{conv}(\text{Aut}(M_n(\mathbb{C}))) \subset \mathcal{FM}(n)$ , any non-factoriz channel fails **AQBC**.



► (Kümmerer '83):  $\text{UCPT}(2) = \text{conv}(\text{Aut}(M_2(\mathbb{C})))$ . However,

$$\text{UCPT}(n) \not\supseteq \text{conv}(\text{Aut}(M_n(\mathbb{C}))), \quad n \geq 3,$$

cf. (Kümmerer '86, Landau-Streater '93, Kümmerer-Maasen '87).

► **Asymptotic Quantum Birkhoff Conj** (Smolin-Verstraete-Winter '05):  
Any  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ , satisfies

$$\lim_{k \rightarrow \infty} d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \text{conv}(\text{Aut}(\bigotimes_{i=1}^k M_n(\mathbb{C}))) \right) = 0.$$

**Theorem** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ . For  $k \geq 1$ ,

$$d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \mathcal{F} \left( \bigotimes_{i=1}^k M_n(\mathbb{C}) \right) \right) \geq d_{\text{cb}}(T, \mathcal{FM}(n)).$$

► If  $T \notin \mathcal{FM}(n)$ , then  $d_{\text{cb}}(T, \mathcal{FM}(n)) > 0$ .

As  $\text{conv}(\text{Aut}(M_n(\mathbb{C}))) \subset \mathcal{FM}(n)$ , any non-factoriz channel fails **AQBC**.

► (Kümmerer '83):  $\text{UCPT}(2) = \text{conv}(\text{Aut}(M_2(\mathbb{C})))$ . However,

$$\text{UCPT}(n) \not\supseteq \text{conv}(\text{Aut}(M_n(\mathbb{C}))), \quad n \geq 3,$$

cf. (Kümmerer '86, Landau-Streater '93, Kümmerer-Maasen '87).

► **Asymptotic Quantum Birkhoff Conj** (Smolin-Verstraete-Winter '05):

Any  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ , satisfies

$$\lim_{k \rightarrow \infty} d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \text{conv}(\text{Aut}(\bigotimes_{i=1}^k M_n(\mathbb{C}))) \right) = 0.$$

**Theorem** (Haagerup-M '11): Let  $T \in \text{UCPT}(n)$ ,  $n \geq 3$ . For  $k \geq 1$ ,

$$d_{\text{cb}} \left( \bigotimes_{i=1}^k T, \mathcal{F} \left( \bigotimes_{i=1}^k M_n(\mathbb{C}) \right) \right) \geq d_{\text{cb}}(T, \mathcal{FM}(n)).$$

► If  $T \notin \mathcal{FM}(n)$ , then  $d_{\text{cb}}(T, \mathcal{FM}(n)) > 0$ .

As  $\text{conv}(\text{Aut}(M_n(\mathbb{C}))) \subset \mathcal{FM}(n)$ , any non-factoriz channel fails **AQBC**.

► For  $T \in \text{UCPT}(n)$ ,  $\text{Choi-rank}(T) = 1$  **iff**  $T \in \text{Aut}(M_n(\mathbb{C}))$ .

Set  $\partial_e^*(\text{CPT}(n)) = \partial_e(\text{CPT}(n)) \setminus \text{Aut}(M_n(\mathbb{C}))$ , and similarly  $\partial_e^*(\text{UCP}(n))$ .

**Cor:**  $T \in (\partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))) \cap \text{UCPT}(n) \Rightarrow T \notin \mathcal{FM}(n)$ .

**Remark:** Not easy to characterize non-factorizability in terms of the convex structure of  $\text{UCPT}(n)$ :

- $\partial_e(\text{UCPT}(n)) \setminus ((\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n))$  can contain both factorizable and non-factorizable maps.
- $T \in \partial_e^*(\text{UCPT}(n))$ ,  $\text{Choi-rank} > n \Rightarrow T \notin \partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))$ .
- (Ohno '09):  $\exists T \in \partial_e^*(\text{UCPT}(3))$ , Choi-rank 4; (H-M-R):  $T \notin \mathcal{FM}(3)$ .
- (H-M-R '21): Explicit family  $T_t \in \partial_e^*(\text{UCPT}(3)) \cap \mathcal{FM}(3)$ , Choi-rank 4.

► For  $T \in \text{UCPT}(n)$ ,  $\text{Choi-rank}(T) = 1$  **iff**  $T \in \text{Aut}(M_n(\mathbb{C}))$ .

Set  $\partial_e^*(\text{CPT}(n)) = \partial_e(\text{CPT}(n)) \setminus \text{Aut}(M_n(\mathbb{C}))$ , and similarly  $\partial_e^*(\text{UCP}(n))$ .

**Cor:**  $T \in (\partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))) \cap \text{UCPT}(n) \Rightarrow T \notin \mathcal{FM}(n)$ .

**Remark:** Not easy to characterize non-factorizability in terms of the convex structure of  $\text{UCPT}(n)$ :

- $\partial_e(\text{UCPT}(n)) \setminus ((\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n))$  can contain both factorizable and non-factorizable maps.
- $T \in \partial_e^*(\text{UCPT}(n))$ ,  $\text{Choi-rank} > n \Rightarrow T \notin \partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))$ .
- (Ohno '09):  $\exists T \in \partial_e^*(\text{UCPT}(3))$ , Choi-rank 4; (H-M-R):  $T \notin \mathcal{FM}(3)$ .
- (H-M-R '21): Explicit family  $T_t \in \partial_e^*(\text{UCPT}(3)) \cap \mathcal{FM}(3)$ , Choi-rank 4.

► For  $T \in \text{UCPT}(n)$ ,  $\text{Choi-rank}(T) = 1$  **iff**  $T \in \text{Aut}(M_n(\mathbb{C}))$ .

Set  $\partial_e^*(\text{CPT}(n)) = \partial_e(\text{CPT}(n)) \setminus \text{Aut}(M_n(\mathbb{C}))$ , and similarly  $\partial_e^*(\text{UCP}(n))$ .

**Cor:**  $T \in (\partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))) \cap \text{UCPT}(n) \Rightarrow T \notin \mathcal{FM}(n)$ .

**Remark:** Not easy to characterize non-factorizability in terms of the convex structure of  $\text{UCPT}(n)$ :

- $\partial_e(\text{UCPT}(n)) \setminus ((\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n))$  can contain both factorizable and non-factorizable maps.
- $T \in \partial_e^*(\text{UCPT}(n))$ ,  $\text{Choi-rank} > n \Rightarrow T \notin \partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))$ .
- (Ohno '09):  $\exists T \in \partial_e^*(\text{UCPT}(3))$ , Choi-rank 4; (H-M-R):  $T \notin \mathcal{FM}(3)$ .
- (H-M-R '21): Explicit family  $T_t \in \partial_e^*(\text{UCPT}(3)) \cap \mathcal{FM}(3)$ , Choi-rank 4.

► For  $T \in \text{UCPT}(n)$ ,  $\text{Choi-rank}(T) = 1$  **iff**  $T \in \text{Aut}(M_n(\mathbb{C}))$ .

Set  $\partial_e^*(\text{CPT}(n)) = \partial_e(\text{CPT}(n)) \setminus \text{Aut}(M_n(\mathbb{C}))$ , and similarly  $\partial_e^*(\text{UCP}(n))$ .

**Cor:**  $T \in (\partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))) \cap \text{UCPT}(n) \Rightarrow T \notin \mathcal{FM}(n)$ .

**Remark:** Not easy to characterize non-factorizability in terms of the convex structure of  $\text{UCPT}(n)$ :

- $\partial_e(\text{UCPT}(n)) \setminus ((\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n))$  can contain both factorizable and non-factorizable maps.
- $T \in \partial_e^*(\text{UCPT}(n))$ ,  $\text{Choi-rank} > n \Rightarrow T \notin \partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))$ .
- (Ohno '09):  $\exists T \in \partial_e^*(\text{UCPT}(3))$ , Choi-rank 4; (H-M-R):  $T \notin \mathcal{FM}(3)$ .
- (H-M-R '21): Explicit family  $T_t \in \partial_e^*(\text{UCPT}(3)) \cap \mathcal{FM}(3)$ , Choi-rank 4.

► For  $T \in \text{UCPT}(n)$ ,  $\text{Choi-rank}(T) = 1$  **iff**  $T \in \text{Aut}(M_n(\mathbb{C}))$ .

Set  $\partial_e^*(\text{CPT}(n)) = \partial_e(\text{CPT}(n)) \setminus \text{Aut}(M_n(\mathbb{C}))$ , and similarly  $\partial_e^*(\text{UCP}(n))$ .

**Cor:**  $T \in (\partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))) \cap \text{UCPT}(n) \Rightarrow T \notin \mathcal{FM}(n)$ .

**Remark:** Not easy to characterize non-factorizability in terms of the convex structure of  $\text{UCPT}(n)$ :

- $\partial_e(\text{UCPT}(n)) \setminus ((\partial_e(\text{UCP}(n)) \cup \partial_e(\text{CPT}(n))) \cap \text{UCPT}(n))$  can contain both factorizable and non-factorizable maps.
- $T \in \partial_e^*(\text{UCPT}(n))$ ,  $\text{Choi-rank} > n \Rightarrow T \notin \partial_e^*(\text{UCP}(n)) \cup \partial_e^*(\text{CPT}(n))$ .
- (Ohno '09):  $\exists T \in \partial_e^*(\text{UCPT}(3))$ , Choi-rank 4; (H-M-R):  $T \notin \mathcal{FM}(3)$ .
- (H-M-R '21): Explicit family  $T_t \in \partial_e^*(\text{UCPT}(3)) \cap \mathcal{FM}(3)$ , Choi-rank 4.

A class of  $\text{UCPT}(n)$  maps constructed in (Haagerup-M-Ruskai '21):

Given  $n \geq 3$ ,  $V_1, \dots, V_n \in \mathcal{U}(n-1)$  and  $t \in [-1, 1]$ ,  $t \neq -1/(n-1)$ , set

$$A_m = \frac{1}{\sqrt{n-1-t^2}} S^{-m+1} \begin{pmatrix} t & 0 \\ 0 & V_m \end{pmatrix} S^{m-1}, \quad 1 \leq m \leq n.$$

Here  $S$  is the canonical shift on  $\mathbb{C}^n$ .

► Can verify  $\sum_{m=1}^n A_m^* A_m = I_n = \sum_{m=1}^n A_m A_m^*$ . Thus, if

$$Tx = \sum_{m=1}^n A_m^* x A_m, \quad x \in M_n(\mathbb{C}),$$

then  $T \in \text{UCPT}(n)$ . Moreover, the Choi-rank of  $T$  is  $n$ .



A class of  $\text{UCPT}(n)$  maps constructed in (Haagerup-M-Ruskai '21):

Given  $n \geq 3$ ,  $V_1, \dots, V_n \in \mathcal{U}(n-1)$  and  $t \in [-1, 1]$ ,  $t \neq -1/(n-1)$ , set

$$A_m = \frac{1}{\sqrt{n-1-t^2}} S^{-m+1} \begin{pmatrix} t & 0 \\ 0 & V_m \end{pmatrix} S^{m-1}, \quad 1 \leq m \leq n.$$

Here  $S$  is the canonical shift on  $\mathbb{C}^n$ .

► Can verify  $\sum_{m=1}^n A_m^* A_m = I_n = \sum_{m=1}^n A_m A_m^*$ . Thus, if

$$Tx = \sum_{m=1}^n A_m^* x A_m, \quad x \in M_n(\mathbb{C}),$$

then  $T \in \text{UCPT}(n)$ . Moreover, the Choi-rank of  $T$  is  $n$ .

**Theorem A** (H-M-R '21): For  $n \geq 3$  and  $t \in (-1, 1)$ ,  $t \neq -1/(n-1)$ , there exists  $W = W^* \in \mathcal{U}(n-1)$  such that if  $V_1 = \dots = V_n = W$  and  $\{A_m\}_{m=1}^n$  are as above, then  $\{A_i^* A_j\}_{i,j=1}^n$  **linearly independent**. Hence

$$T \in \partial_e^*(UCP(n)) \cap \partial_e^*(CPT(n)),$$

thus  $T$  is non-factorizable.

**Proof:** Lots of linear algebra.

**Theorem B** (H-M-R '21): For  $n \geq 3$  and  $t \in (-1, 1)$ ,  $t \neq -1/(n-1)$ , the set of  $n$ -tuples  $(V_1, \dots, V_n) \in \mathcal{U}(n-1)^n$  such that  $\{A_i^* A_j\}_{i,j=1}^n$  is linearly indep has co-measure 0 w.r.t. Haar measure. Hence **almost all** quantum channels  $T$  arising in this way are **non-factorizable**.

**Proof:** Theorem A + Algebraic Geometry.

**Theorem A** (H-M-R '21): For  $n \geq 3$  and  $t \in (-1, 1)$ ,  $t \neq -1/(n-1)$ , there exists  $W = W^* \in \mathcal{U}(n-1)$  such that if  $V_1 = \dots = V_n = W$  and  $\{A_m\}_{m=1}^n$  are as above, then  $\{A_i^* A_j\}_{i,j=1}^n$  **linearly independent**. Hence

$$T \in \partial_e^*(UCP(n)) \cap \partial_e^*(CPT(n)),$$

thus  $T$  is non-factorizable.

**Proof:** Lots of linear algebra.

**Theorem B** (H-M-R '21): For  $n \geq 3$  and  $t \in (-1, 1)$ ,  $t \neq -1/(n-1)$ , the set of  $n$ -tuples  $(V_1, \dots, V_n) \in \mathcal{U}(n-1)^n$  such that  $\{A_i^* A_j\}_{i,j=1}^n$  is linearly indep has co-measure 0 w.r.t. Haar measure. Hence **almost all** quantum channels  $T$  arising in this way are **non-factorizable**.

**Proof:** Theorem A + Algebraic Geometry.

## Infinite dimensional phenomena in QIT:

**Question:** Do we need (inf dim) vN algs to describe factorizable channels?

► For a factorizable channel, *minimal* ancilla (and its size) **not** unique.  
E.g., consider the **completely depolarizing** channel  $S_n$ ,  $n \geq 2$

$$S_n(x) := \operatorname{tr}_n(x) 1_n = \int_{\mathcal{U}(n)} u^* x u \, d\mu(u), \quad x \in \mathbb{M}_n(\mathbb{C}).$$

It's factorizable, and **possible ancillas** are:  $\mathbb{C}^{n^2}$ ,  $M_n(\mathbb{C})$ , but also (a corner of) the reduced free product von Neumann alg  $(M_n(\mathbb{C}), \operatorname{tr}_n) * (M_n(\mathbb{C}), \operatorname{tr}_n)$ .

Let  $\mathcal{FM}_{\text{fin}}(n)$  = factoriz channels on  $M_n(\mathbb{C})$  admitting a **finite dim** ancilla.

**Theorem** (Rørdam-M '19):  $\mathcal{FM}_{\text{fin}}(n)$  is **not** closed, whenever  $n \geq 11$ .  
Moreover, for each such  $n$ , there exist factorizable quantum channels on  $M_n(\mathbb{C})$  which do require infinite dimensional (even type  $\text{II}_1$ ) ancilla.

Infinite dimensional phenomena in QIT:

**Question:** Do we need (inf dim) vN algs to describe factorizable channels?

► For a factorizable channel, *minimal* ancilla (and its size) **not** unique.  
E.g., consider the **completely depolarizing** channel  $S_n$ ,  $n \geq 2$

$$S_n(x) := \operatorname{tr}_n(x) 1_n = \int_{\mathcal{U}(n)} u^* x u \, d\mu(u), \quad x \in \mathbb{M}_n(\mathbb{C}).$$

It's factorizable, and **possible ancillas** are:  $\mathbb{C}^{n^2}$ ,  $M_n(\mathbb{C})$ , but also (a corner of) the reduced free product von Neumann alg  $(M_n(\mathbb{C}), \operatorname{tr}_n) * (M_n(\mathbb{C}), \operatorname{tr}_n)$ .

Let  $\mathcal{FM}_{\text{fin}}(n)$  = factoriz channels on  $M_n(\mathbb{C})$  admitting a **finite dim** ancilla.

**Theorem** (Rørdam-M '19):  $\mathcal{FM}_{\text{fin}}(n)$  is **not** closed, whenever  $n \geq 11$ .  
Moreover, for each such  $n$ , there exist factorizable quantum channels on  $M_n(\mathbb{C})$  which do require infinite dimensional (even type II<sub>1</sub>) ancilla.

**Theorem** (Rørdam-M '19):  $\mathcal{FM}_{\text{fin}}(n)$  is **not** closed, whenever  $n \geq 11$ . Moreover, for each such  $n$ , there exist factorizable quantum channels on  $M_n(\mathbb{C})$  which do require infinite dimensional (even type II<sub>1</sub>) ancilla.

**Proof:** Consider following sets of  $n \times n$  complex matrices,  $n \geq 2$ :

$$\begin{aligned} \mathcal{G}_{\text{fin}}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary} \right. \\ &\quad \left. \text{finite dimens } C^*\text{-alg } (\mathcal{A}, \tau) \right\}, \\ \cap \\ \mathcal{G}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary finite} \right. \\ &\quad \left. \text{vN alg } (M, \tau) \right\}. \end{aligned}$$

► (Rørdam-M):  $\mathcal{G}_{\text{fin}}(n)$  is convex  $\forall n \geq 2$ , but **not** closed whenever  $n \geq 11$ .

**Note:** If  $B \in M_n(\mathbb{C})$  *correlation matrix* (pos definite, diag entries equal 1), the Schur multiplier  $T_B \in UCPT(n)$ . (Ricard '08): If  $B$  has **real** entries, then  $T_B$  is factorizable.

► (Haagerup-M '11): A Schur multiplier  $T_B$  is **factorizable** iff  $B \in \mathcal{G}(n)$ . Furthermore,  $T_B \in \mathcal{FM}_{\text{fin}}(n)$  iff  $B \in \mathcal{G}_{\text{fin}}(n)$ .

The map  $B \mapsto T_B$  is an affine homeo, hence theorem follows.

**Theorem** (Rørdam-M '19):  $\mathcal{FM}_{\text{fin}}(n)$  is **not** closed, whenever  $n \geq 11$ .  
 Moreover, for each such  $n$ , there exist factorizable quantum channels on  $M_n(\mathbb{C})$  which do require infinite dimensional (even type II<sub>1</sub>) ancilla.

**Proof:** Consider following sets of  $n \times n$  complex matrices,  $n \geq 2$ :

$$\begin{aligned} \mathcal{G}_{\text{fin}}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary} \right. \\ &\quad \left. \text{finite dimens } C^*\text{-alg } (\mathcal{A}, \tau) \right\}, \\ \cap \\ \mathcal{G}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary finite} \right. \\ &\quad \left. \text{vN alg } (M, \tau) \right\}. \end{aligned}$$

► (Rørdam-M):  $\mathcal{G}_{\text{fin}}(n)$  is convex  $\forall n \geq 2$ , but **not** closed whenever  $n \geq 11$ .

**Note:** If  $B \in M_n(\mathbb{C})$  *correlation matrix* (pos definite, diag entries equal 1), the Schur multiplier  $T_B \in UCPT(n)$ . (Ricard '08): If  $B$  has **real** entries, then  $T_B$  is factorizable.

► (Haagerup-M '11): A Schur multiplier  $T_B$  is **factorizable** iff  $B \in \mathcal{G}(n)$ .  
 Furthermore,  $T_B \in \mathcal{FM}_{\text{fin}}(n)$  iff  $B \in \mathcal{G}_{\text{fin}}(n)$ .

The map  $B \mapsto T_B$  is an affine homeo, hence theorem follows.

**Theorem** (Rørdam-M '19):  $\mathcal{FM}_{\text{fin}}(n)$  is **not** closed, whenever  $n \geq 11$ . Moreover, for each such  $n$ , there exist factorizable quantum channels on  $M_n(\mathbb{C})$  which do require infinite dimensional (even type II<sub>1</sub>) ancilla.

**Proof:** Consider following sets of  $n \times n$  complex matrices,  $n \geq 2$ :

$$\begin{aligned} \mathcal{G}_{\text{fin}}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary} \right. \\ &\quad \left. \text{finite dimens } C^*\text{-alg } (\mathcal{A}, \tau) \right\}, \\ \cap \\ \mathcal{G}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary finite} \right. \\ &\quad \left. \text{vN alg } (M, \tau) \right\}. \end{aligned}$$

► (Rørdam-M):  $\mathcal{G}_{\text{fin}}(n)$  is convex  $\forall n \geq 2$ , but **not** closed whenever  $n \geq 11$ .

**Note:** If  $B \in M_n(\mathbb{C})$  *correlation matrix* (pos definite, diag entries equal 1), the Schur multiplier  $T_B \in UCPT(n)$ . (Ricard '08): If  $B$  has **real** entries, then  $T_B$  is factorizable.

► (Haagerup-M '11): A Schur multiplier  $T_B$  is factorizable iff  $B \in \mathcal{G}(n)$ . Furthermore,  $T_B \in \mathcal{FM}_{\text{fin}}(n)$  iff  $B \in \mathcal{G}_{\text{fin}}(n)$ .

The map  $B \mapsto T_B$  is an affine homeo, hence theorem follows.



**Theorem** (Rørdam-M '19):  $\mathcal{FM}_{\text{fin}}(n)$  is **not** closed, whenever  $n \geq 11$ . Moreover, for each such  $n$ , there exist factorizable quantum channels on  $M_n(\mathbb{C})$  which do require infinite dimensional (even type II<sub>1</sub>) ancilla.

**Proof:** Consider following sets of  $n \times n$  complex matrices,  $n \geq 2$ :

$$\begin{aligned} \mathcal{G}_{\text{fin}}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary} \right. \\ &\quad \left. \text{finite dimens } C^*\text{-alg } (\mathcal{A}, \tau) \right\}, \\ \cap \\ \mathcal{G}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary finite} \right. \\ &\quad \left. \text{vN alg } (M, \tau) \right\}. \end{aligned}$$

► (Rørdam-M):  $\mathcal{G}_{\text{fin}}(n)$  is convex  $\forall n \geq 2$ , but **not** closed whenever  $n \geq 11$ .

**Note:** If  $B \in M_n(\mathbb{C})$  *correlation matrix* (pos definite, diag entries equal 1), the Schur multiplier  $T_B \in UCPT(n)$ . (Ricard '08): If  $B$  has **real** entries, then  $T_B$  is factorizable.

► (Haagerup-M '11): A Schur multiplier  $T_B$  is **factorizable** **iff**  $B \in \mathcal{G}(n)$ . Furthermore,  $T_B \in \mathcal{FM}_{\text{fin}}(n)$  **iff**  $B \in \mathcal{G}_{\text{fin}}(n)$ .

The map  $B \mapsto T_B$  is an affine homeo, hence theorem follows.

**Theorem** (Rørdam-M '19):  $\mathcal{FM}_{\text{fin}}(n)$  is **not** closed, whenever  $n \geq 11$ .  
 Moreover, for each such  $n$ , there exist factorizable quantum channels on  $M_n(\mathbb{C})$  which do require infinite dimensional (even type II<sub>1</sub>) ancilla.

**Proof:** Consider following sets of  $n \times n$  complex matrices,  $n \geq 2$ :

$$\begin{aligned} \mathcal{G}_{\text{fin}}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary} \right. \\ &\quad \left. \text{finite dimens } C^*\text{-alg } (\mathcal{A}, \tau) \right\}, \\ \cap \\ \mathcal{G}(n) &= \left\{ [\tau(u_j^* u_i)] : u_1, \dots, u_n \text{ unitaries in arbitrary finite} \right. \\ &\quad \left. \text{vN alg } (M, \tau) \right\}. \end{aligned}$$

► (Rørdam-M):  $\mathcal{G}_{\text{fin}}(n)$  is convex  $\forall n \geq 2$ , but **not** closed whenever  $n \geq 11$ .

**Note:** If  $B \in M_n(\mathbb{C})$  *correlation matrix* (pos definite, diag entries equal 1), the Schur multiplier  $T_B \in UCPT(n)$ . (Ricard '08): If  $B$  has **real** entries, then  $T_B$  is factorizable.

► (Haagerup-M '11): A Schur multiplier  $T_B$  is **factorizable** **iff**  $B \in \mathcal{G}(n)$ .  
 Furthermore,  $T_B \in \mathcal{FM}_{\text{fin}}(n)$  **iff**  $B \in \mathcal{G}_{\text{fin}}(n)$ .

The map  $B \mapsto T_B$  is an affine homeo, hence theorem follows.



(Rørdam-M '20): A new view-point on factorizable channels, leading to further connections (and interesting open problems in  $C^*$ -algebras):

►  $\mathcal{FM}(n)$  is *parametrized by* simplex of tracial states  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .

More precisely, if  $\tau \in T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ , let

$$C_\tau(i, j; k, \ell) = n\tau(\iota_2(e_{k\ell})^* \iota_1(e_{ij})), \quad 1 \leq i, j, k, \ell \leq n,$$

where  $\iota_1, \iota_2: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C}) * M_n(\mathbb{C})$  are the canonical inclusions. Then  $C_\tau \in M_{n^2}(\mathbb{C})$  is *positive*, hence it is the Choi matrix of some c.p. lin map  $T_\tau: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ , which turns out to be a *factoriz* quantum channel!

In fact, the map  $\Phi: T(M_n(\mathbb{C}) * M_n(\mathbb{C})) \rightarrow \mathcal{FM}(n), \tau \mapsto \Phi(\tau) := T_\tau$  is an affine continuous surjection, satisfying, moreover,

$$\Phi(T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))) = \mathcal{FM}_{\text{fin}}(n),$$

where  $T_{\text{fin}}$  = tracial states that factor through finite dim.  $C^*$ -alg.

(Rørdam-M '20): A new view-point on factorizable channels, leading to further connections (and interesting open problems in  $C^*$ -algebras):

►  $\mathcal{FM}(n)$  is *parametrized by* simplex of tracial states  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .

More precisely, if  $\tau \in T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ , let

$$C_\tau(i, j; k, \ell) = n\tau(\iota_2(e_{k\ell})^* \iota_1(e_{ij})), \quad 1 \leq i, j, k, \ell \leq n,$$

where  $\iota_1, \iota_2: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C}) * M_n(\mathbb{C})$  are the canonical inclusions. Then  $C_\tau \in M_{n^2}(\mathbb{C})$  is *positive*, hence it is the Choi matrix of some c.p. lin map  $T_\tau: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ , which turns out to be a *factoriz* quantum channel!

In fact, the map  $\Phi: T(M_n(\mathbb{C}) * M_n(\mathbb{C})) \rightarrow \mathcal{FM}(n), \tau \mapsto \Phi(\tau) := T_\tau$  is an affine continuous surjection, satisfying, moreover,

$$\Phi(T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))) = \mathcal{FM}_{\text{fin}}(n),$$

where  $T_{\text{fin}}$  = tracial states that factor through finite dim.  $C^*$ -alg.

(Rørdam-M '20): A new view-point on factorizable channels, leading to further connections (and interesting open problems in  $C^*$ -algebras):

►  $\mathcal{FM}(n)$  is *parametrized by* simplex of tracial states  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .

More precisely, if  $\tau \in T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ , let

$$C_\tau(i, j; k, \ell) = n\tau(\iota_2(e_{k\ell})^* \iota_1(e_{ij})), \quad 1 \leq i, j, k, \ell \leq n,$$

where  $\iota_1, \iota_2: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C}) * M_n(\mathbb{C})$  are the canonical inclusions. Then  $C_\tau \in M_{n^2}(\mathbb{C})$  is *positive*, hence it is the Choi matrix of some c.p. lin map  $T_\tau: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ , which turns out to be a *factoriz* quantum channel!

In fact, the map  $\Phi: T(M_n(\mathbb{C}) * M_n(\mathbb{C})) \rightarrow \mathcal{FM}(n), \tau \mapsto \Phi(\tau) := T_\tau$  is an affine continuous surjection, satisfying, moreover,

$$\Phi(T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))) = \mathcal{FM}_{\text{fin}}(n),$$

where  $T_{\text{fin}}$  = tracial states that factor through finite dim.  $C^*$ -alg.

The affine cont surj  $\Phi: T(M_n(\mathbb{C}) * M_n(\mathbb{C})) \rightarrow \mathcal{FM}(n), \tau \mapsto T_\tau$ , satisfies

- $\Phi(T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))) = \mathcal{FM}_{\text{fin}}(n),$
- $\Phi(\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C})))) = \overline{\mathcal{FM}_{\text{fin}}(n)},$

where  $T_{\text{fin}}$  = tracial states that factor through finite dim.  $C^*$ -alg.

**Thm** (Haagerup-M '15) Connes Embedding Problem (CEP) has positive answer iff  $\overline{\mathcal{FM}_{\text{fin}}(n)} = \mathcal{FM}(n), \forall n \geq 3.$

**Question:** What can we say about  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))}$  ?

- (Exel–Loring '92):  $M_n(\mathbb{C}) * M_n(\mathbb{C})$  residually finite dim. (RFD)
- (Blackadar '85):  $M_n(\mathbb{C}) * M_n(\mathbb{C})$  semi-projective.

The affine cont surj  $\Phi: T(M_n(\mathbb{C}) * M_n(\mathbb{C})) \rightarrow \mathcal{FM}(n), \tau \mapsto T_\tau$ , satisfies

- $\Phi(T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))) = \mathcal{FM}_{\text{fin}}(n)$ ,
- $\Phi(\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C})))) = \overline{\mathcal{FM}_{\text{fin}}(n)}$ ,

where  $T_{\text{fin}}$  = tracial states that factor through finite dim.  $C^*$ -alg.

**Thm** (Haagerup-M '15) Connes Embedding Problem (CEP) has positive answer **iff**  $\overline{\mathcal{FM}_{\text{fin}}(n)} = \mathcal{FM}(n), \forall n \geq 3$ .

**Question:** What can we say about  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))}$  ?

- (Exel–Loring '92):  $M_n(\mathbb{C}) * M_n(\mathbb{C})$  residually finite dim. (RFD)
- (Blackadar '85):  $M_n(\mathbb{C}) * M_n(\mathbb{C})$  semi-projective.

The affine cont surj  $\Phi: T(M_n(\mathbb{C}) * M_n(\mathbb{C})) \rightarrow \mathcal{FM}(n), \tau \mapsto T_\tau$ , satisfies

- $\Phi(T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))) = \mathcal{FM}_{\text{fin}}(n),$
- $\Phi(\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C})))) = \overline{\mathcal{FM}_{\text{fin}}(n)},$

where  $T_{\text{fin}}$  = tracial states that factor through finite dim.  $C^*$ -alg.

**Thm** (Haagerup-M '15) Connes Embedding Problem (CEP) has positive answer **iff**  $\overline{\mathcal{FM}_{\text{fin}}(n)} = \mathcal{FM}(n), \forall n \geq 3.$

**Question:** What can we say about  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))}$  ?

- (Exel–Loring '92):  $M_n(\mathbb{C}) * M_n(\mathbb{C})$  **residually finite dim.** (RFD)
- (Blackadar '85):  $M_n(\mathbb{C}) * M_n(\mathbb{C})$  **semi-projective.**



In general, given  $A = (\text{sep})$  unital tracial  $C^*$ -algebra, we have inclusions:

$$T_{\text{fin}}(A) \subseteq \overline{T_{\text{fin}}(A)} \subseteq T_{\text{qd}}(A) \subseteq T_{\text{am}}(A) \subseteq T_{\text{hyp}}(A) \subseteq T(A),$$

where  $T_{\text{qd}}(A)$  = quasi-diagonal traces,  $T_{\text{am}}(A)$  = amenable (liftable) traces,  $T_{\text{hyp}}(A)$  = hyperlinear traces (i.e., traces  $\tau$  st  $\pi_\tau(A)'' \hookrightarrow \mathcal{R}^\omega$ ).

- CEP pos answer iff  $T_{\text{hyp}}(A) = T(A)$ , for all  $C^*$ -alg  $A$ .

**Reformulation of CEP:** For all sep. unital tracial  $C^*$ -algs  $(A, \tau)$ , there is a unital trace-preserving  $*$ -hom  $\varphi: A \rightarrow \prod_{n=1}^{\infty} M_{k_n}/I^\omega$ , for some  $k_n \geq 1$ .

- (N. Brown '06):  $\exists$  exact RFD  $C^*$ -alg  $A$  s.t.  $T_{\text{am}}(A) \neq T_{\text{hyp}}(A)$ .
- **Open** if  $T_{\text{qd}}(A) = T_{\text{am}}(A)$ . Strong pos results: Tikuisis-Winter-White, Schafhauser, Gabe.
- $A$  (weakly) semi-projective  $\implies \overline{T_{\text{fin}}(A)} = T_{\text{qd}}(A)$
- (Hadwin–Shulman '17):  $\exists$  RFD  $C^*$ -alg  $A$  s.t.  $\overline{T_{\text{fin}}(A)} \neq T_{\text{qd}}(A)$ .

**Thm** (Rørdam-M '20):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .



In general, given  $A = (\text{sep})$  unital tracial  $C^*$ -algebra, we have inclusions:

$$T_{\text{fin}}(A) \subseteq \overline{T_{\text{fin}}(A)} \subseteq T_{\text{qd}}(A) \subseteq T_{\text{am}}(A) \subseteq T_{\text{hyp}}(A) \subseteq T(A),$$

where  $T_{\text{qd}}(A)$  = quasi-diagonal traces,  $T_{\text{am}}(A)$  = amenable (liftable) traces,  $T_{\text{hyp}}(A)$  = hyperlinear traces (i.e., traces  $\tau$  st  $\pi_\tau(A)'' \hookrightarrow \mathcal{R}^\omega$ ).

- CEP pos answer **iff**  $T_{\text{hyp}}(A) = T(A)$ , for all  $C^*$ -alg  $A$ .

**Reformulation of CEP:** For all sep. unital tracial  $C^*$ -algs  $(A, \tau)$ , there is a unital trace-preserving  $*$ -hom  $\varphi: A \rightarrow \prod_{n=1}^{\infty} M_{k_n}/I^\omega$ , for some  $k_n \geq 1$ .

- (N. Brown '06):  $\exists$  exact RFD  $C^*$ -alg  $A$  s.t.  $T_{\text{am}}(A) \neq T_{\text{hyp}}(A)$ .
- **Open** if  $T_{\text{qd}}(A) = T_{\text{am}}(A)$ . Strong pos results: Tikuisis-Winter-White, Schafhauser, Gabe.
- $A$  (weakly) semi-projective  $\implies \overline{T_{\text{fin}}(A)} = T_{\text{qd}}(A)$
- (Hadwin–Shulman '17):  $\exists$  RFD  $C^*$ -alg  $A$  s.t.  $\overline{T_{\text{fin}}(A)} \neq T_{\text{qd}}(A)$ .

**Thm** (Rørdam-M '20):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .



In general, given  $A = (\text{sep})$  unital tracial  $C^*$ -algebra, we have inclusions:

$$T_{\text{fin}}(A) \subseteq \overline{T_{\text{fin}}(A)} \subseteq T_{\text{qd}}(A) \subseteq T_{\text{am}}(A) \subseteq T_{\text{hyp}}(A) \subseteq T(A),$$

where  $T_{\text{qd}}(A)$  = quasi-diagonal traces,  $T_{\text{am}}(A)$  = amenable (liftable) traces,  $T_{\text{hyp}}(A)$  = hyperlinear traces (i.e., traces  $\tau$  st  $\pi_\tau(A)'' \hookrightarrow \mathcal{R}^\omega$ ).

- CEP pos answer **iff**  $T_{\text{hyp}}(A) = T(A)$ , for all  $C^*$ -alg  $A$ .

**Reformulation of CEP:** For all sep. unital tracial  $C^*$ -algs  $(A, \tau)$ , there is a unital trace-preserving  $*$ -hom  $\varphi: A \rightarrow \prod_{n=1}^{\infty} M_{k_n}/I^\omega$ , for some  $k_n \geq 1$ .

- (N. Brown '06):  $\exists$  exact RFD  $C^*$ -alg  $A$  s.t.  $T_{\text{am}}(A) \neq T_{\text{hyp}}(A)$ .
- **Open** if  $T_{\text{qd}}(A) = T_{\text{am}}(A)$ . Strong pos results: Tikuisis-Winter-White, Schafhauser, Gabe.
- $A$  (weakly) semi-projective  $\implies \overline{T_{\text{fin}}(A)} = T_{\text{qd}}(A)$
- (Hadwin–Shulman '17):  $\exists$  RFD  $C^*$ -alg  $A$  s.t.  $\overline{T_{\text{fin}}(A)} \neq T_{\text{qd}}(A)$ .

**Thm** (Rørdam-M '20):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .



In general, given  $A = (\text{sep})$  unital tracial  $C^*$ -algebra, we have inclusions:

$$T_{\text{fin}}(A) \subseteq \overline{T_{\text{fin}}(A)} \subseteq T_{\text{qd}}(A) \subseteq T_{\text{am}}(A) \subseteq T_{\text{hyp}}(A) \subseteq T(A),$$

where  $T_{\text{qd}}(A)$  = quasi-diagonal traces,  $T_{\text{am}}(A)$  = amenable (liftable) traces,  $T_{\text{hyp}}(A)$  = hyperlinear traces (i.e., traces  $\tau$  st  $\pi_\tau(A)'' \hookrightarrow \mathcal{R}^\omega$ ).

- CEP pos answer **iff**  $T_{\text{hyp}}(A) = T(A)$ , for all  $C^*$ -alg  $A$ .

**Reformulation of CEP:** For all sep. unital tracial  $C^*$ -algs  $(A, \tau)$ , there is a unital trace-preserving  $*$ -hom  $\varphi: A \rightarrow \prod_{n=1}^{\infty} M_{k_n}/I^\omega$ , for some  $k_n \geq 1$ .

- (N. Brown '06):  $\exists$  exact RFD  $C^*$ -alg  $A$  s.t.  $T_{\text{am}}(A) \neq T_{\text{hyp}}(A)$ .
- **Open** if  $T_{\text{qd}}(A) = T_{\text{am}}(A)$ . Strong pos results: Tikuisis-Winter-White, Schafhauser, Gabe.
- $A$  (weakly) semi-projective  $\implies \overline{T_{\text{fin}}(A)} = T_{\text{qd}}(A)$
- (Hadwin-Shulman '17):  $\exists$  RFD  $C^*$ -alg  $A$  s.t.  $\overline{T_{\text{fin}}(A)} \neq T_{\text{qd}}(A)$ .

**Thm** (Rørdam-M '20):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .



In general, given  $A = (\text{sep})$  unital tracial  $C^*$ -algebra, we have inclusions:

$$T_{\text{fin}}(A) \subseteq \overline{T_{\text{fin}}(A)} \subseteq T_{\text{qd}}(A) \subseteq T_{\text{am}}(A) \subseteq T_{\text{hyp}}(A) \subseteq T(A),$$

where  $T_{\text{qd}}(A)$  = quasi-diagonal traces,  $T_{\text{am}}(A)$  = amenable (liftable) traces,  $T_{\text{hyp}}(A)$  = hyperlinear traces (i.e., traces  $\tau$  st  $\pi_\tau(A)'' \hookrightarrow \mathcal{R}^\omega$ ).

- CEP pos answer **iff**  $T_{\text{hyp}}(A) = T(A)$ , for all  $C^*$ -alg  $A$ .

**Reformulation of CEP:** For all sep. unital tracial  $C^*$ -algs  $(A, \tau)$ , there is a unital trace-preserving  $*$ -hom  $\varphi: A \rightarrow \prod_{n=1}^{\infty} M_{k_n}/I^\omega$ , for some  $k_n \geq 1$ .

- (N. Brown '06):  $\exists$  exact RFD  $C^*$ -alg  $A$  s.t.  $T_{\text{am}}(A) \neq T_{\text{hyp}}(A)$ .
- **Open** if  $T_{\text{qd}}(A) = T_{\text{am}}(A)$ . Strong pos results: Tikuisis-Winter-White, Schafhauser, Gabe.
- $A$  (weakly) semi-projective  $\implies \overline{T_{\text{fin}}(A)} = T_{\text{qd}}(A)$
- (Hadwin-Shulman '17):  $\exists$  RFD  $C^*$ -alg  $A$  s.t.  $\overline{T_{\text{fin}}(A)} \neq T_{\text{qd}}(A)$ .

**Thm** (Rørdam-M '20):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .



In general, given  $A = (\text{sep})$  unital tracial  $C^*$ -algebra, we have inclusions:

$$T_{\text{fin}}(A) \subseteq \overline{T_{\text{fin}}(A)} \subseteq T_{\text{qd}}(A) \subseteq T_{\text{am}}(A) \subseteq T_{\text{hyp}}(A) \subseteq T(A),$$

where  $T_{\text{qd}}(A)$  = quasi-diagonal traces,  $T_{\text{am}}(A)$  = amenable (liftable) traces,  $T_{\text{hyp}}(A)$  = hyperlinear traces (i.e., traces  $\tau$  st  $\pi_\tau(A)'' \hookrightarrow \mathcal{R}^\omega$ ).

- CEP pos answer **iff**  $T_{\text{hyp}}(A) = T(A)$ , for all  $C^*$ -alg  $A$ .

**Reformulation of CEP:** For all sep. unital tracial  $C^*$ -algs  $(A, \tau)$ , there is a unital trace-preserving  $*$ -hom  $\varphi: A \rightarrow \prod_{n=1}^{\infty} M_{k_n}/I^\omega$ , for some  $k_n \geq 1$ .

- (N. Brown '06):  $\exists$  exact RFD  $C^*$ -alg  $A$  s.t.  $T_{\text{am}}(A) \neq T_{\text{hyp}}(A)$ .
- **Open** if  $T_{\text{qd}}(A) = T_{\text{am}}(A)$ . Strong pos results: Tikuisis-Winter-White, Schafhauser, Gabe.
- $A$  (weakly) semi-projective  $\implies \overline{T_{\text{fin}}(A)} = T_{\text{qd}}(A)$
- (Hadwin–Shulman '17):  $\exists$  RFD  $C^*$ -alg  $A$  s.t.  $\overline{T_{\text{fin}}(A)} \neq T_{\text{qd}}(A)$ .

**Thm** (Rørdam-M '20):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .



In general, given  $A = (\text{sep})$  unital tracial  $C^*$ -algebra, we have inclusions:

$$T_{\text{fin}}(A) \subseteq \overline{T_{\text{fin}}(A)} \subseteq T_{\text{qd}}(A) \subseteq T_{\text{am}}(A) \subseteq T_{\text{hyp}}(A) \subseteq T(A),$$

where  $T_{\text{qd}}(A)$  = quasi-diagonal traces,  $T_{\text{am}}(A)$  = amenable (liftable) traces,  $T_{\text{hyp}}(A)$  = hyperlinear traces (i.e., traces  $\tau$  st  $\pi_\tau(A)'' \hookrightarrow \mathcal{R}^\omega$ ).

- CEP pos answer **iff**  $T_{\text{hyp}}(A) = T(A)$ , for all  $C^*$ -alg  $A$ .

**Reformulation of CEP:** For all sep. unital tracial  $C^*$ -algs  $(A, \tau)$ , there is a unital trace-preserving  $*$ -hom  $\varphi: A \rightarrow \prod_{n=1}^{\infty} M_{k_n}/I^\omega$ , for some  $k_n \geq 1$ .

- (N. Brown '06):  $\exists$  exact RFD  $C^*$ -alg  $A$  s.t.  $T_{\text{am}}(A) \neq T_{\text{hyp}}(A)$ .
- **Open** if  $T_{\text{qd}}(A) = T_{\text{am}}(A)$ . Strong pos results: Tikuisis-Winter-White, Schafhauser, Gabe.
- $A$  (weakly) semi-projective  $\implies \overline{T_{\text{fin}}(A)} = T_{\text{qd}}(A)$
- (Hadwin–Shulman '17):  $\exists$  RFD  $C^*$ -alg  $A$  s.t.  $\overline{T_{\text{fin}}(A)} \neq T_{\text{qd}}(A)$ .

**Thm** (Rørdam-M '20):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ .



**Thm** (Rørdam-M):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})).$

**Cor:** CEP pos iff  $T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})) = T(M_n(\mathbb{C}) * M_n(\mathbb{C})), \forall n \geq 3.$

**Further results:** Let  $A$  be a unital  $C^*$ -algebra.

- If  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ , then  $A$  generated by  $n^2$  elem.
- If  $A$  gen by  $n - 1$  unitaries, then  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ .

As unital separable  $\mathcal{Z}$ -stable  $C^*$ -alg are singly generated (Thiel-Winter), we deduce: If  $A$  is a **simple unital inf dim AF-algebra**, then  $M_n(A)$  is a **quotient** of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ ,  $n \geq 3$ .

**Thm** (Rørdam-M): Each metrizable Choquet simplex is affinely homeo to a face of  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ ,  $n \geq 3$ .

**(Open) question:** Is  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$  the Poulsen simplex?



**Thm** (Rørdam-M):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})).$

**Cor:** CEP pos iff  $T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})) = T(M_n(\mathbb{C}) * M_n(\mathbb{C})), \forall n \geq 3.$

**Further results:** Let  $A$  be a unital  $C^*$ -algebra.

- If  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ , then  $A$  generated by  $n^2$  elem.
- If  $A$  gen by  $n - 1$  unitaries, then  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ .

As unital separable  $\mathcal{Z}$ -stable  $C^*$ -alg are singly generated (Thiel-Winter), we deduce: If  $A$  is a **simple unital inf dim AF-algebra**, then  $M_n(A)$  is a **quotient** of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ ,  $n \geq 3$ .

**Thm** (Rørdam-M): Each metrizable Choquet simplex is affinely homeo to a face of  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ ,  $n \geq 3$ .

**(Open) question:** Is  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$  the Poulsen simplex?

**Thm** (Rørdam-M):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})).$

**Cor:** CEP pos iff  $T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})) = T(M_n(\mathbb{C}) * M_n(\mathbb{C})), \forall n \geq 3.$

**Further results:** Let  $A$  be a unital  $C^*$ -algebra.

- If  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ , then  $A$  generated by  $n^2$  elem.
- If  $A$  gen by  $n - 1$  unitaries, then  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ .

As unital separable  $\mathcal{Z}$ -stable  $C^*$ -alg are singly generated (Thiel-Winter), we deduce: If  $A$  is a **simple unital inf dim AF-algebra**, then  $M_n(A)$  is a **quotient** of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ ,  $n \geq 3$ .

**Thm** (Rørdam-M): Each metrizable Choquet simplex is affinely homeo to a face of  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ ,  $n \geq 3$ .

**(Open) question:** Is  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$  the Poulsen simplex?

**Thm** (Rørdam-M):  $\overline{T_{\text{fin}}(M_n(\mathbb{C}) * M_n(\mathbb{C}))} = T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})).$

**Cor:** CEP pos iff  $T_{\text{hyp}}(M_n(\mathbb{C}) * M_n(\mathbb{C})) = T(M_n(\mathbb{C}) * M_n(\mathbb{C})), \forall n \geq 3.$

**Further results:** Let  $A$  be a unital  $C^*$ -algebra.

- If  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ , then  $A$  generated by  $n^2$  elem.
- If  $A$  gen by  $n - 1$  unitaries, then  $M_n(A)$  is a quotient of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ .

As unital separable  $\mathcal{Z}$ -stable  $C^*$ -alg are singly generated (Thiel-Winter), we deduce: If  $A$  is a **simple unital inf dim AF-algebra**, then  $M_n(A)$  is a **quotient** of  $M_n(\mathbb{C}) * M_n(\mathbb{C})$ ,  $n \geq 3$ .

**Thm** (Rørdam-M): Each metrizable Choquet simplex is affinely homeo to a face of  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$ ,  $n \geq 3$ .

**(Open) question:** Is  $T(M_n(\mathbb{C}) * M_n(\mathbb{C}))$  the Poulsen simplex?

## Further connections: Analysis of quantum correlations and CEP

- $\Gamma := \mathbb{Z}_k * \mathbb{Z}_k * \cdots * \mathbb{Z}_k$ ,  $n$  copies,  $n, k \geq 2$ .
- $C^*(\Gamma) = C^*(q_{j,x} \mid q_{j,x} = q_{j,x}^* = q_{j,x}^2, \sum_{j=1}^k q_{j,x} = 1)$ .

(Schafhauser, AIM '21): For all  $n, k \geq 2$ , we have

$$\overline{T_{\text{fin}}(C^*(\Gamma))} = T_{\text{hyp}}(C^*(\Gamma)).$$

**Definition:** A "correlation"  $[(p(i,j \mid x,y))]$  is *synchronous* if  $\forall 1 \leq x \leq n$ ,  $p(i,j \mid x,x) = 0$  whenever  $i \neq j$ .

**Theorem** (Paulsen-Severini-Stalke-Todorov-Winter '16):

$$\begin{aligned} C_{qc}^s(n, k) &= \left\{ [\tau(q_{j,x} q_{i,y})]_{(i,x;j,y)} \mid \tau \in T(C^*(\Gamma)) \right\} \\ C_q^s(n, k) &= \left\{ [\tau(q_{j,x} q_{i,y})]_{(i,x;j,y)} \mid \tau \in T_{\text{fin}}(C^*(\Gamma)) \right\}. \end{aligned}$$

## Further connections: Analysis of quantum correlations and CEP

- $\Gamma := \mathbb{Z}_k * \mathbb{Z}_k * \cdots * \mathbb{Z}_k$ ,  $n$  copies,  $n, k \geq 2$ .
- $C^*(\Gamma) = C^*(q_{j,x} \mid q_{j,x} = q_{j,x}^* = q_{j,x}^2, \sum_{j=1}^k q_{j,x} = 1)$ .

(Schafhauser, AIM '21): For all  $n, k \geq 2$ , we have

$$\overline{T_{\text{fin}}(C^*(\Gamma))} = T_{\text{hyp}}(C^*(\Gamma)).$$

**Definition:** A "correlation"  $[(p(i,j \mid x,y))]$  is *synchronous* if  $\forall 1 \leq x \leq n$ ,  $p(i,j \mid x,x) = 0$  whenever  $i \neq j$ .

**Theorem** (Paulsen-Severini-Stalke-Todorov-Winter '16):

$$\begin{aligned} C_{qc}^s(n, k) &= \left\{ [\tau(q_{j,x} q_{i,y})]_{(i,x;j,y)} \mid \tau \in T(C^*(\Gamma)) \right\} \\ C_q^s(n, k) &= \left\{ [\tau(q_{j,x} q_{i,y})]_{(i,x;j,y)} \mid \tau \in T_{\text{fin}}(C^*(\Gamma)) \right\}. \end{aligned}$$

## Further connections: Analysis of quantum correlations and CEP

- $\Gamma := \mathbb{Z}_k * \mathbb{Z}_k * \cdots * \mathbb{Z}_k$ ,  $n$  copies,  $n, k \geq 2$ .
- $C^*(\Gamma) = C^*(q_{j,x} \mid q_{j,x} = q_{j,x}^* = q_{j,x}^2, \sum_{j=1}^k q_{j,x} = 1)$ .

(Schafhauser, AIM '21): For all  $n, k \geq 2$ , we have

$$\overline{T_{\text{fin}}(C^*(\Gamma))} = T_{\text{hyp}}(C^*(\Gamma)).$$

**Definition:** A "correlation"  $[(p(i, j \mid x, y))]$  is *synchronous* if  $\forall 1 \leq x \leq n$ ,  $p(i, j \mid x, x) = 0$  whenever  $i \neq j$ .

**Theorem** (Paulsen-Severini-Stalke-Todorov-Winter '16):

$$\begin{aligned} C_{qc}^s(n, k) &= \left\{ [\tau(q_{j,x} q_{i,y})]_{(i,x;j,y)} \mid \tau \in T(C^*(\Gamma)) \right\} \\ C_q^s(n, k) &= \left\{ [\tau(q_{j,x} q_{i,y})]_{(i,x;j,y)} \mid \tau \in T_{\text{fin}}(C^*(\Gamma)) \right\}. \end{aligned}$$

Set  $C_{qa}(n, k) := \overline{C_q(n, k)}$ .

(Kim-Paulsen-Schafhauser '17):  $C_{qa}^s(n, k) = \overline{C_q^s(n, k)}$ .

**Theorem** (Kim-Paulsen-Schafhauser '17, Ozawa '13): TFAE

- (1) Connes Embedding Problem has positive answer.
- (2)  $C_{qa}^s(n, k) = C_{qc}^s(n, k)$ ,  $\forall n, k \geq 2$ .
- (3) Tsirelson's conjecture is true, i.e.,  $C_{qa}(n, k) = C_{qc}(n, k)$ ,  $\forall n, k \geq 2$ .

**Theorem** (Fritz/Junge et. al. '09):

- $C_{qa}(n, k) = \left\{ \left[ \varphi(q_{j,x} \otimes q_{i,y}) \right] : \varphi \text{ state on } C^*(\Gamma) \otimes_{\min} C^*(\Gamma) \right\}$ .
  - $C_{qc}(n, k) = \left\{ \left[ \varphi(q_{j,x} \otimes q_{i,y}) \right] : \varphi \text{ state on } C^*(\Gamma) \otimes_{\max} C^*(\Gamma) \right\}$ .
- $A := C^*(\Gamma) \otimes_{\min} C^*(\Gamma)$  is RFD  $[\Rightarrow S_{\text{fin}}(A) \stackrel{\text{dense}}{\subseteq} S(A)]$ .

Set  $C_{qa}(n, k) := \overline{C_q(n, k)}$ .

(Kim-Paulsen-Schafhauser '17):  $C_{qa}^s(n, k) = \overline{C_q^s(n, k)}$ .

**Theorem** (Kim-Paulsen-Schafhauser '17, Ozawa '13): TFAE

- (1) Connes Embedding Problem has positive answer.
- (2)  $C_{qa}^s(n, k) = C_{qc}^s(n, k)$ ,  $\forall n, k \geq 2$ .
- (3) Tsirelson's conjecture is true, i.e.,  $C_{qa}(n, k) = C_{qc}(n, k)$ ,  $\forall n, k \geq 2$ .

**Theorem** (Fritz/Junge et. al. '09):

- $C_{qa}(n, k) = \left\{ \left[ \varphi(q_{j,x} \otimes q_{i,y}) \right] : \varphi \text{ state on } C^*(\Gamma) \otimes_{\min} C^*(\Gamma) \right\}$ .
- $C_{qc}(n, k) = \left\{ \left[ \varphi(q_{j,x} \otimes q_{i,y}) \right] : \varphi \text{ state on } C^*(\Gamma) \otimes_{\max} C^*(\Gamma) \right\}$ .

- $A := C^*(\Gamma) \otimes_{\min} C^*(\Gamma)$  is RFD  $[\Rightarrow S_{\text{fin}}(A) \stackrel{\text{dense}}{\subseteq} S(A)]$ .



Set  $C_{qa}(n, k) := \overline{C_q(n, k)}$ .

(Kim-Paulsen-Schafhauser '17):  $C_{qa}^s(n, k) = \overline{C_q^s(n, k)}$ .

**Theorem** (Kim-Paulsen-Schafhauser '17, Ozawa '13): TFAE

- (1) Connes Embedding Problem has positive answer.
- (2)  $C_{qa}^s(n, k) = C_{qc}^s(n, k)$ ,  $\forall n, k \geq 2$ .
- (3) Tsirelson's conjecture is true, i.e.,  $C_{qa}(n, k) = C_{qc}(n, k)$ ,  $\forall n, k \geq 2$ .

**Theorem** (Fritz/Junge et. al. '09):

- $C_{qa}(n, k) = \left\{ \left[ \varphi(q_{j,x} \otimes q_{i,y}) \right] : \varphi \text{ state on } C^*(\Gamma) \otimes_{\min} C^*(\Gamma) \right\}$ .
- $C_{qc}(n, k) = \left\{ \left[ \varphi(q_{j,x} \otimes q_{i,y}) \right] : \varphi \text{ state on } C^*(\Gamma) \otimes_{\max} C^*(\Gamma) \right\}$ .

- $A := C^*(\Gamma) \otimes_{\min} C^*(\Gamma)$  is RFD  $[\Rightarrow S_{\text{fin}}(A) \stackrel{\text{dense}}{\subseteq} S(A)]$ .

► **Posted on arXiv, Jan 2020:**  $MIP^* = RE$ , Ji, Natarajan, Vidick, Wright, Yuen, 165 pp.

Proving that the complexity class  $MIP^*$  (quantum version of complexity class  $MIP$ =languages with a Multiprover Interactive Proof) contains an undecidable language, they conclude that Tsirelson's Conjecture is **false!**

New version (with corrections) 206 pp., posted on arXiv, Sept 2020.

► Further recent applications of/connections to factorizability:

Gangbo-Jeckel-Nam-Shlyakhtenko, May 2021: “Duality for optimal couplings in free probability”.

► **Posted on arXiv, Jan 2020:**  $MIP^* = RE$ , Ji, Natarajan, Vidick, Wright, Yuen, 165 pp.

Proving that the complexity class  $MIP^*$  (quantum version of complexity class  $MIP$ =languages with a Multiprover Interactive Proof) contains an undecidable language, they conclude that Tsirelson's Conjecture is **false!**

New version (with corrections) 206 pp., posted on arXiv, Sept 2020.

► Further recent applications of/connections to factorizability:

Gangbo-Jeckel-Nam-Shlyakhtenko, May 2021: “Duality for optimal couplings in free probability”.