

# The ideal intersection property for essential groupoid $C^*$ -algebras

Matthew Kennedy  
joint work with Se-Jin Kim, Xin Li, Sven Raum and Dan Ursu

University of Waterloo, Waterloo, Canada

June 1, 2022

# Advertisement

## Fields Thematic Program on Operator Algebras and Applications

**Dates:** July 1 - December 31, 2023



**Workshops include:** Connections with Logic, Symmetry and Structure, Groups and Group Actions, Free Probability, Non-Commutative Geometry

# Motivation

Let  $A$  be a  $C^*$ -algebra constructed in terms of some “combinatorial” data.  
For example, a group or group action.

# Motivation

Let  $A$  be a  $C^*$ -algebra constructed in terms of some “combinatorial” data. For example, a group or group action.

**Basic question:** When is  $A$  simple? Prefer an answer in terms of the data.

# Motivation

Let  $A$  be a  $C^*$ -algebra constructed in terms of some “combinatorial” data. For example, a group or group action.

**Basic question:** When is  $A$  simple? Prefer an answer in terms of the data.

## Theorem (Murray-von Neumann 1936)

*For a discrete group  $G$ , the group von Neumann algebra  $L(G)$  is simple (i.e. factorial) iff  $G$  is ICC.*

# Reduced discrete group $C^*$ -algebras

**Theorem (Powers 1975, ..., Olshanski-Osin 2014)**

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if  $G$  satisfies a “Powers-type” condition.*

# Reduced discrete group $C^*$ -algebras

## Theorem (Powers 1975, ..., Olshanski-Osin 2014)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if  $G$  satisfies a “Powers-type” condition.*

Note: First established by Powers for  $\mathbb{F}_2$ . Culminating in work of Olshanskii-Osin for certain free Burnside groups.

# Reduced discrete group $C^*$ -algebras

## Theorem (Powers 1975, ..., Olshanski-Osin 2014)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if  $G$  satisfies a “Powers-type” condition.*

Note: First established by Powers for  $\mathbb{F}_2$ . Culminating in work of Olshanskii-Osin for certain free Burnside groups.

## Theorem (Kalanatar-K 2017)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if and only if  $G$  has a topologically free boundary action.*



# Reduced discrete group $C^*$ -algebras

## Theorem (Powers 1975, ..., Olshanski-Osin 2014)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if  $G$  satisfies a “Powers-type” condition.*

Note: First established by Powers for  $\mathbb{F}_2$ . Culminating in work of Olshanskii-Osin for certain free Burnside groups.

## Theorem (Kalanatar-K 2017)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if and only if  $G$  has a topologically free boundary action.*

Note: Equivalent to freeness of the Furstenberg boundary  $\partial_F G$ .

# Reduced discrete group $C^*$ -algebras

## Theorem (Powers 1975, ..., Olshanski-Osin 2014)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if  $G$  satisfies a “Powers-type” condition.*

Note: First established by Powers for  $\mathbb{F}_2$ . Culminating in work of Olshanskii-Osin for certain free Burnside groups.

## Theorem (Kalanatar-K 2017)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if and only if  $G$  has a topologically free boundary action.*

Note: Equivalent to freeness of the Furstenberg boundary  $\partial_F G$ .

## Theorem (K 2020)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if and only if  $G$  has no amenable confined subgroups.*

# Reduced discrete group $C^*$ -algebras

## Theorem (Powers 1975, ..., Olshanski-Osin 2014)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if  $G$  satisfies a “Powers-type” condition.*

Note: First established by Powers for  $\mathbb{F}_2$ . Culminating in work of Olshanskii-Osin for certain free Burnside groups.

## Theorem (Kalanatar-K 2017)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if and only if  $G$  has a topologically free boundary action.*

Note: Equivalent to freeness of the Furstenberg boundary  $\partial_F G$ .

## Theorem (K 2020)

*The  $C^*$ -algebra  $C_\lambda^*(G)$  is simple if and only if  $G$  has no amenable confined subgroups.*

Note:  $H \leq G$  is confined if it is non-trivial and “almost normal” in the sense that

$$1 \notin \overline{\{gHg^{-1}\}}.$$

# Reduced discrete crossed product $C^*$ -algebras

## Theorem (Elliott 1980, ..., Archbold-Spielberg 1993)

*If  $G$  is amenable then  $C(X) \rtimes_\lambda G$  is simple if and only if  $G \curvearrowright X$  is minimal and topologically free.*

# Reduced discrete crossed product $C^*$ -algebras

## Theorem (Elliott 1980, ..., Archbold-Spielberg 1993)

*If  $G$  is amenable then  $C(X) \rtimes_\lambda G$  is simple if and only if  $G \curvearrowright X$  is minimal and topologically free.*

## Theorem (Kawabe 2017)

*The  $C^*$ -algebra  $C(X) \rtimes_\lambda G$  is simple if and only if  $X$  is minimal and for every  $x \in X$ ,  $G_x$  has no amenable confined subgroups.*

# Reduced discrete crossed product $C^*$ -algebras

## Theorem (Elliott 1980, ..., Archbold-Spielberg 1993)

*If  $G$  is amenable then  $C(X) \rtimes_\lambda G$  is simple if and only if  $G \curvearrowright X$  is minimal and topologically free.*

## Theorem (Kawabe 2017)

*The  $C^*$ -algebra  $C(X) \rtimes_\lambda G$  is simple if and only if  $X$  is minimal and for every  $x \in X$ ,  $G_x$  has no amenable confined subgroups.*

Note: Here  $G_x$  is the stabilizer subgroup

$$G_x = \{g \in G : gx = x\}.$$

# Reduced discrete crossed product $C^*$ -algebras

## Theorem (Elliott 1980, ..., Archbold-Spielberg 1993)

*If  $G$  is amenable then  $C(X) \rtimes_\lambda G$  is simple if and only if  $G \curvearrowright X$  is minimal and topologically free.*

## Theorem (Kawabe 2017)

*The  $C^*$ -algebra  $C(X) \rtimes_\lambda G$  is simple if and only if  $X$  is minimal and for every  $x \in X$ ,  $G_x$  has no amenable confined subgroups.*

Note: Here  $G_x$  is the stabilizer subgroup

$$G_x = \{g \in G : gx = x\}.$$

## Theorem (Kalantar-Scarparo 2021)

*The  $C^*$ -algebra  $C_0(X) \rtimes_\lambda G$  is simple if and only if  $X$  is minimal and for every  $x \in X$ ,  $G_x$  has no amenable confined subgroups.*

# Reduced discrete crossed product $C^*$ -algebras

## Theorem (Elliott 1980, ..., Archbold-Spielberg 1993)

*If  $G$  is amenable then  $C(X) \rtimes_\lambda G$  is simple if and only if  $G \curvearrowright X$  is minimal and topologically free.*

## Theorem (Kawabe 2017)

*The  $C^*$ -algebra  $C(X) \rtimes_\lambda G$  is simple if and only if  $X$  is minimal and for every  $x \in X$ ,  $G_x$  has no amenable confined subgroups.*

Note: Here  $G_x$  is the stabilizer subgroup

$$G_x = \{g \in G : gx = x\}.$$

## Theorem (Kalantar-Scarparo 2021)

*The  $C^*$ -algebra  $C_0(X) \rtimes_\lambda G$  is simple if and only if  $X$  is minimal and for every  $x \in X$ ,  $G_x$  has no amenable confined subgroups.*

Note: Key idea is to consider the inclusion  $C_0(X) \rtimes_\lambda G \subseteq C(X^*) \rtimes_\lambda G$ , where  $X^* = X \sqcup \{\infty\}$ .




## Next step: reduced étale groupoid $C^*$ -algebras

Why? Many (or even most) separable  $C^*$ -algebras arise as the reduced  $C^*$ -algebra of a (twisted) étale groupoid.


# Next step: reduced étale groupoid $C^*$ -algebras

Why? Many (or even most) separable  $C^*$ -algebras arise as the reduced  $C^*$ -algebra of a (twisted) étale groupoid.



# Next step: reduced étale groupoid $C^*$ -algebras

Why? Many (or even most) separable  $C^*$ -algebras arise as the reduced  $C^*$ -algebra of a (twisted) étale groupoid.



For example, Xin Li showed that every separable simple nuclear  $\mathcal{Z}$ -stable UCT (i.e. classifiable within Elliott's program)  $C^*$ -algebra arises as the reduced  $C^*$ -algebra of a twisted étale Hausdorff groupoid.

# Étale groupoids

A **groupoid** is an algebraic structure  $(\mathcal{G}, ^{-1}, *)$  consisting of a set of objects  $\mathcal{G}$ , an inverse map  $^{-1} : G \rightarrow G$  and a (potentially only partially defined) multiplication  $* : G \times G \rightarrow G$ .

# Étale groupoids

A **groupoid** is an algebraic structure  $(\mathcal{G}, ^{-1}, *)$  consisting of a set of objects  $\mathcal{G}$ , an inverse map  $^{-1} : G \rightarrow G$  and a (potentially only partially defined) multiplication  $* : G \times G \rightarrow G$ .

The range and source maps  $r : \mathcal{G} \rightarrow \mathcal{G}$  and  $s : \mathcal{G} \rightarrow \mathcal{G}$  are defined by

$$r(g) = gg^{-1}, \quad s(g) = g^{-1}g.$$

For  $g, h \in \mathcal{G}$ , the product  $gh$  is defined if  $r(h) = s(g)$ .

# Étale groupoids

A **groupoid** is an algebraic structure  $(\mathcal{G}, ^{-1}, *)$  consisting of a set of objects  $\mathcal{G}$ , an inverse map  $^{-1} : \mathcal{G} \rightarrow \mathcal{G}$  and a (potentially only partially defined) multiplication  $*$  :  $\mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$ .

The range and source maps  $r : \mathcal{G} \rightarrow \mathcal{G}$  and  $s : \mathcal{G} \rightarrow \mathcal{G}$  are defined by

$$r(g) = gg^{-1}, \quad s(g) = g^{-1}g.$$

For  $g, h \in \mathcal{G}$ , the product  $gh$  is defined if  $r(h) = s(g)$ .

The unit space is  $\mathcal{G}^{(0)} = r(\mathcal{G})$ . Note that  $\mathcal{G} = \sqcup_{x \in \mathcal{G}^{(0)}} \mathcal{G}_x$ , where  $\mathcal{G}_x = \{g \in \mathcal{G} : s(g) = x\}$ .

# Étale groupoids

A **groupoid** is an algebraic structure  $(\mathcal{G}, ^{-1}, *)$  consisting of a set of objects  $\mathcal{G}$ , an inverse map  $^{-1} : \mathcal{G} \rightarrow \mathcal{G}$  and a (potentially only partially defined) multiplication  $*$  :  $\mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$ .

The range and source maps  $r : \mathcal{G} \rightarrow \mathcal{G}$  and  $s : \mathcal{G} \rightarrow \mathcal{G}$  are defined by

$$r(g) = gg^{-1}, \quad s(g) = g^{-1}g.$$

For  $g, h \in \mathcal{G}$ , the product  $gh$  is defined if  $r(h) = s(g)$ .

The unit space is  $\mathcal{G}^{(0)} = r(\mathcal{G})$ . Note that  $\mathcal{G} = \sqcup_{x \in \mathcal{G}^{(0)}} \mathcal{G}_x$ , where  $\mathcal{G}_x = \{g \in \mathcal{G} : s(g) = x\}$ .

The groupoid  $\mathcal{G}$  is **topological** if it is equipped with a locally compact topology for which the above maps are continuous and  $\mathcal{G}^{(0)}$  is Hausdorff in the relative topology. It is **étale** if, in addition,  $r$  is a local homeomorphism.

# Étale groupoids

A **groupoid** is an algebraic structure  $(\mathcal{G}, ^{-1}, *)$  consisting of a set of objects  $\mathcal{G}$ , an inverse map  $^{-1} : \mathcal{G} \rightarrow \mathcal{G}$  and a (potentially only partially defined) multiplication  $*$  :  $\mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$ .

The range and source maps  $r : \mathcal{G} \rightarrow \mathcal{G}$  and  $s : \mathcal{G} \rightarrow \mathcal{G}$  are defined by

$$r(g) = gg^{-1}, \quad s(g) = g^{-1}g.$$

For  $g, h \in \mathcal{G}$ , the product  $gh$  is defined if  $r(h) = s(g)$ .

The unit space is  $\mathcal{G}^{(0)} = r(\mathcal{G})$ . Note that  $\mathcal{G} = \sqcup_{x \in \mathcal{G}^{(0)}} \mathcal{G}_x$ , where  $\mathcal{G}_x = \{g \in \mathcal{G} : s(g) = x\}$ .

The groupoid  $\mathcal{G}$  is **topological** if it is equipped with a locally compact topology for which the above maps are continuous and  $\mathcal{G}^{(0)}$  is Hausdorff in the relative topology. It is **étale** if, in addition,  $r$  is a local homeomorphism.

**Intuition:** Think about homeomorphisms between open subsets of a topological space with composition as multiplication, defined whenever it makes sense.



# Étale groupoids

A **groupoid** is an algebraic structure  $(\mathcal{G}, ^{-1}, *)$  consisting of a set of objects  $G$ , an inverse map  $^{-1} : G \rightarrow G$  and a (potentially only partially defined) multiplication  $* : G \times G \rightarrow G$ .

The range and source maps  $r : \mathcal{G} \rightarrow G$  and  $s : \mathcal{G} \rightarrow G$  are defined by

$$r(g) = gg^{-1}, \quad s(g) = g^{-1}g.$$

For  $g, h \in \mathcal{G}$ , the product  $gh$  is defined if  $r(h) = s(g)$ .

The unit space is  $\mathcal{G}^{(0)} = r(\mathcal{G})$ . Note that  $\mathcal{G} = \sqcup_{x \in \mathcal{G}^{(0)}} \mathcal{G}_x$ , where  $\mathcal{G}_x = \{g \in \mathcal{G} : s(g) = x\}$ .

The groupoid  $\mathcal{G}$  is **topological** if it is equipped with a locally compact topology for which the above maps are continuous and  $\mathcal{G}^{(0)}$  is Hausdorff in the relative topology. It is **étale** if, in addition,  $r$  is a local homeomorphism.

**Intuition:** Think about homeomorphisms between open subsets of a topological space with composition as multiplication, defined whenever it makes sense.

Connes and Renault showed that **étale** groupoids give rise to an extremely rich class of  $C^*$ -algebras.

# Reduced $C^*$ -algebra of a (Hausdorff) étale groupoid

Let  $\mathcal{G}$  be a Hausdorff étale groupoid. For  $x \in \mathcal{G}^{(0)}$ , obtain a  $*$ -representation  $\lambda_x : C_c(\mathcal{G}) \rightarrow \mathcal{B}(\ell^2(\mathcal{G}_x))$  defined by

$$\lambda_x(f)\delta_g = \sum_{h \in \mathcal{G}_{r(g)}} f(h)\delta_{hg}.$$

# Reduced $C^*$ -algebra of a (Hausdorff) étale groupoid

Let  $\mathcal{G}$  be a Hausdorff étale groupoid. For  $x \in \mathcal{G}^{(0)}$ , obtain a  $*$ -representation  $\lambda_x : C_c(\mathcal{G}) \rightarrow \mathcal{B}(\ell^2(\mathcal{G}_x))$  defined by

$$\lambda_x(f)\delta_g = \sum_{h \in \mathcal{G}_{r(g)}} f(h)\delta_{hg}.$$

The regular  $*$ -representation  $\lambda : C_c(\mathcal{G}) \rightarrow \oplus_{x \in \mathcal{G}^{(0)}} \mathcal{B}(\ell^2(\mathcal{G}_x))$  of  $\mathcal{G}$  is

$$\lambda = \oplus_{x \in \mathcal{G}^{(0)}} \lambda_x.$$

# Reduced $C^*$ -algebra of a (Hausdorff) étale groupoid

Let  $\mathcal{G}$  be a Hausdorff étale groupoid. For  $x \in \mathcal{G}^{(0)}$ , obtain a  $*$ -representation  $\lambda_x : C_c(\mathcal{G}) \rightarrow \mathcal{B}(\ell^2(\mathcal{G}_x))$  defined by

$$\lambda_x(f)\delta_g = \sum_{h \in \mathcal{G}_{r(g)}} f(h)\delta_{hg}.$$

The regular  $*$ -representation  $\lambda : C_c(\mathcal{G}) \rightarrow \oplus_{x \in \mathcal{G}^{(0)}} \mathcal{B}(\ell^2(\mathcal{G}_x))$  of  $\mathcal{G}$  is

$$\lambda = \oplus_{x \in \mathcal{G}^{(0)}} \lambda_x.$$

The **reduced  $C^*$ -algebra**  $C_{\lambda}^*(\mathcal{G})$  of  $\mathcal{G}$  is the  $C^*$ -algebra generated by  $\lambda(C_c(\mathcal{G}))$ .

# Reduced (Hausdorff) étale groupoid $C^*$ -algebras

**Hausdorff case:**  $\mathcal{G}$  is a Hausdorff étale groupoid

# Reduced (Hausdorff) étale groupoid $C^*$ -algebras

**Hausdorff case:**  $\mathcal{G}$  is a Hausdorff étale groupoid

## Theorem (Brown-Clark-Farthing-Sims 2014)

*If  $\mathcal{G}$  is amenable, then  $C_\lambda^*(\mathcal{G})$  is simple if and only if  $\mathcal{G}$  is minimal and topologically principal.*

# Reduced (Hausdorff) étale groupoid $C^*$ -algebras

**Hausdorff case:**  $\mathcal{G}$  is a Hausdorff étale groupoid

## Theorem (Brown-Clark-Farthing-Sims 2014)

*If  $\mathcal{G}$  is amenable, then  $C_\lambda^*(\mathcal{G})$  is simple if and only if  $\mathcal{G}$  is minimal and topologically principal.*

## Theorem (Borys 2019)

*The  $C^*$ -algebra  $C_\lambda^*(\mathcal{G})$  is simple if  $\mathcal{G}$  has compact unit space, is minimal and has no amenable confined subgroups of isotropy groups.*

# Reduced (Hausdorff) étale groupoid $C^*$ -algebras

**Hausdorff case:**  $\mathcal{G}$  is a Hausdorff étale groupoid

## Theorem (Brown-Clark-Farthing-Sims 2014)

*If  $\mathcal{G}$  is amenable, then  $C_\lambda^*(\mathcal{G})$  is simple if and only if  $\mathcal{G}$  is minimal and topologically principal.*

## Theorem (Borys 2019)

*The  $C^*$ -algebra  $C_\lambda^*(\mathcal{G})$  is simple if  $\mathcal{G}$  has compact unit space, is minimal and has no amenable confined subgroups of isotropy groups.*

Note: One direction only. We will return to this point.



# Reduced (potentially non-Hausdorff) étale groupoid $C^*$ -algebras

**General case:**  $\mathcal{G}$  is a (potentially non-Hausdorff) étale groupoid

# Reduced (potentially non-Hausdorff) étale groupoid $C^*$ -algebras

**General case:**  $\mathcal{G}$  is a (potentially non-Hausdorff) étale groupoid

More appropriate to work with the **essential  $C^*$ -algebra**  $C_{\text{ess}}^*(\mathcal{G})$  introduced by Exel-Pitts and Kwaśniewski-Meyer.

# Reduced (potentially non-Hausdorff) étale groupoid $C^*$ -algebras

**General case:**  $\mathcal{G}$  is a (potentially non-Hausdorff) étale groupoid

More appropriate to work with the **essential  $C^*$ -algebra**  $C_{\text{ess}}^*(\mathcal{G})$  introduced by Exel-Pitts and Kwaśniewski-Meyer.

This is the quotient of  $C_\lambda^*(\mathcal{G})$  by ideal of “singular” elements

$$I = \{a \in C_\lambda^*(\mathcal{G}) : E(a^*a) = 0\},$$

where  $E : C_\lambda^*(\mathcal{G}) \rightarrow B^\infty(\mathcal{G}^{(0)})/M^\infty(\mathcal{G}^{(0)})$  is a conditional expectation onto the Dixmier algebra of bounded Borel functions modulo the functions with meager support.

# Reduced (potentially non-Hausdorff) étale groupoid $C^*$ -algebras

**General case:**  $\mathcal{G}$  is a (potentially non-Hausdorff) étale groupoid

More appropriate to work with the **essential  $C^*$ -algebra**  $C_{\text{ess}}^*(\mathcal{G})$  introduced by Exel-Pitts and Kwaśniewski-Meyer.

This is the quotient of  $C_\lambda^*(\mathcal{G})$  by ideal of “singular” elements

$$I = \{a \in C_\lambda^*(\mathcal{G}) : E(a^*a) = 0\},$$

where  $E : C_\lambda^*(\mathcal{G}) \rightarrow B^\infty(\mathcal{G}^{(0)})/M^\infty(\mathcal{G}^{(0)})$  is a conditional expectation onto the Dixmier algebra of bounded Borel functions modulo the functions with meager support.

Sufficient conditions for simplicity of  $C_{\text{ess}}^*(\mathcal{G})$  established in work of Clark-Exel-Pardo-Sims-Starling from 2019 and Kwaśniewski-Meyer from 2021.

# Preliminary strategy for characterizing simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

# Preliminary strategy for characterizing simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

**Preliminary strategy:** Motivated by the strategy for groups.

# Preliminary strategy for characterizing simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

**Preliminary strategy:** Motivated by the strategy for groups.

1. Construct Furstenberg boundary  $\partial_F \mathcal{G}$  of  $\mathcal{G}$ , i.e. the minimal injective object  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.

# Preliminary strategy for characterizing simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

**Preliminary strategy:** Motivated by the strategy for groups.

1. Construct Furstenberg boundary  $\partial_F \mathcal{G}$  of  $\mathcal{G}$ , i.e. the minimal injective object  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.
2. Replace  $\mathcal{G}$  by the “Furstenberg groupoid”  $\partial_F \mathcal{G} \rtimes \mathcal{G}$ , which is more tractable. Essentiality of the inclusion  $C_{\text{ess}}^*(\mathcal{G}) \subseteq C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  implies simplicity of smaller  $C^*$ -algebra equivalent to simplicity of larger  $C^*$ -algebra.



# Preliminary strategy for characterizing simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

**Preliminary strategy:** Motivated by the strategy for groups.

1. Construct Furstenberg boundary  $\partial_F \mathcal{G}$  of  $\mathcal{G}$ , i.e. the minimal injective object  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.
2. Replace  $\mathcal{G}$  by the “Furstenberg groupoid”  $\partial_F \mathcal{G} \rtimes \mathcal{G}$ , which is more tractable. Essentiality of the inclusion  $C_{\text{ess}}^*(\mathcal{G}) \subseteq C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  implies simplicity of smaller  $C^*$ -algebra equivalent to simplicity of larger  $C^*$ -algebra.
3. Classify  $\mathcal{G}$ -maps from  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  to  $C(\partial_F \mathcal{G})$ . Minimality of  $C(\partial_F \mathcal{G})$  implies correspondence between maps and quotients of  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$ .

# Preliminary strategy for characterizing simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

**Preliminary strategy:** Motivated by the strategy for groups.

1. Construct Furstenberg boundary  $\partial_F \mathcal{G}$  of  $\mathcal{G}$ , i.e. the minimal injective object  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.
2. Replace  $\mathcal{G}$  by the “Furstenberg groupoid”  $\partial_F \mathcal{G} \rtimes \mathcal{G}$ , which is more tractable. Essentiality of the inclusion  $C_{\text{ess}}^*(\mathcal{G}) \subseteq C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  implies simplicity of smaller  $C^*$ -algebra equivalent to simplicity of larger  $C^*$ -algebra.
3. Classify  $\mathcal{G}$ -maps from  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  to  $C(\partial_F \mathcal{G})$ . Minimality of  $C(\partial_F \mathcal{G})$  implies correspondence between maps and quotients of  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$ .

**Major obstruction:**  $C_{\text{ess}}^*(\mathcal{G})$  is typically not a  $\mathcal{G}$ - $C^*$ -algebra in the traditional sense, even when  $\mathcal{G}$  is Hausdorff.

# Preliminary strategy for characterizing simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

**Preliminary strategy:** Motivated by the strategy for groups.

1. Construct Furstenberg boundary  $\partial_F \mathcal{G}$  of  $\mathcal{G}$ , i.e. the minimal injective object  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.
2. Replace  $\mathcal{G}$  by the “Furstenberg groupoid”  $\partial_F \mathcal{G} \rtimes \mathcal{G}$ , which is more tractable. Essentiality of the inclusion  $C_{\text{ess}}^*(\mathcal{G}) \subseteq C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  implies simplicity of smaller  $C^*$ -algebra equivalent to simplicity of larger  $C^*$ -algebra.
3. Classify  $\mathcal{G}$ -maps from  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  to  $C(\partial_F \mathcal{G})$ . Minimality of  $C(\partial_F \mathcal{G})$  implies correspondence between maps and quotients of  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$ .

**Major obstruction:**  $C_{\text{ess}}^*(\mathcal{G})$  is typically not a  $\mathcal{G}$ - $C^*$ -algebra in the traditional sense, even when  $\mathcal{G}$  is Hausdorff.

Note: Traditionally, a  $\mathcal{G}$ - $C^*$ -algebra is fibered over  $\mathcal{G}^{(0)}$  and so contains a central copy of  $C(\mathcal{G}^{(0)})$ .

## New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the “pseudogroup”  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  on which the range and source maps restrict to homeomorphisms.

## New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the “pseudogroup”  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  on which the range and source maps restrict to homeomorphisms.

$$A \cdot B := \{g \cdot h : g \in a, h \in B, s(g) = r(h)\}$$

# New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the “pseudogroup”  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  on which the range and source maps restrict to homeomorphisms.

$$A \cdot B := \{g \cdot h : g \in a, h \in B, s(g) = r(h)\}$$

## Definition

A  $\mathcal{G}$ - $C^*$ -algebra  $A$  is a  $C^*$ -algebra containing a (not necessarily central) copy of  $C(\mathcal{G}^{(0)})$  along with compatible families of hereditary subalgebras  $(A_U)$  and  $*$ -isomorphisms  $(\alpha_\gamma)_{\gamma \in \Gamma(\mathcal{G})}$  satisfying

$$\alpha_\gamma : A_{\text{supp}(\gamma)} \rightarrow A_{\text{im}(\gamma)}.$$

# New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the “pseudogroup”  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  on which the range and source maps restrict to homeomorphisms.

$$A \cdot B := \{g \cdot h : g \in a, h \in B, s(g) = r(h)\}$$

## Definition

A  $\mathcal{G}$ - $C^*$ -algebra  $A$  is a  $C^*$ -algebra containing a (not necessarily central) copy of  $C(\mathcal{G}^{(0)})$  along with compatible families of hereditary subalgebras  $(A_U)$  and  $*$ -isomorphisms  $(\alpha_\gamma)_{\gamma \in \Gamma(\mathcal{G})}$  satisfying

$$\alpha_\gamma : A_{\text{supp}(\gamma)} \rightarrow A_{\text{im}(\gamma)}.$$

## Theorem (KKLRU 2021)

*There is a minimal injective  $\mathcal{G}$ - $C^*$ -algebra  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.*

## New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the pseudogroup  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  upon which the range and source maps restrict to homeomorphisms.



# New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the pseudogroup  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  upon which the range and source maps restrict to homeomorphisms.

## Definition (sketch)

A  $\mathcal{G}$ - $C^*$ -algebra  $A$  is a  $C^*$ -algebra containing a copy of  $C(\mathcal{G}^{(0)})$  along with compatible families of hereditary sub- $C^*$ -algebras  $(A_U)$  and  $*$ -isomorphisms  $(\alpha_\gamma)_{\gamma \in \Gamma(\mathcal{G})}$  satisfying

$$\alpha_\gamma : A_{\text{supp}(\gamma)} \rightarrow A_{\text{im}(\gamma)}.$$

# New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the pseudogroup  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  upon which the range and source maps restrict to homeomorphisms.

## Definition (sketch)

A  $\mathcal{G}$ - $C^*$ -algebra  $A$  is a  $C^*$ -algebra containing a copy of  $C(\mathcal{G}^{(0)})$  along with compatible families of hereditary sub- $C^*$ -algebras  $(A_U)$  and  $*$ -isomorphisms  $(\alpha_\gamma)_{\gamma \in \Gamma(\mathcal{G})}$  satisfying

$$\alpha_\gamma : A_{\text{supp}(\gamma)} \rightarrow A_{\text{im}(\gamma)}.$$

## Theorem (KKLRU 2021)

*There is a minimal injective  $\mathcal{G}$ - $C^*$ -algebra  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.*

# New category of groupoid $C^*$ -algebras

**Solution:** Instead of acting by elements of  $\mathcal{G}$ , act by elements of the pseudogroup  $\Gamma(\mathcal{G})$  consisting of open bisections of  $\mathcal{G}$ , i.e. open subsets of  $\mathcal{G}$  upon which the range and source maps restrict to homeomorphisms.

## Definition (sketch)

A  $\mathcal{G}$ - $C^*$ -algebra  $A$  is a  $C^*$ -algebra containing a copy of  $C(\mathcal{G}^{(0)})$  along with compatible families of hereditary sub- $C^*$ -algebras  $(A_U)$  and  $*$ -isomorphisms  $(\alpha_\gamma)_{\gamma \in \Gamma(\mathcal{G})}$  satisfying

$$\alpha_\gamma : A_{\text{supp}(\gamma)} \rightarrow A_{\text{im}(\gamma)}.$$

## Theorem (KKLRU 2021)

*There is a minimal injective  $\mathcal{G}$ - $C^*$ -algebra  $C(\partial_F \mathcal{G})$  in the category of  $\mathcal{G}$ - $C^*$ -algebras.*

We refer to  $\partial_F \mathcal{G}$  as the **Furstenberg boundary** of  $\mathcal{G}$ . For Hausdorff  $\mathcal{G}$ , coincides with Furstenberg boundary constructed by Borys.

# Ideal intersection property

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

# Ideal intersection property

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

More generally, we can ask when the ideals in  $C_{\text{ess}}^*(\mathcal{G})$  are completely determined by the dynamics of  $\mathcal{G}$ . The following property was first considered by Kawamura-Tomiyama (1990) and Sierakowski (2010) for crossed products.

# Ideal intersection property

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

More generally, we can ask when the ideals in  $C_{\text{ess}}^*(\mathcal{G})$  are completely determined by the dynamics of  $\mathcal{G}$ . The following property was first considered by Kawamura-Tomiyama (1990) and Sierakowski (2010) for crossed products.

## Definition

The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  has the **ideal intersection property** if every nonzero ideal of  $C_{\text{ess}}^*(\mathcal{G})$  has nonzero intersection with  $C_0(\mathcal{G}^{(0)})$ .

# Ideal intersection property

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

More generally, we can ask when the ideals in  $C_{\text{ess}}^*(\mathcal{G})$  are completely determined by the dynamics of  $\mathcal{G}$ . The following property was first considered by Kawamura-Tomiyama (1990) and Sierakowski (2010) for crossed products.

## Definition

The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  has the **ideal intersection property** if every nonzero ideal of  $C_{\text{ess}}^*(\mathcal{G})$  has nonzero intersection with  $C_0(\mathcal{G}^{(0)})$ .

Invariant subspaces of  $\mathcal{G}^{(0)}$  clearly give rise to ideals of  $C_{\text{ess}}^*(\mathcal{G})$ . If  $\mathcal{G}$  is Hausdorff and inner exact, then the ideal intersection property is equivalent to the statement that every ideal in  $C_{\lambda}^*(\mathcal{G})$  arises in this way. Note: inner exactness is necessary.

# Dynamical characterization of simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .



# Dynamical characterization of simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

## Theorem (KKLRU 2021)

*The following are equivalent:*

1. *The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  has the ideal intersection property.*
2. *The  $C^*$ -algebra  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  has the ideal intersection property.*
3. *The Furstenberg boundary  $\partial_F \mathcal{G}$  is free.*
4. *There is a unique  $\mathcal{G}$ -pseudoexpectation from  $C_{\text{ess}}^*(\mathcal{G})$  to  $C(\partial_F \mathcal{G})$ .*

# Dynamical characterization of simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

## Theorem (KKLRU 2021)

*The following are equivalent:*

1. *The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  has the ideal intersection property.*
2. *The  $C^*$ -algebra  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  has the ideal intersection property.*
3. *The Furstenberg boundary  $\partial_F \mathcal{G}$  is free.*
4. *There is a unique  $\mathcal{G}$ -pseudoexpectation from  $C_{\text{ess}}^*(\mathcal{G})$  to  $C(\partial_F \mathcal{G})$ .*

## Corollary

*The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  is simple if and only if  $\partial_F \mathcal{G}$  is minimal and free.*

# Dynamical characterization of simplicity

Let  $\mathcal{G}$  be an étale groupoid with compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

## Theorem (KKLRU 2021)

*The following are equivalent:*

1. *The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  has the ideal intersection property.*
2. *The  $C^*$ -algebra  $C_{\text{ess}}^*(\partial_F \mathcal{G} \rtimes \mathcal{G})$  has the ideal intersection property.*
3. *The Furstenberg boundary  $\partial_F \mathcal{G}$  is free.*
4. *There is a unique  $\mathcal{G}$ -pseudoexpectation from  $C_{\text{ess}}^*(\mathcal{G})$  to  $C(\partial_F \mathcal{G})$ .*

## Corollary

*The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  is simple if and only if  $\partial_F \mathcal{G}$  is minimal and free.*

Note: We can apply this result when  $\mathcal{G}^{(0)}$  is not compact by replacing  $\mathcal{G}$  with a suitable one-point compactification.

# Intrinsic characterization of simplicity

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

# Intrinsic characterization of simplicity

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

For  $x \in \mathcal{G}^{(0)}$ , the **isotropy group**  $\mathcal{G}_x^\times$  is

$$\mathcal{G}_x^\times = \{g \in \mathcal{G} : s(g) = r(g) = x\}.$$

An **isotropy subgroup** is a subgroup of some  $\mathcal{G}_x^\times$ .

# Intrinsic characterization of simplicity

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

For  $x \in \mathcal{G}^{(0)}$ , the **isotropy group**  $\mathcal{G}_x^\times$  is

$$\mathcal{G}_x^\times = \{g \in \mathcal{G} : s(g) = r(g) = x\}.$$

An **isotropy subgroup** is a subgroup of some  $\mathcal{G}_x^\times$ .

## Theorem (KKLRU 2021)

*The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  is simple if and only if  $\mathcal{G}$  is minimal and has no amenable confined isotropy subgroups.*

# **Intrinsic characterization of the ideal intersection property**

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

# Intrinsic characterization of the ideal intersection property

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

A **section** of isotropy subgroups a collection

$$\Lambda = \{(x, H_x) : x \in \mathcal{G}^{(0)}, H_x \leq \mathcal{G}_x^x\}.$$

The section  $\Lambda$  is **essentially confined** if  $\mathcal{G}^{(0)} \not\subseteq \overline{\mathcal{G}\Lambda}$ .



# Intrinsic characterization of the ideal intersection property

Let  $\mathcal{G}$  be an étale groupoid with locally compact Hausdorff unit space  $\mathcal{G}^{(0)}$ .

A **section** of isotropy subgroups is a collection

$$\Lambda = \{(x, H_x) : x \in \mathcal{G}^{(0)}, H_x \leq \mathcal{G}_x^x\}.$$

The section  $\Lambda$  is **essentially confined** if  $\mathcal{G}^{(0)} \not\subseteq \overline{\mathcal{G}\Lambda}$ .

## Theorem (KKLRU 2021)

*The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  has the ideal intersection property if and only if  $\mathcal{G}$  has no amenable essentially confined sections of isotropy subgroups.*

# Relative Powers Averaging Property

We generalize K-Haagerup and Amrutan-Ursu's characterization of simplicity in terms of Powers-type averaging properties.

# Relative Powers Averaging Property

We generalize K-Haagerup and Amrutan-Ursu's characterization of simplicity in terms of Powers-type averaging properties.

## Theorem (KKLRU 2021)

*Let  $\mathcal{G}$  be a minimal étale groupoid with compact Hausdorff unit space. The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  is simple if and only if for every  $a \in C_{\text{ess}}^*(\mathcal{G})$ ,*

$$0 \in \overline{\{\mu * a : \mu \text{ a "generalized" probability measure}\}}$$

# Relative Powers Averaging Property

We generalize K-Haagerup and Amrutan-Ursu's characterization of simplicity in terms of Powers-type averaging properties.

## Theorem (KKLRU 2021)

*Let  $\mathcal{G}$  be a minimal étale groupoid with compact Hausdorff unit space. The  $C^*$ -algebra  $C_{\text{ess}}^*(\mathcal{G})$  is simple if and only if for every  $a \in C_{\text{ess}}^*(\mathcal{G})$ ,*

$$0 \in \overline{\{\mu * a : \mu \text{ a "generalized" probability measure}\}}$$

Note: The terminology “generalized” probability measure is used in the sense of Amrutan-Ursu.

# Related open problems

1. In the non-Hausdorff case, what is the relationship between the ideal structure of  $C_{ess}^*(\mathcal{G})$  and  $C_\lambda^*(G)$ ?

# Related open problems

1. In the non-Hausdorff case, what is the relationship between the ideal structure of  $C_{ess}^*(\mathcal{G})$  and  $C_\lambda^*(G)$ ?
2. How does the triviality of the Furstenberg boundary  $\partial_F \mathcal{G}$  relate to the amenability of  $\mathcal{G}$ ?

# Related open problems

1. In the non-Hausdorff case, what is the relationship between the ideal structure of  $C_{ess}^*(\mathcal{G})$  and  $C_\lambda^*(G)$ ?
2. How does the triviality of the Furstenberg boundary  $\partial_F \mathcal{G}$  relate to the amenability of  $\mathcal{G}$ ?
3. When is the reduced  $C^*$ -algebra of a twisted group simple?

Thanks!