

Estimating winter balance and its uncertainty from direct measurements of snow depth and density on alpine glaciers

Alexandra PULWICKI,¹ Gwenn E. FLOWERS,¹ Valentina RADIC,²

¹ *Department of Earth Sciences, Faculty of Science, Simon Fraser University, Burnaby, BC, Canada*

² *Department of Earth, Ocean and Atmospheric Sciences, Faculty of Science, University of British Columbia, Vancouver, BC, Canada*

Correspondence: Alexandra Pulwicksi <apulwick@sfu.ca>

ABSTRACT. Accurately estimating winter surface mass balance on glaciers is central to assessing glacier health and predicting glacier runoff. However, measuring and modelling snow distribution is inherently difficult in mountainous terrain. Here we explore rigorous statistical methods of estimating winter balance and its uncertainty from multiscale measurements of snow depth and density. In May 2016 we collected over 9000 manual measurements of snow depth across three glaciers in the St. Elias Mountains, Yukon, Canada. Linear regression, combined with cross correlation and Bayesian model averaging, as well as simple kriging are used to interpolate point-scale values to glacier-wide estimates of winter balance. Elevation and a wind-redistribution parameter exhibit the highest correlations with winter balance, but the relationship varies considerably between glaciers. A Monte Carlo analysis reveals that the interpolation itself introduces more uncertainty than the assignment of snow density or the representation of grid-scale variability. For our study glaciers, the winter balance uncertainty from all assessed sources ranges from 0.03 m w.e. (8%) to 0.15 m w.e. (54%). Despite the challenges associated with estimating winter balance, our results are consistent with a regional-scale winter-balance gradient.

INTRODUCTION

Winter surface mass balance, or “winter balance”, is the net accumulation and ablation of snow over the winter season (Cogley and others, 2011), which constitutes glacier mass input. Winter balance is half of the seasonally resolved mass balance, initializes summer ablation conditions and must be estimated to simulate energy and mass exchange between the land and atmosphere (e.g. Hock, 2005; Réveillet and others, 2016). Effectively representing the spatial distribution of snow is also central to monitoring surface runoff and its downstream effects (e.g. Clark and others, 2011).

Winter balance (WB) is notoriously difficult to estimate (e.g. Dadić and others, 2010; Cogley and others, 2011). Snow distribution in alpine regions is highly variable with short correlation length scales (e.g. Anderton and others, 2004; Egli and others, 2011; Grünewald and others, 2010; Helbig and van Herwijnen, 2017; López-Moreno and others, 2011, 2013; Machguth and others, 2006; Marshall and others, 2006) and is influenced by dynamic interactions between the atmosphere and complex topography, operating on multiple spatial and temporal scales (e.g. Barry, 1992; Liston and Elder, 2006; Clark and others, 2011). Simultaneously extensive, high resolution and accurate snow distribution measurements on glaciers are therefore difficult to obtain (e.g. Cogley and others, 2011; McGrath and others, 2015) and is further complicated by the inaccessibility of many glacierized regions during the winter. Physically based models are computationally intensive and require detailed meteorological data to drive them (Dadić and others, 2010). As a result, there is significant uncertainty in estimates of winter balance, thus limiting the ability of models to represent current and projected glacier conditions.

Studies that have focused on obtaining detailed estimates of WB have used a wide range of observational techniques, including direct measurement of snow depth and density (e.g. Cullen and others, 2017), lidar or photogrammetry (e.g. Sold and others, 2013) and ground-penetrating radar (e.g. Machguth and others, 2006; Gusmeroli and others, 2014; McGrath and others, 2015). Spatial coverage of direct measurements is generally limited and comprises an elevation transect along the glacier centreline (e.g. Kaser and others, 2003). Measurements are often interpolated using linear regression on only a few topographic parameters (e.g. MacDougall and Flowers, 2011), with elevation being the most common. Other established techniques include hand contouring (e.g. Tangborn and others, 1975), kriging (e.g. Hock and Jensen, 1999) and attributing measured winter balance values to elevation bands (e.g. Thibert and others, 2008). Physical snow models have been used to estimate spatial patterns of winter balance (e.g. Mott and others, 2008; Schuler and others, 2008; Dadić and others, 2010) but availability of the required meteorological data generally prohibits their

widespread application. Error analysis is rarely undertaken and few studies have thoroughly investigated uncertainty in spatially distributed estimates of winter balance (c.f. Schuler and others, 2008).

More sophisticated snow-survey designs and statistical models of snow distribution are widely used in the field of snow science. Surveys described in the snow science literature are generally spatially extensive and designed to measure snow depth and density throughout a basin, ensuring that all terrain types are sampled. A wide array of measurement interpolation methods are used, including linear (e.g. López-Moreno and others, 2010) and non-linear regressions (e.g. Molotch and others, 2005) that include numerous terrain parameters, as well as geospatial interpolation (e.g. Erxleben and others, 2002; Cullen and others, 2017) including various forms of kriging. Different interpolation methods are also combined; for example, regression kriging adds kriged residuals to a field obtained with linear regression (e.g. Balk and Elder, 2000). Physical snow models such as SnowTran-3D (Liston and Sturm, 1998), Alpine3D (Lehning and others, 2006), and SnowDrift3D (Schneiderbauer and Prokop, 2011) are widely used, and errors in estimating snow distribution have been examined from theoretical (e.g. Trujillo and Lehning, 2015) and applied perspectives (e.g. Turcan and Loijens, 1975; Woo and Marsh, 1978; Deems and Painter, 2006).

The goals of this study are to (1) critically examine methods of converting direct snow depth and density measurements to distributed estimates of winter balance and (2) identify sources of uncertainty, evaluate their magnitude and assess their combined contribution to uncertainty in glacier-wide winter balance. We focus on commonly applied, low-complexity methods of measuring and estimating winter balance in the interest of making our results broadly applicable.

STUDY SITE

The St. Elias Mountains (Fig. 1a) rise sharply from the Pacific Ocean, creating a significant climatic gradient between coastal maritime conditions, generated by Aleutian–Gulf of Alaska low-pressure systems, and interior continental conditions, driven by the Yukon–Mackenzie high-pressure system (Taylor-Barge, 1969). The boundary between the two climatic zones is generally aligned with the divide between the Hubbard and Kaskawulsh Glaciers, approximately 130 km from the coast. Research on snow distribution and glacier mass balance in this area is limited. A series of research programs, including Project “Snow Cornice” and the Icefield Ranges Research Project, were operational in the 1950s and 60s (Wood, 1948; Danby and others, 2003) and in the last 30 years, there have been a few long-term studies on selected alpine glaciers (e.g. Clarke, 2014) as well as several regional studies of glacier mass balance and dynamics (e.g. Arendt and others, 2008; Burgess and others, 2013; Waechter and others, 2015).




Fig. 1. Study area location and sampling design for Glaciers 4, 2 and 13. (a) Study region in the Donjek Range of the St. Elias Mountains of Yukon, Canada. (b) Study glaciers located along a southwest-northeast transect through the Donjek Range. The local topographic divide is shown as a dashed line. Imagery from Landsat8 (5 September 2013, data available from the U.S. Geological Survey). (c) Details of the snow-survey sampling design, with centreline and transverse transects (blue dots), hourglass and circle designs (green dots) and locations of snow density measurements (orange squares). Arrows indicate ice-flow directions. Approximate location of ELA on each glacier is shown as a black dashed line. (d) Close up of linear and curvilinear transects. (e) Configuration of navigator and observers. (f) Point-scale snow-depth sampling. (g) Linear-random snow-depth measurements in ‘zigzag’ design (purple dots) with one density measurement (orange square) per zigzag.

86 We carried out winter balance surveys on three unnamed glaciers in the Donjek Range of the St. Elias
 87 Mountains. The Donjek Range is located approximately 40 km to the east of the regional mountain divide
 88 and has an area of about $30 \times 30 \text{ km}^2$. Glacier 4, Glacier 2 and Glacier 13 (labelling adopted from Crompton
 89 and Flowers (2016)) are located along a southwest-northeast transect through the range (Fig. 1b, Table 1).
 90 These small alpine glaciers are generally oriented southeast-northwest, with Glacier 4 having a predominantly
 91 southeast aspect and Glaciers 2 and 13 have generally northwest aspects. The glaciers are situated in valleys
 92 with steep walls and have simple geometries. Based on a detailed study of Glacier 2 (Wilson and others,

Table 1. Physical characteristics of the study glaciers.

	Location		Elevation (m a.s.l.)			Slope (°)	Area
	UTM Zone 7		<i>Mean</i>	<i>Range</i>	<i>ELA</i>	<i>Mean</i>	(km ²)
Glacier 4	595470 E	6740730 N	2344	1958–2809	~2500	12.8	3.8
Glacier 2	601160 E	6753785 N	2495	1899–3103	~2500	13.0	7.0
Glacier 13	604602 E	6763400 N	2428	1923–3067	~2380	13.4	12.6

2013) and related theoretical modelling (Wilson and Flowers, 2013) we suspect all of the study glaciers to be polythermal.

METHODS

Estimating glacier-wide winter balance involves transforming measurements of snow depth and density into values of winter balance distributed across a defined grid. We do this in four steps. (1) Obtain direct measurements of snow depth and density in the field. (2) Assign density values to all depth-measurement locations to calculate point-scale values of WB at each location. Winter balance can be estimated as the product of snow depth and depth-averaged density. (3) Average all point-scale values within each gridcell of a digital elevation model (DEM) to obtain the gridcell-averaged WB. (4) Interpolate and extrapolate these gridcell-averaged WB values to obtain estimates of WB (in m w.e.) in each gridcell across the domain. Glacier-wide WB is then calculated by taking the average of all gridcell-averaged WB values for each glacier. For brevity, we refer to these four steps as (1) field measurements, (2) density assignment, (3) gridcell-averaged

Table 2. Details of the May 2016 winter-balance survey, including number of snow-depth measurement locations along transects (n_T), total length of transects (d_T), number of combined snow pit and Federal Sampler density measurement locations (n_ρ), number of zigzag surveys (n_{zz}), number (and as percent of total number of gridcells) of gridcells sampled (n_S) and the elevation range (and as percent of total elevations range).

	Date	n_T	d_T (km)	n_ρ	n_{zz}	n_S	Elevation range (m a.s.l.)
Glacier 4	4–7 May 2016	649	13.1	7	3	295 (12%, 21%)	2015–2539 (62%, 97%)
Glacier 2	8–11 May 2016	762	13.6	7	3	353 (8%, 14%)	2151–2541 (32%, 47%)
Glacier 13	12–15 May 2016	941	18.1	19	4	468 (6%, 14%)	2054–2574 (45%, 62%)

WB and (4) distributed WB. Detailed methodology for each step is outlined below. We use the SPIRIT
 SPOT-5 DEM (40×40 m) from 2005 (Korona and others, 2009) throughout this study.

Field measurements

Our sampling campaign involved four people and occurred between 5–15 May 2016, which falls within the
 period of historical peak snow accumulation in southwestern Yukon (Yukon Snow Survey Bulletin and Water
 Supply Forecast, May 1, 2016). Snow depth is generally accepted to be more variable than density (Elder
 and others, 1991; Clark and others, 2011; López-Moreno and others, 2013) so we chose a sampling design
 that resulted in a high ratio ($\sim 55:1$) of snow depth to density measurements. In total, we collected more
 than 9000 snow-depth measurements and more than 100 density measurements throughout the study area
 (Table 1).

During the field campaign there were two small accumulation events. The first, on 6 May 2016, also involved
 high winds so accumulation could not be determined. The second, on 10 May 2016, resulted in 0.01 m w.e
 accumulation measured at one location on Glacier 2. Assuming both accumulation events contributed a
 uniform 0.01 m w.e accumulation to all study glaciers then our survey did not capture $\sim 3\%$ and $\sim 2\%$ of
 estimated glacier-wide winter balance on Glaciers 4 and 2, respectively. We therefore assume that these
 accumulation events were negligible. Positive temperatures and clear skies occurred between 11–16 May
 2016, which we suspect resulted in melt occurring on Glacier 13. The snow in the lower part of the ablation
 area of Glacier 13 was isothermal and showed clear signs of melt and metamorphosis. The total amount of
 melt during the study period was not estimated so no corrections were made.

Sampling design

The snow surveys were designed to capture variability in snow depth at regional, basin, gridcell and point
 scales (Clark and others, 2011). To capture variability at the regional scale we chose three glaciers along
 a transect aligned with the dominant precipitation gradient (Fig. 1) (Taylor-Barge, 1969). To account for
 basin-scale variability, snow depth was measured along linear and curvilinear transects on each glacier (Fig.
 1c) with a sample spacing of 10–60 m (Fig. 1d). Sample spacing was constrained by protocols for safe glacier
 travel, while survey scope was constrained by the need to complete all surveys within the period of peak
 accumulation. We selected centreline and transverse transects as the most commonly used survey designs
 in winter balance studies (e.g. Kaser and others, 2003; Machguth and others, 2006) as well as an hourglass
 pattern with an inscribed circle, which allows for sampling in multiple directions and easy travel (personal
 communication from C. Parr, 2016). To capture variability at the grid scale, we densely sampled up to four

135 gridcells on each glacier using a linear-random sampling design (Shea and Jamieson, 2010) we term a ‘zigzag’.
136 To capture point-scale variability, each observer made 3–4 depth measurements within ~ 1 m (Fig. 1f) at
137 each transect measurement location.

138 *Snow depth: transects*

139 While roped-up for glacier travel with fixed distances between observers, the lead observer used a single-
140 frequency GPS unit (Garmin GPSMAP 64s) to navigate between predefined transect measurement locations
141 (Fig. 1e). The remaining three observers used 3.2 m graduated aluminum avalanche probes to make snow-
142 depth measurements. The locations of each set of depth measurements, made by the second, third and fourth
143 observers, are estimated using the recorded location of the first observer, the approximate distance between
144 observers and the direction of travel. The 3–4 point-scale depth measurements are averaged to obtain a single
145 depth measurement at each transect measurement location. When considering snow variability at the point
146 scale as a source of uncertainty in snow depth measurements, we find that the mean standard deviation of
147 point-scale snow depth measurements is found to be $< 7\%$ of the mean snow depth for all study glaciers.

148 Snow-depth sampling was concentrated in the ablation area to ensure that only snow from the current
149 accumulation season was measured. The boundary between snow and firn in the accumulation area can be
150 difficult to detect and often misinterpreted, especially when using an avalanche probe (Grünwald and others,
151 2010; Sold and others, 2013). We intended to use a firn corer to measure winter balance in the accumulation
152 area, but cold snow combined with positive air temperatures led to cores being unrecoverable. Successful
153 snow depth measurements within the accumulation area were made either in snow pits or using a Federal
154 Sampler (described below) to unambiguously identify the snow–firn transition.

155 *Snow depth: zigzags*

156 We measured depth at random intervals of 0.3–3.0 m along two ‘Z’-shaped patterns (Shea and Jamieson,
157 2010), resulting in 135–191 measurements per zigzag, within three to four 40×40 m gridcells (Fig. 1g) per
158 glacier. Random intervals were machine-generated from a uniform distribution in sufficient numbers that
159 each survey was unique. Zigzag locations were randomly chosen within the upper, middle and lower regions
160 of the ablation area of each glacier. Extra time in the field allowed us to measure a fourth zigzag on Glacier
161 13 in the central ablation area at ~ 2200 m a.s.l.

162 *Snow density*

163 Snow density was measured using a Snowmetrics wedge cutter in three snow pits on each glacier. Within
 164 the snow pits (SP), we measured a vertical density profile (in 5 cm increments) with the $5 \times 10 \times 10$ cm
 165 wedge-shaped cutter (250 cm^3) and a Presola 1000 g spring scale (e.g. Gray and Male, 1981; Fierz and others,
 166 2009). Wedge-cutter error is approximately $\pm 6\%$ (e.g. Proksch and others, 2016; Carroll, 1977). Uncertainty
 167 in estimating density from SP measurements also stems from incorrect assignment of density to layers that
 168 cannot be sampled (e.g. ice lenses and hard layers). We attempt to quantify this uncertainty by varying
 169 estimated ice-layer thickness by ± 1 cm ($\leq 100\%$) of the recorded thickness, ice layer density between 700
 170 and 900 kg m^{-3} and the density of layers identified as being too hard to sample (but not ice) between 600
 171 and 700 kg m^{-3} . When considering all three sources of uncertainty, the range of integrated density values
 172 is always less than 15% of the reference density. Depth-averaged densities for shallow pits (< 50 cm) that
 173 contain ice lenses are particularly sensitive to changes in prescribed density and ice-lens thickness.

174 While SP provide the most accurate measure of snow density, digging and sampling a SP is time and
 175 labour intensive. Therefore, a Geo Scientific Ltd. metric Federal Sampler (FS) (Clyde, 1932) with a 3.2–
 176 3.8 cm diameter sampling tube, which directly measures depth-integrated snow-water equivalent, was used
 177 to augment the snow pit measurements. A minimum of three FS measurements were taken at each of 7–19
 178 locations on each glacier and an additional eight FS measurements were co-located with each snow pit profile.
 179 Measurements for which the snow core length inside the sampling tube was less than 90% of the snow depth
 180 were discarded. Densities at each measurement location (eight at each snow pit, three elsewhere) were then
 181 averaged, with the standard deviation taken to represent the uncertainty. The mean standard deviation of
 182 FS-derived density was $\leq 4\%$ of the mean density for all glaciers.

183 **Density assignment**

184 Measured snow density must be interpolated or extrapolated to estimate point-scale winter balance at each
 185 snow-depth sampling location. We employ four commonly used methods to interpolate and extrapolate
 186 density (Table 3): (1) calculate mean density over an entire mountain range (e.g. Cullen and others, 2017),
 187 (2) calculate mean density for each glacier (e.g. Elder and others, 1991; McGrath and others, 2015), (3)
 188 linear regression of density on elevation for each glacier (e.g. Elder and others, 1998; Molotch and others,
 189 2005) and (4) calculate mean density using inverse-distance weighting (e.g. Molotch and others, 2005) for
 190 each glacier. Densities derived from SP and FS measurements are treated separately, for reasons explained
 191 below, resulting in eight possible methods of assigning density.

Table 3. Eight methods used to estimate snow density at unmeasured locations. Total number of resulting density values given in parentheses, with n_T the total number of snow-depth measurement locations along transects (Table 1).

Method code	Source of measured snow density		Density assignment method
	<i>Snow pit</i>	<i>Federal</i>	
		<i>Sampler</i>	
S1	■		Mean of measurements
F1		■	across all glaciers (1)
S2	■		Mean of measurements
F2		■	for each glacier (3)
S3	■		Regression of density on
F3		■	elevation for a glacier (n_T)
S4	■		Inverse distance weighted
F4		■	mean for a glacier (n_T)

Gridcell-averaged winter balance

We average one to six (mean of 2.1 measurements) point-scale values of WB within each 40×40 m DEM gridcell to obtain the gridcell-averaged WB. The locations of individual measurements have uncertainty due to the error in the horizontal position given by the GPS unit and the estimation of observer location based on the recorded GPS positions of the navigator. This location uncertainty could result in the incorrect assignment of a point-scale WB to a particular gridcell. However, this source of error is not further investigated because we assume that the uncertainty in gridcell-averaged WB is captured in the zigzag measurements described below. Error due to having multiple observers is also evaluated by conducting an analysis of variance (ANOVA) of snow-depth measurement along a transect and testing for differences between observers. We find no significant differences between snow-depth measurements made by observers along any transect ($p > 0.05$), with the exception of the first transect on Glacier 4 (51 measurements), where snow depth values collected by one observer were, on average, greater than the snow depth measurements taken by the other two observers ($p < 0.01$). Since this was the first transect completed and the only one to show differences by observer, this difference can be considered an anomaly. This result shows that observer bias is likely to not affect the results of this study and no corrections to the data based on observer were applied.

Distributed winter balance

Gridcell-averaged values of WB are interpolated and extrapolated across each glacier using linear regression (LR) and simple kriging (SK). The LR relates gridcell-averaged WB and various topographic parameters, as this method is simple and has precedent for success (e.g. McGrath and others, 2015). Instead of a basic LR however, we use cross-validation and model averaging to test all combinations of the topographic parameters. We compare the LR approach with SK, a data-driven interpolation method free of any physical interpretation (e.g. Hock and Jensen, 1999).

Linear regression

In the LR, we use commonly applied topographic parameters as in McGrath and others (2015), including elevation, slope, aspect, curvature, “northness” and a wind-redistribution parameter (Sx from Winstral and others (2002)); we add distance-from-centreline as an additional parameter. Topographic parameters are standardized for use in the LR. For details on data and methods used to estimate the topographic parameters see the Supplementary Material. Our sampling design ensured that the ranges of topographic parameters associated with our measurement locations represent more than 70% of the total area of each glacier (except elevation on Glacier 2, where our measurements captured only 50%).

The goal of the LR is to obtain a set of fitted regression coefficients (β_i) that correspond to each topographic parameter and to a model intercept. The LR implemented in this study is an extension of a basic multiple linear regression; we use cross-validation to avoid overfitting the data and model averaging to incorporate every possible combination of topographic parameters.

First, cross-validation is used to obtain a set of regression coefficients that have the greatest predictive ability (Kohavi and others, 1995). We randomly select 1000 subsets of the data (2/3 of the values) and fit a basic multiple linear regression (implemented in MATLAB) to the data subsets, thus obtaining 1000 sets of regression coefficients. The basic multiple linear regression calculates a set of regression coefficients by minimizing the sum of squares of the vertical deviations of each datum from the regression line (Davis and Sampson, 1986). Distributed winter balance is then calculated by weighting each topographic parameter by the corresponding regression coefficient for all DEM gridcells. We then use the remaining data (1/3 of the values) to calculate a root mean squared error (RMSE) between the estimated WB and the observed WB for corresponding locations. From the 1000 sets of β_i values, we select the set that results in the lowest RMSE.

Second, we use model averaging to account for uncertainty when selecting predictors and to maximize the model’s predictive ability (Madigan and Raftery, 1994). Models are generated by calculating a set of

237 regression coefficients as described above for all possible combinations of topographic parameters (resulting in
 238 2^7 models, or sets of regression coefficients). Using a Bayesian framework, model averaging involves weighting
 239 all models by their posterior model probabilities (Raftery and others, 1997). The models (i.e. sets of regression
 240 coefficients) are weighted according to the relative predictive success of the model, as assessed by the value
 241 of the Bayesian Information Criterion (BIC) (Burnham and Anderson, 2004). BIC penalizes more complex
 242 models which further reduces the risk of overfitting. The final set of regression coefficients is then the weighted
 243 sum of regression coefficients from all models. Distributed WB is obtained by applying the final set of
 244 regression coefficients to the topographic parameters associated with each gridcell.

245 *Simple kriging*

246 SK is a data-driven method of estimating variables at unsampled locations by using the isotropic spatial
 247 correlation (covariance) of measured values to find a set of optimal weights (Davis and Sampson, 1986; Li
 248 and Heap, 2008). SK assumes spatial correlation between sampling locations that are distributed across a
 249 surface and then applies the correlation to interpolate between these locations. We used the **DiceKriging**
 250 R package (Roustant and others, 2012) to calculate the maximum likelihood covariance matrix, as well as
 251 the range distance (θ) and nugget for gridcell-averaged values of winter balance. The range distance is a
 252 measure of data correlation length and the nugget is the residual that encompasses sampling-error variance
 253 as well as the spatial variance at distances less than the minimum sample spacing (Li and Heap, 2008). A
 254 Matérn covariance function with $\nu=5/2$ is used to define a stationary and isotropic covariance and covariance
 255 kernels are parameterized as in Rasmussen and Williams (2006). Unlike LR, SK is not useful for generating
 256 hypotheses to explain the physical controls on snow distribution, nor can it be used to estimate winter
 257 balance on unmeasured glaciers.

258 **Uncertainty analysis using a Monte Carlo approach**

259 Three sources of uncertainty are considered separately: the uncertainty due to (1) grid-scale variability of
 260 WB (σ_{GS}), (2) the assignment of snow density (σ_{ρ}) and (3) interpolating and extrapolating gridcell-averaged
 261 values of WB (σ_{INT}). To quantify the uncertainty of grid-scale and interpolation uncertainty on estimates of
 262 glacier-wide WB we conduct a Monte Carlo analysis, which uses repeated random sampling of input variables
 263 to calculate a distribution of output variables (Metropolis and Ulam, 1949). We repeat the random sampling
 264 process 1000 times, resulting in a distribution of values of the glacier-wide WB based on uncertainties
 265 associated with the four steps outlined above. Density assignment uncertainty is calculated as the standard
 266 deviation of the eight resulting values of glacier-wide winter balance. Individual sources of uncertainty are

propagated through the conversion of snow depth and density measurements to glacier-wide WB. Finally, the combined effect of all three sources of uncertainty on the glacier-wide WB is quantified. We use the standard deviation of this distribution as a useful metric of uncertainty on the glacier-wide WB. We calculate a relative uncertainty as the normalized sum of differences between every pair of one hundred distributed WB estimates including σ_{GS} and σ_{INT} .

Grid-scale uncertainty (σ_{GS})

We make use of the zigzag surveys to quantify the true variability of WB at the grid scale. Our limited data do not permit a spatially-resolved assessment of grid-scale uncertainty, so we characterize this uncertainty as uniform across each glacier and represent it by a normal distribution. The distribution is centred at zero and has a standard deviation equal to the mean standard deviation of all zigzag measurements for each glacier. For each iteration of the Monte Carlo, WB values are randomly chosen from the distribution and added to the values of gridcell-averaged WB. These perturbed gridcell-averaged values of WB are then used in the interpolation. We represent uncertainty in glacier-wide WB due to grid-scale uncertainty (σ_{GS}) as the standard deviation of the resulting distribution of glacier-wide WB estimates.

Density assignment uncertainty (σ_{ρ})

We incorporate uncertainty due to the method of density assignment by carrying forward all eight density interpolation methods (Table 3) when estimating glacier-wide WB. By choosing to retain even the least plausible options, as well as the questionable FS data, this approach results in a generous assessment of uncertainty. We represent the glacier-wide WB uncertainty due to density assignment uncertainty (σ_{ρ}) as the standard deviation of glacier-wide WB estimates calculated using each density assignment method.

Interpolation uncertainty (σ_{INT})

We represent the uncertainty due to interpolation of gridcell-averaged WB in different ways for LR and SK. LR interpolation uncertainty is represented by a multivariate normal distribution of possible regression coefficients (β_i). The standard deviation of each distribution is calculated using the covariance of regression coefficients as outlined in Bagos and Adam (2015), which ensures that regression coefficients are internally consistent. The β_i distributions are randomly sampled and used to calculate gridcell-estimated WB.

SK interpolation uncertainty is represented by the standard deviation for each gridcell-estimated value of WB generated by the `DiceKriging` package. The standard deviation of glacier-wide WB is then found by taking the square root of the average variance of each gridcell-estimated WB. The final distribution of




Fig. 2. Measured snow depth and density. (a) Boxplot of measured snow depth on Glaciers 4, 2 and 13 with the first quartiles (box), median (line within box), minimum and maximum values excluding outliers (bar) and outliers (circles), which are defined as being outside of the range of 1.5 times the quartiles (approximately $\pm 2.7\sigma$). (b) Comparison of depth-averaged densities estimated using Federal Sampler (FS) measurements and a wedge cutter in a snow pit (SP) for Glacier 4 (G4), Glacier 2 (G2) and Glacier 13 (G13). Labels indicate SP locations in the accumulation area (ASP), upper ablation area (USP) and lower ablation area (LSP). Error bars for SP-derived densities are calculated by varying 100 the thickness and density of layers that are too hard to sample, and error bars for FS-derived densities are the standard deviation of measurements taken at one location. One-to-one line is dashed.

296 glacier-wide WB values is centred at the glacier-wide WB estimated with SK. For simplicity, the standard
 297 deviation of glacier-wide WB values that result from either LR or SK interpolation uncertainty is referred
 298 to as σ_{INT} .




Fig. 3. Distributions of estimated winter-balance values for each zigzag survey in lower (L), middle (M) and upper (U) ablation areas (insets). Local mean has been subtracted. (a) Glacier 4. (b) Glacier 2. (c) Glacier 13.

RESULTS AND DISCUSSION

Field measurements

Snow depth

Mean snow depth varied systematically across the study region, with Glacier 4 having the highest mean snow depth and Glacier 13 having the lowest (Fig. 2a). At each measurement location, the median range of measured depths (3–4 points) as a percent of the mean local depth is 2%, 11% and 12%, for Glaciers 4, 2 and 13, respectively. While Glacier 4 has the lowest point-scale variability, as assessed above, it also has the highest proportion of outliers, indicating a more variable snow depth across the glacier. The average standard deviation of all zigzag depth measurements is 0.07 m, 0.17 m and 0.14 m, for Glaciers 4, 2 and 13, respectively. When converted to values of WB using the local FS-derived density measurement, the average standard deviation is 0.027 m w.e., 0.035 m w.e. and 0.040 m w.e. WB data for each zigzag are not normally distributed (Fig. 3).

Snow density

Contrary to expectation, co-located FS and SP measurements are found to be uncorrelated ($R^2 = 0.25$, Fig. 2b). The FS appears to oversample in deep snow and undersample in shallow snow. Oversampling by small-diameter sampling tubes has been observed in previous studies, with a percent error between 6.8% and 11.8% (e.g. Work and others, 1965; Fames and others, 1982; Conger and McClung, 2009). Studies that use FS often apply a 10% correction to all measurements for this reason (e.g. Molotch and others, 2005). Oversampling has been attributed to slots “shaving” snow into the tube as it is rotated (e.g. Dixon and Boon, 2012) and to snow falling into the slots, particularly for snow samples with densities $>400 \text{ kg m}^{-3}$ and snow

depths >1 m (e.g. Beaumont and Work, 1963). Undersampling is likely to occur due to loss of snow from the bottom of the sampler (Turcan and Loijens, 1975). Loss by this mechanism may have occurred in our study, given the isothermal and melt-affected snow conditions observed over the lower reaches of Glaciers 2 and 13. Relatively poor FS spring-scale sensitivity also calls into question the reliability of measurements for snow depths <20 cm.

Our FS-derived density values are positively correlated with snow depth ($R^2 = 0.59$). This relationship could be a result of physical processes, such as compaction in deep snow and preferential formation of depth hoar in shallow snow, but is more likely a result of measurement artefacts for a number of reasons. First, the total range of densities measured by the FS seems improbably large (227–431 kg m⁻³). At the time of sampling the snow pack had little fresh snow, which confounds the low density values, and was not yet saturated and had few ice lenses, which confounds the high density values. Moreover, the range of FS-derived values is much larger than that of SP-derived values when co-located measurements are compared. Second, compaction effects of the magnitude required to explain the density differences between SP and FS measurements would not be expected at the measured snow depths (up to 340 cm). Third, no linear relationship exists between depth and SP-derived density ($R^2 = 0.05$). These findings suggest that the FS measurements have a bias for which we have not identified a suitable correction. Despite this bias, we use FS-derived densities to generate a range of possible WB estimates and to provide a generous estimate of uncertainty arising from density assignment.

Density assignment

Given the lack of correlation between co-located SP- and FS-derived densities (Fig. 2), we use the densities derived from these two methods separately (Table 3). SP-derived regional (S1) and glacier-mean (S2) densities are within one standard deviation of the corresponding FS-derived densities (F1 and F2) although SP-derived density values are larger (see Supplementary Material, Table S2). For both SP- and FS-derived densities, the mean density for any given glacier (S2 or F2) is within one standard deviation of the mean across all glaciers (S1 or F1). Correlations between elevation and SP- and FS-derived densities are generally high ($R^2 > 0.5$) but vary between glaciers (Supplementary material, Table S2). For any given glacier, the standard deviation of the 3–4 SP- or FS-derived densities is $<13\%$ of the mean of those values (S2 or F2) (Supplementary material, Table S2). We adopt S2 (glacier-wide mean of SP-derived densities) as the reference method of density assignment. Though the method described by S2 does not account for known basin-scale spatial variability




Fig. 4. Distribution of coefficients (β_i) determined by linear regression of gridcell-averaged WB on DEM-derived topographic parameters for the eight different density assignment methods (Table 3). Coefficients are calculated using standardized data, so values can be compared across parameters. Regression coefficients that are not significant are assigned a value of zero. Topographic parameters include elevation (z), distance from centreline (d_C), slope (m), curvature (κ) and wind redistribution (Sx). Aspect (α) and “northness” (N) are not shown because coefficient values are zero in every case. The box plot shows first quartiles (box), median (line within box), mean (circle within box), minimum and maximum values excluding outliers (bars) and outliers (gray dots), which are defined as being outside of the range of 1.5 times the quartiles (approximately $\pm 2.7\sigma$).

in snow density (e.g. Wetlaufer and others, 2016), it is commonly used in winter balance studies (e.g. Elder and others, 1991; McGrath and others, 2015; Cullen and others, 2017).

Gridcell-averaged winter balance

The distributions of gridcell-averaged WB values for the individual glaciers are similar to those in Fig. 2a but with fewer outliers (see Supplementary Material). The standard deviations of WB values determined from the zigzag surveys are almost twice as large as the mean standard deviation of point-scale WB values within a gridcell measured along transects (see Supplementary Material). However, a small number of gridcells sampled in transect surveys have standard deviations in WB that exceed 0.25 m w.e. (~ 10 times greater than those for zigzag surveys).




Fig. 5. Spatial distribution of winter balance (WB) estimated using linear regression (top row) and simple kriging (bottom row) with densities assigned as per S2 (Table 3). The linear regression method involves multiplying regression coefficients, found using cross validation and model averaging, by topographic parameters for each gridcell. Simple kriging uses the covariance of measured values to find a set of optimal weights for estimating values at unmeasured locations. Locations of snow-depth measurements taken in May 2016 are shown as black dots. Ice-flow directions are indicated by arrows. Values of glacier-wide WB are given below labels.

Distributed winter balance

Linear Regression

Of the topographic parameters in the LR, elevation (z) is the most significant predictor of gridcell-averaged WB for Glaciers 2 and 13, while wind redistribution (Sx) is the most significant predictor for Glacier 4

Table 4. Glacier-wide winter balance (WB, m w.e.) estimated using linear regression and simple kriging for the three study glaciers. Root mean squared error (RMSE, m w.e.) is computed as the average of all RMSE values between gridcell-averaged values of WB (the data) that were randomly selected and excluded from interpolation (1/3 of all data) and those estimated by interpolation. RMSE as a percent of the glacier-wide WB is shown in brackets.

	Linear regression		Simple kriging	
	WB	RMSE	WB	RMSE
G4	0.58	0.15 (26%)	0.62	0.13 (21%)
G2	0.58	0.10 (17%)	0.37	0.07 (19%)
G13	0.38	0.08 (21%)	0.27	0.07 (26%)




Fig. 6. Relative uncertainty in distributed winter balance (WB) (Fig. 5) found using linear regression (top row) and simple kriging (bottom row). Values closer to one indicate higher relative uncertainty. Ice-flow directions are indicated by arrows.

361 (Fig. 4). As expected, gridcell-averaged WB is positively correlated with elevation where the correlation
 362 is significant. It is possible that the elevation correlation was accentuated due to melt onset for Glacier
 363 13 in particular. Our results are consistent with many studies that have found elevation to be the most
 364 significant predictor of winter-balance data (e.g. Machguth and others, 2006; McGrath and others, 2015).
 365 The WB–elevation gradient is 13 mm/100 m on Glacier 2 and 7 mm/100 m on Glacier 13. These gradients
 366 are consistent with those reported for a few glaciers in Svalbard (Winther and others, 1998) but considerably
 367 smaller than many reported WB–elevation gradients, which range from about 60–240 mm/100 m (e.g. Hagen
 368 and Liestøl, 1990; Tveit and Killingtveit, 1994; Winther and others, 1998). Extrapolating linear relationships
 369 to unmeasured locations typically results in large uncertainties, as seen by the large WB values (Fig. 5) and
 370 large relative uncertainty (Fig. 6) in the high-elevation regions of the accumulation areas of Glaciers 2 and
 371 13. The low of correlation between WB and elevation Glacier 4 is consistent with Grabiec and others (2011)
 372 and López-Moreno and others (2011), who conclude that highly variable distributions of snow are attributed
 373 to complex interactions between topography and the atmosphere that could not be easily quantified.




Fig. 7. Winter balance (WB) estimated by linear regression (LR, top row) or simple kriging (SK, bottom row) versus the grid-cell averaged WB data for Glacier 4 (left), Glacier 2 (middle) and Glacier 13 (right). Each circle represents a single gridcell. Best-fit (solid) and one-to-one (dashed) lines are shown.

Gridcell-averaged WB is negatively correlated with Sx on Glacier 4, counter-intuitively indicating less snow in what would be interpreted as sheltered areas. Gridcell-averaged WB is positively correlated with Sx on Glaciers 2 and 13. Similarly, gridcell-averaged WB is positively correlated with curvature on Glacier 4 and negatively correlated on Glaciers 2 and 13. Our results corroborate those of McGrath and others (2015) in a study of six glaciers in Alaska (DEM resolutions of 5 m) where elevation and Sx were the only significant parameters for all glaciers; Sx regression coefficients were smaller than elevation regression coefficients, and in some cases, negative. While our results point to wind having an impact on snow distribution, the wind redistribution parameter (Sx) may not adequately capture these effects at our study sites. For example, Glacier 4 is located in a curved valley with steep side walls, so specifying a single cardinal direction for wind may not be adequate. Further, the scale of deposition may be smaller than the resolution of the Sx parameter estimated from the DEM. Creation of a parametrization for sublimation from blowing snow, which has been shown to be an important mechanism of mass loss from ridges (e.g. Musselman and others, 2015), may also improve explanatory power of LR for our study sites.

We find that transfer of LR coefficients between glaciers results in large estimation errors. Regression coefficients from Glacier 4 produce the highest root mean squared error (0.38 m w.e. on Glacier 2 and 0.40 m w.e. on Glacier 13, see Table 4 for comparison) and glacier-wide WB values are the same for all

glaciers (0.64 m w.e.) due to the dominance of the regression intercept. Even if the LR is performed with WB values from all glaciers combined, the resulting coefficients produce large root mean squared errors when applied to individual glaciers (0.31 m w.e., 0.15 m w.e. and 0.14 m w.e. for Glaciers 4, 2 and 13, respectively). Our results are consistent with those of Grünewald and others (2013), who found that local statistical models cannot be transferred across basins and that regional-scale models are not able to explain the majority of observed variance in winter balance.

Simple kriging

Fitted kriging parameters, including the nugget and spatial correlation length, can provide insight into important scales of winter-balance variability. The model fitted to the gridcell-averaged values of WB for Glacier 4 has a short correlation length (90 m) and large nugget (see Supplementary Material Table S3), suggesting variability in winter balance at smaller scales. Conversely, Glaciers 2 and 13 have longer correlation lengths (~ 450 m) and smaller nuggets, suggesting variability at larger scales. Additionally, SK is better able to estimate values of WB for Glaciers 2 and 13 than for Glacier 4 (Fig. 7). Due to a paucity of data, SK produces almost uniform gridcell-estimated values of winter balance in the accumulation area of each glacier, inconsistent with observations described in the literature (e.g. Machguth and others, 2006; Grabiec and others, 2011). As expected, extrapolation using SK leads to large uncertainty (Fig. 6), further emphasizing the need for spatially distributed point-scale measurements.

LR and SK comparison

Glacier-wide WB estimates found using both LR and SK are ~ 0.58 m w.e. for Glacier 4 but both are poor predictors of WB in measured gridcells (Table 4). For Glaciers 2 and 13, SK estimates are more than 0.1 m w.e. (up to 40%) lower than LR estimates (Table 4). RMSE as a percentage of the glacier-wide WB are comparable between LR and SK (Table 4) with an average RMSE of 22%. This comparability is interesting, given that all of the data were used to generate the SK model, while only 2/3 were used in the LR (consistent with the best SK model estimated with 2/3 of the data). Gridcell-estimated values of WB found using LR and SK differ markedly in the upper accumulation areas of Glaciers 2 and 13 (Fig. 5), where observations are sparse and topographic parameters, such as elevation, vary considerably. The influence of elevation results in substantially higher LR-estimated values of WB at high elevation, whereas SK-estimated values approximate the nearest data. Estimates of ablation-area-wide WB differ by $<7\%$ between LR and SK on each glacier, further emphasizing the combined role of interpolation method and measurement scarcity in the accumulation area on glacier-wide WB estimates.




Fig. 8. Distributions of glacier-wide winter balance (WB) for Glaciers 4 (G4), 2 (G2) and 13 (G13) that arise from various sources of uncertainty. WB distribution arising from grid-scale uncertainty (σ_{GS}) (left column). WB distribution arising from interpolation uncertainty (σ_{INT}) (middle column). WB distribution arising from a combination of σ_{GS} , σ_{INT} and density assignment uncertainty (σ_{ρ}) (right column). Results are shown for interpolation by linear regression (LR, top row) and simple kriging (SK, bottom row). Left two columns include eight distributions per glacier (colour) and each corresponds to a density assignment method (S1–S4 and F1–F4).

Uncertainty analysis

Glacier-wide winter balance is affected by uncertainty introduced by the representativeness of gridcell-averaged values of WB (σ_{GS}), choosing a method of density assignment (σ_{ρ}), and interpolating/extrapolating WB values across the domain (σ_{INT}). Using a Monte Carlo analysis, we find that interpolation uncertainty contributes more to WB uncertainty than grid-scale uncertainty or density assignment method. In other words, the distribution of glacier-wide WB that arises from grid-scale uncertainty and the differences in distributions between methods of density assignment are smaller than the distribution that arises from interpolation uncertainty (Fig. 8 and Table 5). The WB distributions obtained using LR and SK overlap for a given glacier, but the distribution modes differ (Fig. 8). For reasons outlined above, SK-estimated values of WB in the accumulation area are generally lower, which lowers the glacier-wide WB estimate. The uncertainty in SK-estimated values of WB is large, and unrealistic glacier-wide values of WB of 0 m w.e. can be estimated (Fig. 8). Our results caution strongly against including interpolated/extrapolated values of WB in comparisons with remote sensing- or model-derived estimates of WB. If possible, such comparisons should be restricted to point-scale data.

Table 5. Standard deviation ($\times 10^{-2}$ m w.e.) of glacier-wide winter balance (WB) distributions arising from uncertainties in grid-scale WB (σ_{GS}), density assignment (σ_{ρ}), interpolation (σ_{INT}) and all three sources combined (σ_{ALL}) for linear regression (left columns) and simple kriging (right columns)

	Linear regression				Simple kriging			
	σ_{GS}	σ_{ρ}	σ_{INT}	σ_{ALL}	σ_{GS}	σ_{ρ}	σ_{INT}	σ_{ALL}
Glacier 4	0.86	1.90	2.13	2.90	0.85	2.15	14.05	14.72
Glacier 2	1.80	3.37	3.09	4.90	2.53	2.03	13.78	13.44
Glacier 13	1.12	1.68	2.80	3.20	1.15	1.27	9.65	10.43

Grid-scale uncertainty (σ_{GS}) is the smallest assessed contributor to overall WB uncertainty. This result is consistent with the generally smoothly-varying snow depths encountered in zigzag surveys, and previously reported ice-roughness lengths on the order of centimetres (e.g. Hock, 2005) compared to snow depths on the order of decimetres to metres. Given our assumption that zigzags are an adequate representation of grid-scale variability, the low WB uncertainty arising from σ_{GS} implies that subgrid-scale sampling need not be a high priority for reducing overall uncertainty. Our assumption that the 3–4 zigzag surveys can be used to estimate glacier-wide σ_{GS} may be flawed, particularly in areas with debris cover, crevasses and steep slopes.

Our analysis did not include uncertainty arising from density measurement errors associated with the FS, wedge cutters and spring scales, from vertical and horizontal errors in the DEM or from error associated with estimating measurement locations based on the GPS position of the lead observer. We assume that these sources of uncertainty are either encompassed by the sources investigated or negligible.

The values of glacier-wide WB for our study glaciers (using LR and S2 density assignment method), with an uncertainty equal to one standard deviation of the distribution found with Monte Carlo analysis, are: 0.59 ± 0.03 m w.e. for Glacier 4, 0.61 ± 0.05 m w.e. for Glacier 2 and 0.40 ± 0.03 m w.e. for Glacier 13. The glacier-wide WB uncertainty from the three investigated sources of uncertainty ranges from 0.03 m w.e (5%) to 0.05 m w.e (8%) for LR estimates and from 0.10 m w.e (37%) to 0.15 m w.e (24%) for simple-kriging estimates (Table 4).

Context and caveats

Regional winter-balance gradient

Although we find considerable inter- and intra-basin variability in winter balance, our results are consistent with a regional-scale winter-balance gradient for the continental side of the St. Elias Mountains (Fig. 9). Winter-balance data are compiled from Taylor-Barge (1969), the three glaciers presented in this paper and




Fig. 9. Relationship between winter balance (WB) and linear distance from the regional topographic divide between the Kaskawulsh and Hubbard Glaciers in the St. Elias Mountains. Point-scale values of WB from snow-pit data reported by Taylor-Barge (1969) (blue boxes, P-WB). LR-estimated glacier-wide WB calculated using density assignment S2 for Glaciers 4 (G4), 2 (G2) and 13 (G13) with errors bars calculated as the standard deviation of Monte Carlo-derived WB distributions (this study) (open orange circles, G-WB). Point-scale WB estimated from snow-pit data at two locations in the accumulation area of the Kaskawulsh Glacier, collected in May 2016 (unpublished data, SFU Glaciology Group) (filled orange dots, P-WB). Black line indicates best fit ($R^2 = 0.85$).

two snow pits we analyzed near the head of the Kaskawulsh Glacier between 20–21 May 2016. The data show a linear decrease of $0.024 \text{ m w.e. km}^{-1}$ ($R^2 = 0.85$) in winter balance with distance from the regional topographic divide between the Kaskawulsh and Hubbard Glaciers, as identified by Taylor-Barge (1969). While the three study glaciers fit the regional trend, the same relationship would not apply if just the Donjek Range were considered. We hypothesize that interaction between meso-scale weather patterns and large-scale mountain topography is a major driver of regional-scale winter balance. Further insight into regional-scale patterns of winter balance in the St. Elias Mountains could be gained by investigating moisture source trajectories and the contribution of orographic precipitation.

Limitations and future work

The potential limitations of our work include the restriction of our data to a single year, minimal sampling in the accumulation area, the problem of uncorrelated SP- and FS-derived densities, a sampling design that could not be optimized a priori, the assumption of spatially uniform subgrid variability and lack of more finely resolved DEMs.

Inter-annual variability in winter balance is not considered in our study. A number of studies have found temporal stability in spatial patterns of snow distribution and that statistical models based on topographic

parameters could be applied reliably between years (e.g. Grünewald and others, 2013). For example, Walmsley (2015) analyzed more than 40 years of winter balance recorded on two Norwegian glaciers and found that snow distribution is spatially heterogeneous yet exhibits robust temporal stability. Contrary to this, Crochet and others (2007) found that snow distribution in Iceland differed considerably between years and depended primarily on the dominant wind direction over the course of a winter. Therefore, multiple years of snow depth and density measurements, that are not necessarily consecutive, are needed to better understand inter-annual variability in winter-balance distribution within the Donjek Range.

There is a conspicuous lack of data in the accumulation areas of our study glaciers. With increased sampling in the accumulation area, interpolation uncertainties would be reduced where they are currently greatest and the LR would be better constrained. Although certain regions of the glaciers remain inaccessible for direct measurements, other methods of obtaining winter-balance, including ground-penetrating radar and DEM differencing with photogrammetry or lidar, could be used in conjunction with manual probing to increase the spatial coverage of measurements.

The lack of correlation between SP- and FS-derived densities needs to be reconciled. Contrary to our results, most studies that compare SP- and FS-derived densities report minimal discrepancy (e.g. Dixon and Boon, 2012, and sources within). Additional co-located density measurements are needed to better compare the two methods of obtaining density values. Comparison with other FS would also be informative. Even with this limitation, density assignment was, fortunately, not the largest source of uncertainty in estimating glacier-wide winter balance.

Our sampling design was chosen to achieve broad spatial coverage of the ablation area, but is likely too finely resolved along transects for many mass-balance surveys to replicate. An optimal sampling design would minimize uncertainty in winter balance while reducing the number of required measurements. Analysis of the estimated winter balance obtained using subsets of the data is underway to make recommendations on optimal transect configuration and along-track spacing of measurements. López-Moreno and others (2010) found that 200–400 observations are needed within a non-glacierized alpine basin (6 km²) to obtain accurate and robust snow distribution models. Similar guidelines would be useful for glacierized environments.

In this study, we assume that the subgrid variability of winter balance is uniform across a given glacier. Contrary to this assumption, McGrath and others (2015) found greater variability of winter-balance values close to the terminus. Testing our assumption could be a simple matter of prioritizing the labour-intensive

500 zigzags surveys. To ensure consistent quantification of subgrid variability, zigzag survey measurements could
501 also be tested against other measurements methods, such as lidar.

502 DEM gridcell size is known to influence values of computed topographic parameters (Zhang and
503 Montgomery, 1994; Garbrecht and Martz, 1994; Guo-an and others, 2001; López-Moreno and others, 2010).
504 The relationship between topographic parameters and winter balance is, therefore, not independent of DEM
505 gridcell size. For example, Kienzle (2004) and López-Moreno and others (2010) found that a decrease in
506 spatial resolution of the DEM results in a decrease in the importance of curvature and an increase in the
507 importance of elevation in LR of snow distribution on topographic parameters in non-glacierized basins.
508 The importance of curvature in our study is affected by the DEM smoothing that we applied to obtain a
509 spatially continuous curvature field (see Supplementary Material). A comparison of regression coefficients
510 from high-resolution DEMs obtained from various sources and sampled with various gridcell sizes could be
511 used to characterize the dependence of topographic parameters on DEMs, and therefore assess the robustness
512 of inferred relationships between winter balance and topographic parameters.

513 CONCLUSION

514 We estimate winter balance for three glaciers (termed Glacier 2, Glacier 4 and Glacier 13) in the St. Elias
515 Mountains, Yukon, Canada from multiscale snow depth and density measurements. Linear regression and
516 simple kriging are used to obtain estimates of distributed winter balance. We use Monte Carlo analysis to
517 evaluate the contributions of interpolation, the assignment of snow density and grid-scale variability of winter
518 balance to uncertainty in glacier-wide winter balance.

519 Values of glacier-wide winter balance estimated using linear regression and simple kriging differ by up
520 to 0.24 m w.e. ($\sim 50\%$). We find that interpolation uncertainty is the largest assessed source of uncertainty
521 in glacier-wide winter balance (5% for linear-regression estimates and 32% for simple-kriging estimates).
522 Uncertainty resulting from the method of density assignment is comparatively low, despite the wide range of
523 methods explored. Given our representation of grid-scale variability, the resulting winter balance uncertainty
524 is small indicating that extensive subgrid-scale sampling is not required to reduce overall uncertainty.

525 Our results suggest that processes governing distributed winter balance differ between glaciers, highlighting
526 the importance of regional-scale winter-balance studies. The estimated distribution of winter balance on
527 Glacier 4 is characterized by high variability, as indicated by the poor correlation between estimated and
528 observed values and large number of data outliers. Glaciers 2 and 13 appear to have lower spatial variability,
529 with elevation being the dominant predictor of gridcell-averaged winter balance. A wind-redistribution

parameter is found to be a weak but significant predictor of winter balance, though conflicting relationships between glaciers make it difficult to interpret. The major limitations of our work include the restriction of our data to a single year and minimal sampling in the accumulation area. Although challenges persist when estimating winter balance, our data are consistent with a regional-scale winter-balance gradient for the continental side of the St. Elias Mountains.

AUTHOR CONTRIBUTION STATEMENT

AP planned and executed the data collection, performed all calculations and drafted the manuscript. GF conceived of the study, contributed to field planning and data collection, oversaw all stages of the work and edited the manuscript. VR provided guidance with statistical methods and edited the manuscript.

ACKNOWLEDGEMENTS

We thank the Kluane First Nation (KFN), Parks Canada and the Yukon Territorial Government for granting us permission to work in KFN Traditional Territory and Kluane National Park and Reserve. We are grateful for financial support provided by the Natural Sciences and Engineering Research Council of Canada, Simon Fraser University and the Northern Scientific Training Program. We kindly acknowledge Kluane Lake Research Station, Sian Williams, Lance Goodwin and Trans North pilot Dion Parker for facilitating field logistics. We are grateful to Alison Criscitiello and Coline Ariagno for all aspects of field assistance and Sarah Furney for assistance with data entry. Thank you to Etienne Berthier for providing us with the SPIRIT SPOT-5 DEM and for assistance in DEM correction. We are grateful to Derek Bingham and Michael Grosskopf for assistance with the statistics, including simple kriging. Luke Wonneck, Leif Anderson and Jeff Crompton all provided thoughtful and constructive comments on drafts of the manuscript.

REFERENCES

- Anderton S, White S and Alvera B (2004) Evaluation of spatial variability in snow water equivalent for a high mountain catchment. *Hydrological Processes*, **18**(3), 435–453 (doi: 10.1002/hyp.1319)
- Arendt AA, Luthcke SB, Larsen CF, Abdalati W, Krabill WB and Beedle MJ (2008) Validation of high-resolution GRACE mascon estimates of glacier mass changes in the St Elias Mountains, Alaska, USA, using aircraft laser altimetry. *Journal of Glaciology*, **54**(188), 778–787 (doi: 10.3189/002214308787780067)
- Bagos PG and Adam M (2015) On the Covariance of Regression Coefficients. *Open Journal of Statistics*, **5**, 680–701 (doi: 10.4236/ojs.2015.57069)

- 558 Balk B and Elder K (2000) Combining binary decision tree and geostatistical methods to estimate snow distribution
559 in a mountain watershed. *Water Resources Research*, **36**(1), 13–26 (doi: 10.1029/1999WR900251)
- 560 Barry RG (1992) *Mountain weather and climate*. Cambridge University Press, 3rd edition
- 561 Beaumont RT and Work RA (1963) Snow sampling results from three samplers. *International Association of Scientific*
562 *Hydrology. Bulletin*, **8**(4), 74–78 (doi: 10.1080/02626666309493359)
- 563 Burgess EW, Forster RR and Larsen CF (2013) Flow velocities of Alaskan glaciers. *Nature communications*, **4**,
564 2146–2154 (doi: 10.1038/ncomms3146)
- 565 Burnham KP and Anderson DR (2004) Multimodel Inference: Understanding AIC and BIC in Model Selection.
566 *Sociological Methods & Research*, **33**(2), 261–304 (doi: 10.1177/0049124104268644)
- 567 Carroll T (1977) A comparison of the CRREL 500 cm³ tube and the ILTS 200 and 100 cm³ box cutters used for
568 determining snow densities. *Journal of Glaciology*, **18**(79), 334–337 (doi: 10.1017/S0022143000021420)
- 569 Clark MP, Hendrikx J, Slater AG, Kavetski D, Anderson B, Cullen NJ, Kerr T, Örn Hreinsson E and Woods RA
570 (2011) Representing spatial variability of snow water equivalent in hydrologic and land-surface models: A review.
571 *Water Resources Research*, **47**(7) (doi: 10.1029/2011WR010745)
- 572 Clarke GK (2014) A short and somewhat personal history of Yukon glacier studies in the Twentieth Century. *Arctic*,
573 **37**(1), 1–21
- 574 Clyde GD (1932) Circular No. 99-Utah Snow Sampler and Scales for Measuring Water Content of Snow. *UAES*
575 *Circulars*, Paper 90
- 576 Cogley J, Hock R, Rasmussen L, Arendt A, Bauder A, Braithwaite R, Jansson P, Kaser G, Möller M, Nicholson L
577 and others (2011) *Glossary of glacier mass balance and related terms*. UNESCO-IHP, Paris
- 578 Conger SM and McClung DM (2009) Comparison of density cutters for snow profile observations. *Journal of*
579 *Glaciology*, **55**(189), 163–169 (doi: 10.3189/002214309788609038)
- 580 Crochet P, Jóhannesson T, Jónsson T, Sigurðsson O, Björnsson H, Pálsson F and Barstad I (2007) Estimating
581 the Spatial Distribution of Precipitation in Iceland Using a Linear Model of Orographic Precipitation. *Journal of*
582 *Hydrometeorology*, **8**(6), 1285–1306 (doi: 10.1175/2007JHM795.1)
- 583 Crompton JW and Flowers GE (2016) Correlations of suspended sediment size with bedrock lithology and glacier
584 dynamics. *Annals of Glaciology*, **57**(72), 1–9 (doi: 10.1017/aog.2016.6)
- 585 Cullen NJ, Anderson B, Sirguey P, Stumm D, Mackintosh A, Conway JP, Horgan HJ, Dadić R, Fitzsimons SJ
586 and Lorrey A (2017) An 11-year record of mass balance of Brewster Glacier, New Zealand, determined using a
587 geostatistical approach. *Journal of Glaciology*, **63**(238), 199–217 (doi: 10.1017/jog.2016.128)
- 588 Dadić R, Mott R, Lehning M and Burlando P (2010) Parameterization for wind-induced preferential deposition of
589 snow. *Journal of Geophysical Research: Earth Surface*, **24**(14), 1994–2006 (doi: 10.1029/2009JF001261)

- 590 Danby RK, Hik DS, Slocombe DS and Williams A (2003) Science and the St. Elias: an evolving framework
591 for sustainability in North America's highest mountains. *The Geographical Journal*, **169**(3), 191–204 (doi:
592 10.1111/1475-4959.00084)
- 593 Davis JC and Sampson RJ (1986) *Statistics and data analysis in geology*. Wiley New York et al., 2nd edition
- 594 Deems JS and Painter TH (2006) Lidar measurement of snow depth: accuracy and error sources. In *Proceedings of*
595 *the International Snow Science Workshop*
- 596 Dixon D and Boon S (2012) Comparison of the SnowHydro snow sampler with existing snow tube designs. *Hydrological*
597 *Processes*, **26**(17), 2555–2562 (doi: 10.1002/hyp.9317)
- 598 Egli L, Griessinger N and Jonas T (2011) Seasonal development of spatial snow-depth variability across different
599 scales in the Swiss Alps. *Annals of Glaciology*, **52**(58), 216–222 (doi: 10.3189/172756411797252211)
- 600 Elder K, Dozier J and Michaelsen J (1991) Snow accumulation and distribution in an alpine watershed. *Water*
601 *Resources Research*, **27**(7), 1541–1552 (doi: 10.1029/91WR00506)
- 602 Elder K, Rosenthal W and Davis RE (1998) Estimating the spatial distribution of snow water equiv-
603 alence in a montane watershed. *Hydrological Processes*, **12**(1011), 1793–1808 (doi: 10.1002/(SICI)1099-
604 1085(199808/09)12:10<111793::AID-HYP6953.0.CO;2-K)
- 605 Erxleben J, Elder K and Davis R (2002) Comparison of spatial interpolation methods for estimating snow distribution
606 in the Colorado Rocky Mountains. *Hydrological Processes*, **16**(18), 3627–3649 (doi: 10.1002/hyp.1239)
- 607 Fames PE, Peterson N, Goodison B and Richards RP (1982) Metrication of Manual Snow Sampling Equipment. In
608 *Proceedings of the 50th Western Snow Conference*, 120–132
- 609 Fierz C, Armstrong RL, Durand Y, Etchevers P, Greene E, McClung DM, Nishimura K, Satyawali PK and Sokratov
610 SA (2009) *The international classification for seasonal snow on the ground*. UNESCO/IHP, unesco/ihp paris
611 edition
- 612 Garbrecht J and Martz L (1994) Grid size dependency of parameters extracted from digital elevation models.
613 *Computers & Geosciences*, **20**(1), 85–87 (doi: 10.1016/0098-3004(94)90098-1)
- 614 Gerber F, Lehning M, Hoch SW and Mott R (2017) A close-ridge small-scale atmospheric flow field and its
615 influence on snow accumulation. *Journal of Geophysical Research: Atmospheres*, **122**(15), 7737–7754 (doi:
616 10.1002/2016JD026258), 2016JD026258
- 617 Grabiec M, Puczek D, Budzik T and Gajek G (2011) Snow distribution patterns on Svalbard glaciers derived from
618 radio-echo soundings. *Polish Polar Research*, **32**(4), 393–421 (doi: 10.2478/v10183-011-0026-4)
- 619 Gray DM and Male DH (1981) *Handbook of snow: principles, processes, management & use*. Pergamon Press, 1st
620 edition
- 621 Grünewald T, Schirmer M, Mott R and Lehning M (2010) Spatial and temporal variability of snow depth and ablation
622 rates in a small mountain catchment. *Cryosphere*, **4**(2), 215–225 (doi: 10.5194/tc-4-215-2010)

- Grünwald T, Stötter J, Pomeroy J, Dadic R, Moreno Baños I, Marturià J, Spross M, Hopkinson C, Burlando P and Lehning M (2013) Statistical modelling of the snow depth distribution in open alpine terrain. *Hydrology and Earth System Sciences*, **17**(8), 3005–3021 (doi: 10.5194/hess-17-3005-2013)
- Guo-an T, Yang-he H, Strobl J and Wang-qing L (2001) The impact of resolution on the accuracy of hydrologic data derived from DEMs. *Journal of Geographical Sciences*, **11**(4), 393–401 (doi: 10.1007/BF02837966)
- Gusmeroli A, Wolken GJ and Arendt AA (2014) Helicopter-borne radar imaging of snow cover on and around glaciers in Alaska. *Annals of Glaciology*, **55**(67), 78–88 (doi: 10.3189/2014AoG67A029)
- Hagen JO and Liestøl O (1990) Long-term glacier mass-balance investigations in Svalbard, 1950–88. *Annals of Glaciology*, **14**(1), 102–106
- Helbig N and van Herwijnen A (2017) Subgrid parameterization for snow depth over mountainous terrain from flat field snow depth. *Water Resources Research*, **53**(2), 1444–1456 (doi: 10.1002/2016WR019872)
- Hock R (2005) Glacier melt: a review of processes and their modelling. *Progress in Physical Geography*, **29**(3), 362–391 (doi: 10.1191/0309133305pp453ra)
- Hock R and Jensen H (1999) Application of kriging interpolation for glacier mass balance computations. *Geografiska Annaler: Series A, Physical Geography*, **81**(4), 611–619 (doi: 10.1111/1468-0459.00089)
- Kaser G, Fountain A, Jansson P and others (2003) *A manual for monitoring the mass balance of mountain glaciers*. ICSI/UNESCO
- Kienzle S (2004) The Effect of DEM Raster Resolution on First Order, Second Order and Compound Terrain Derivatives. *Transactions in GIS*, **8**(1), 83–111 (doi: 10.1111/j.1467-9671.2004.00169.x)
- Kohavi R and others (1995) A study of cross-validation and bootstrap for accuracy estimation and model selection. In *Proceedings of the Fourteenth International Joint Conference on Artificial Intelligence*, 1137–1145
- Korona J, Berthier E, Bernard M, Rémy F and Thouvenot E (2009) SPIRIT SPOT 5 stereoscopic survey of Polar Ice: Reference images and topographies during the fourth International Polar Year (2007–2009). *ISPRS Journal of Photogrammetry and Remote Sensing*, **64**(2), 204–212 (doi: 10.1016/j.isprsjprs.2008.10.005)
- Lehning M, Völksch I, Gustafsson D, Nguyen TA, Stähli M and Zappa M (2006) ALPINE3D: a detailed model of mountain surface processes and its application to snow hydrology. *Hydrological Processes*, **20**(10), 2111–2128 (doi: 10.1002/hyp.6204)
- Li J and Heap AD (2008) A review of spatial interpolation methods for environmental scientists. *Geoscience Australia*, Record 2008/23
- Liston GE and Elder K (2006) A distributed snow-evolution modeling system (SnowModel). *Journal of Hydrometeorology*, **7**(6), 1259–1276 (doi: 10.1175/JHM548.1)
- Liston GE and Sturm M (1998) A snow-transport model for complex terrain. *Journal of Glaciology*, **44**(148), 498–516

- 655 López-Moreno J, Latron J and Lehmann A (2010) Effects of sample and grid size on the accuracy and stability of
656 regression-based snow interpolation methods. *Hydrological Processes*, **24**(14), 1914–1928 (doi: 10.1002/hyp.7564)
- 657 López-Moreno J, Fassnacht S, Heath J, Musselman K, Revuelto J, Latron J, Morán-Tejeda E and Jonas T (2013)
658 Small scale spatial variability of snow density and depth over complex alpine terrain: Implications for estimating
659 snow water equivalent. *Advances in Water Resources*, **55**, 40–52 (doi: 10.1016/j.advwatres.2012.08.010)
- 660 López-Moreno JJ, Fassnacht S, Beguería S and Latron J (2011) Variability of snow depth at the plot scale: implications
661 for mean depth estimation and sampling strategies. *The Cryosphere*, **5**(3), 617–629 (doi: 10.5194/tc-5-617-2011)
- 662 MacDougall AH and Flowers GE (2011) Spatial and temporal transferability of a distributed energy-balance glacier
663 melt model. *Journal of Climate*, **24**(5), 1480–1498 (doi: 10.1175/2010JCLI3821.1)
- 664 Machguth H, Eisen O, Paul F and Hoelzle M (2006) Strong spatial variability of snow accumulation observed
665 with helicopter-borne GPR on two adjacent alpine glaciers. *Geophysical Research Letters*, **33**(13), 1–5 (doi:
666 10.1029/2006GL026576)
- 667 Madigan D and Raftery AE (1994) Model Selection and Accounting for Model Uncertainty in Graphical Models
668 Using Occam’s Window. *Journal of the American Statistical Association*, **89**(428), 1535–1546
- 669 Marshall HP, Koh G, Sturm M, Johnson J, Demuth M, Landry C, Deems J and Gleason J (2006) Spatial variability of
670 the snowpack: Experiences with measurements at a wide range of length scales with several different high precision
671 instruments. In *Proceedings International Snow Science Workshop*, 359–364
- 672 McGrath D, Sass L, O’Neel S, Arendt A, Wolken G, Gusmeroli A, Kienholz C and McNeil C (2015) End-of-winter
673 snow depth variability on glaciers in Alaska. *Journal of Geophysical Research: Earth Surface*, **120**(8), 1530–1550
674 (doi: 10.1002/2015JF003539)
- 675 Metropolis N and Ulam S (1949) The Monte Carlo Method. *Journal of the American Statistical Association*, **44**(247),
676 335–341
- 677 Molotch N, Colee M, Bales R and Dozier J (2005) Estimating the spatial distribution of snow water equivalent in
678 an alpine basin using binary regression tree models: the impact of digital elevation data and independent variable
679 selection. *Hydrological Processes*, **19**(7), 1459–1479 (doi: 10.1002/hyp.5586)
- 680 Mott R, Faure F, Lehning M, Löwe H, Hynek B, Michlmayer G, Prokop A and Schöner W (2008) Simulation of
681 seasonal snow-cover distribution for glacierized sites on Sonnblick, Austria, with the Alpine3D model. *Annals of*
682 *Glaciology*, **49**(1), 155–160 (doi: 10.3189/172756408787814924)
- 683 Musselman KN, Pomeroy JW, Essery RL and Leroux N (2015) Impact of windflow calculations on simulations
684 of alpine snow accumulation, redistribution and ablation. *Hydrological Processes*, **29**(18), 3983–3999 (doi:
685 10.1002/hyp.10595)
- 686 Proksch M, Rutter N, Fierz C and Schneebeli M (2016) Intercomparison of snow density measurements: bias, precision,
687 and vertical resolution. *The Cryosphere*, **10**(1), 371–384 (doi: 10.5194/tc-10-371-2016)

- 688 Raftery AE, Madigan D and Hoeting JA (1997) Bayesian Model Averaging for Linear Regression Models. *Journal of*
689 *the American Statistical Association*, **92**(437), 179–191 (doi: 10.1080/01621459.1997.10473615)
- 690 Rasmussen CE and Williams CK (2006) *Gaussian processes for machine learning*. MIT press Cambridge
- 691 Réveillet M, Vincent C, Six D and Rabatel A (2016) Which empirical model is best suited to simulate glacier mass
692 balances? *Journal of Glaciology*, **63**(237), 1–16 (doi: 10.1017/jog.2016.110)
- 693 Roustant O, Ginsbourger D and Deville Y (2012) DiceKriging, DiceOptim: Two R packages for the analysis of
694 computer experiments by kriging-based metamodeling and optimization. *Journal of Statistical Software*, **21**, 1–55
- 695 Schneiderbauer S and Prokop A (2011) The atmospheric snow-transport model: SnowDrift3D. *Journal of Glaciology*,
696 **57**(203), 526–542 (doi: 10.3189/002214311796905677)
- 697 Schuler TV, Crochet P, Hock R, Jackson M, Barstad I and Jóhannesson T (2008) Distribution of snow accumulation
698 on the Svartisen ice cap, Norway, assessed by a model of orographic precipitation. *Hydrological Processes*, **22**(19),
699 3998–4008 (doi: 10.1002/hyp.7073)
- 700 Shea C and Jamieson B (2010) Star: an efficient snow point-sampling method. *Annals of Glaciology*, **51**(54), 64–72
701 (doi: 10.3189/172756410791386463)
- 702 Sold L, Huss M, Hoelzle M, Anderegg H, Joerg PC and Zemp M (2013) Methodological approaches to
703 infer end-of-winter snow distribution on alpine glaciers. *Journal of Glaciology*, **59**(218), 1047–1059 (doi:
704 10.3189/2013JoG13J015)
- 705 Tangborn WV, Krimmel RM and Meier MF (1975) A comparison of glacier mass balance by glaciological, hydrological
706 and mapping methods, South Cascade Glacier, Washington. *International Association of Hydrological Sciences*
707 *Publication*, **104**, 185–196
- 708 Taylor-Barge B (1969) *The summer climate of the St. Elias Mountain region*. Montreal: Arctic Institute of North
709 America, Research Paper No. 53
- 710 Thibert E, Blanc R, Vincent C and Eckert N (2008) Glaciological and volumetric mass-balance measurements: error
711 analysis over 51 years for Glacier de Sarennes, French Alps. *Journal of Glaciology*, **54**(186), 522–532
- 712 Trujillo E and Lehning M (2015) Theoretical analysis of errors when estimating snow distribution through point
713 measurements. *The Cryosphere*, **9**(3), 1249–1264 (doi: 10.5194/tc-9-1249-2015)
- 714 Turcan J and Loijens H (1975) Accuracy of snow survey data and errors in snow sampler measurements. In *32nd*
715 *Eastern Snow Conference*
- 716 Tveit J and Killingtveit Å (1994) Snow surveys for studies of water budget on Svalbard 1991–1994. In *Proceedings*
717 *of the 10th International Northern Research Basins Symposium and Workshop, Spitsbergen, Norway. SINTEF*
718 *Report*, volume 22, A96415

- 719 Waechter A, Copland L and Herdes E (2015) Modern glacier velocities across the Icefield Ranges, St Elias
720 Mountains, and variability at selected glaciers from 1959 to 2012. *Journal of Glaciology*, **61**(228), 624–634 (doi:
721 10.3189/2015JoG14J147)
- 722 Walmsley APU (2015) *Long-term observations of snow spatial distributions at Hellstugubreen and Gråsubreen,*
723 *Norway*. Master’s thesis, University of Oslo
- 724 Wetlaufer K, Hendrikx J and Marshall L (2016) Spatial Heterogeneity of Snow Density and Its Influence on Snow Wa-
725 ter Equivalence Estimates in a Large Mountainous Basin. *Hydrology*, **3**(3), 1–17 (doi: 10.3390/hydrology3010003)
- 726 Wilson N and Flowers G (2013) Environmental controls on the thermal structure of alpine glaciers. *The Cryosphere*,
727 **7**(1), 167–182 (doi: 10.5194/tc-7-167-2013)
- 728 Wilson NJ, Flowers GE and Mingo L (2013) Comparison of thermal structure and evolution between neighboring
729 subarctic glaciers. *Journal of Geophysical Research: Earth Surface*, **118**(3), 1443–1459 (doi: 10.1002/jgrf.20096)
- 730 Winstral A, Elder K and Davis RE (2002) Spatial snow modeling of wind-redistributed snow using terrain-based pa-
731 rameters. *Journal of Hydrometeorology*, **3**(5), 524–538 (doi: 10.1175/1525-7541(2002)0030524:SSMOWR2.0.CO;2)
- 732 Winstral A, Marks D and Gurney R (2013) Simulating wind-affected snow accumulations at catchment to basin
733 scales. *Advances in Water Resources*, **55**, 64–79 (doi: 10.1016/j.advwatres.2012.08.011)
- 734 Winther J, Bruland O, Sand K, Killingtveit A and Marechal D (1998) Snow accumulation distribution on Spitsbergen,
735 Svalbard, in 1997. *Polar Research*, **17**, 155–164 (doi: 10.3402/polar.v17i2.6616)
- 736 Woo MK and Marsh P (1978) Analysis of Error in the Determination of Snow Storage for Small High Arctic Basins.
737 *Journal of Applied Meteorology*, **17**(10), 1537–1541 (doi: 10.1175/1520-0450(1978)0171537:AOEITD2.0.CO;2)
- 738 Wood WA (1948) Project “Snow Cornice”: the establishment of the Seward Glacial research station. *Arctic*, **1**(2),
739 107–112
- 740 Work R, Stockwell H, Freeman T and Beaumont R (1965) *Accuracy of field snow surveys*. Cold Regions Research &
741 Engineering Laboratory
- 742 Zhang W and Montgomery DR (1994) Digital elevation model grid size, landscape representation, and hydrologic
743 simulations. *Water Resources Research*, **30**(4), 1019–1028 (doi: 10.1029/93WR03553)