

PROBLEM SET 01

DAY 1

The hardest step in writing an $\epsilon - \delta$ proof is the first step, once you know how to start the proof the rest is mechanical.

- Q.1.** Explain in your own words how the provisional definition is equivalent to the more rigorous $\epsilon - \delta$ definition of limit.

Provisional definition: The function f approaches a limit l near a i.e. $\lim_{x \rightarrow a} f(x) = l$, if we can make $f(x)$ as close to l as we like by requiring that x be sufficiently close to, but unequal to, a .

$\epsilon - \delta$ **definition:** The function f approaches a limit l near a i.e. $\lim_{x \rightarrow a} f(x) = l$, if for every $\epsilon > 0$ there is some $\delta > 0$ such that, for all x , if $0 < |x - a| < \delta$, then $|f(x) - l| < \epsilon$.

Explain how this is equivalent to the standard $\epsilon - \delta$ definition.

- Q.2.** For each of the following functions f and real numbers a ,

- Guess the limit $L = \lim_{x \rightarrow a} f(x)$.
- Find a δ corresponding to $\epsilon = 0.1$ in the $\epsilon - \delta$ definition of limit.
- Find a δ corresponding to an arbitrary real number ϵ and **prove** that L is indeed the limit.

(1) $f(x) = x + 2, \quad a = 0$

(2) $f(x) = 7x + 2, \quad a = 0$

(3) $f(x) = x^2 - 2, \quad a = 0$

(4) $f(x) = 1/x, \quad a = 1$

DAY 2

It's tedious to use $\epsilon - \delta$ proofs in practice, instead we use Theorem 2 and its analogues.

- Q.3.**
- Let n be a positive integer. Prove that the function $f(x) = x^n$ is continuous at 0.
 - Use Theorem 2 to prove that every polynomial $p(x)$ is continuous at 0.
 - Prove that a function $f(x)$ is continuous at a if and only if the function $g(x) = f(x + a)$ is continuous at 0.
 - Prove that x^n is continuous at every real number a .
 - Use Theorem 2 to prove that every polynomial $p(x)$ is continuous at all real numbers.
 - What are the real numbers at which the ratio of two polynomials $\frac{p(x)}{q(x)}$ is continuous?

- Q.4.** Using the $\epsilon - \delta$ definition of limit to prove that if $\lim_{x \rightarrow a} f(x) = l$ and $\lim_{x \rightarrow a} g(x) = m$ then

$$\lim_{x \rightarrow a} (f(x) - g(x)) = l - m$$

Trigonometric functions, exponential functions and logarithms are continuous wherever they are defined. We will assume this fact without proof for now and come back to it later.

DAY 3

To understand continuity it is equally important to understand discontinuity.

- Q.5.**
- Determine, with proof, $\lim_{x \rightarrow \infty} 1/x$.
 - Prove that for no real number l do we have $\lim_{x \rightarrow 0^+} 1/x = l$.

- Q.6.** For the function

$$f(x) = \begin{cases} 0 & \text{if } x < 0 \\ 1 & \text{otherwise} \end{cases}$$

determine, with proof, the limits $\lim_{x \rightarrow 0^+}$, $\lim_{x \rightarrow 0^-}$. How will your answers change if we replace $x < 0$ by $x \leq 0$ in the definition of $f(x)$?

- Q.7.** Prove that the function

$$f(x) = \begin{cases} 0 & \text{if } x \text{ is rational} \\ 1 & \text{otherwise} \end{cases}$$

is discontinuous everywhere. What can you say about the continuity of the function $g(x) = x \cdot f(x)$?

- Q.8.** Give examples to show that the following definitions of $\lim_{x \rightarrow a} f(x) = L$ are not correct. (Hint: Think graphically)

- For all $\delta > 0$ there exists an $\epsilon > 0$ such that if $0 < |x - a| < \delta$ then $|f(x) - L| < \epsilon$.
- For every $\epsilon > 0$ there is some $\delta > 0$ such that, for all x , if $|f(x) - L| < \epsilon$ then $0 < |x - a| < \delta$.