

# **Computer Arithmetic and Errors**

While numerical result is an approximation, this can usually be as accurate as needed.

The necessary accuracy is, of course, determined by the application. To achieve high accuracy, very many separate operations must be carried out, but computers do them so rapidly without ever making mistakes that is no significant problem.

Actually, evaluating an analytical result to get the numerical answer for a specific application is subject to the same errors.

How can we define 'error' in a computation? In its simplest form, it is the difference between the exact answer **True**, say, and the computed answer, **Approximate**.

Hence, we can write,

$$ERROR = True - Approximate$$

Since we are usually interested in the magnitude or absolute value of the error we can also define

$$ABSOLUTE\_ERROR = |True - Approximate|$$

In practical calculations, it is important to obtain an upper bound on the error i.e. a number, **Upper**, such that,

$$|True - Approximate| < Upper(TOLERANCE)$$

Clearly, we would like **Upper** to be small!

In practice we are often more interested in so-called '**relative error**' rather than absolute error and we define,

$$RELATIVE\_ERROR = \frac{|True - Approximate|}{|True|}$$

**The percentage error** is 100% times the relative error.

Hence, an 'error' of  $10^{-5}$  may be a good or bad 'relative error' depending on the answer.

For example,

Answer = 1000	error = $10^{-5}$	very good
Answer = 1	error = $10^{-5}$	good
Answer = $10^{-5}$	error = $10^{-5}$	very bad

**Example:** Find the absolute error and relative error in the following cases.

- Let  $y = 3.141592$ (True) and  $\hat{y} = 3.14$ (Approximate);

Then the absolute error is

$$E_y = |3.141592 - 3.14| = 0.001592$$

and the relative error is

$$R_y = \frac{0.001592}{3.141592} = 0.000507.$$

- Let  $w = 1000000$  and  $\hat{w} = 999996$ ; then the absolute error is

$$E_w = |1000000 - 999996| = 4$$

and the relative error is

$$R_w = \frac{4}{1000000} = 0.000004.$$

- Let  $s = 0.000012$  and  $\hat{s} = 0.000009$ ; then the absolute error is

$$E_s = |0.000012 - 0.000009| = 0.000003$$

and the relative error is

$$R_s = \frac{0.000003}{0.000012} = 0.25$$




# Numerical Integration

## Simple Functions that do not have simple antiderivatives

The normal distribution is a very important functions in statistics. Gaussian noise is one of many ways in which this function is used in engineering and science. The normal distribution function is scaled form of the function  $f(x) = e^{-x^2}$ . The indefinite integral of this function **cannot be represented as a simple function.**

```
> int(exp(-x^2), x);
```

$$\frac{1}{2}\sqrt{\pi} \operatorname{erf}(x)$$




Another function that is important in optics and other applications, but does not have a simple antiderivative is,

$$f(x) = \sin(x)/x.$$


```
> int(sin(x)/x,x);
```

Si(x)







An Approximation of the Integral of  
 $f(x) = \sin(x)$   
on the Interval  $[0, 5.0]$   
Using a Midpoint Riemann Sum  
Approximate Value: .7163378145




An Approximation of the Integral of  
 $f(x) = \sin(x)$   
on the Interval  $[0, 5]$   
Using the Trapezoid Rule  
Approximate Value: .7163378145



An Approximation of the Integral of  
 $f(x) = x^*(x-2)*(x-3)$   
on the Interval  $[0, 5]$   
Using the Trapezoid Rule  
Approximate Value: 22.91666667



**Theorem:** *If a function  $f(x)$  is continuous on a finite interval  $a \leq x \leq b$ , then the definite integral of  $f(x)$  with respect to  $x$  from  $x=a$  to  $x=b$  exists.*




**Theorem:(The Fundamental Theorem of Calculus)** *If  $f(x)$  is continuous on the interval  $a \leq x \leq b$ , and  $F(x)$  is an antiderivative of  $f(x)$  then*

$$\int_a^b f(x)dx = F(b) - F(a)$$

**Evaluate**


$$\int_0^2 x^2 dx$$



$$\int_0^2 x^2 dx = \left. \frac{x^3}{3} \right|_0^2 = \frac{8}{3}$$

**Numerical integration rules are very important because even simple functions may not have exact formulas for their antiderivatives (indefinite integrals). Even when an exact formula for the antiderivative does exist, it may be difficult to find.**

$$\int_0^2 e^{x^2} dx = ?$$



$$y = e^{x^2}, \text{ and } y = x^2$$

$$\int_0^2 e^{x^2} dx = -\frac{1}{2} \operatorname{erf}(2I) \sqrt{\pi}$$

> `int(exp(x^2),x=0..2);`

$$-\frac{1}{2} \operatorname{erf}(2I) \sqrt{\pi}$$

In general, a numerical integration formula approximates a definite integral by a weighted sum of function values at points within the interval of integration has the form

$$\int_a^b f(x)dx \approx \sum_{i=1}^n c_i f(x_i)$$

where the coefficients  $c_i$  depend on the particular method.

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x_i$$

$$\Delta x_i = x_i - x_{i-1}$$

We consider the most common numerical integration formulas that are based on equally spaced data points : these are known as *Newton-Cotes formulas*.

Subdivide the interval into  $n$  equal subintervals by  $n+1$  points.

$$a = x_0 < x_1 < x_2 < \dots < x_i \dots < x_{n-1} < x_n = b$$

**Evaluate**

$$\int_0^2 x^2 dx$$

$$\Delta x = \frac{b-a}{n}$$

$$\Delta x = \frac{2-0}{n} = \frac{2}{n}$$

$$x_i = a + i\Delta x = 0 + i\frac{2}{n} = \frac{2i}{n}$$

$$\begin{aligned}\sum_{i=1}^n f(x_i)\Delta x &= \sum_{i=1}^n f\left(\frac{2i}{n}\right)\left(\frac{2}{n}\right) \\&= \sum_{i=1}^n \left(\frac{2i}{n}\right)^2 \left(\frac{2}{n}\right) \\&= \frac{8}{n^3} \sum_{i=1}^n i^2 \\&= \left(\frac{8}{n^3}\right) \left(\frac{n(n+1)(2n+1)}{6}\right) = \frac{4(n+1)(2n+1)}{3n^2}\end{aligned}$$

$$\int_0^2 x^2 dx = \lim_{n \rightarrow \infty} \frac{4(n+1)(2n+1)}{3n^2} = \frac{8}{3}$$

```
> int(x^2,x=0..2);
```

$$\frac{8}{3}$$

**Evaluate**


$$\int_1^3 (x^3 - 2x^2 + 1)dx$$

$$\Delta x = \frac{3-1}{n} = \frac{2}{n}$$

$$x_i = a + i\Delta x = 1 + i\frac{2}{n} = 1 + \frac{2i}{n}$$

$$\begin{aligned}\sum_{i=1}^n f(x_i)\Delta x &= \sum_{i=1}^n f\left(1 + \frac{2i}{n}\right)\left(\frac{2}{n}\right) \\&= \sum_{i=1}^n \left\{ \left(1 + \frac{2i}{n}\right)^3 - 2\left(1 + \frac{2i}{n}\right)^2 + 1 \right\} \left(\frac{2}{n}\right) \\&= \frac{16}{n^4} \sum_{i=1}^n i^3 + \frac{8}{n^3} \sum_{i=1}^n i^2 - \frac{4}{n^2} \sum_{i=1}^n i \\&= \left(\frac{16}{n^4}\right) \left(\frac{n^2(n+1)^2}{4}\right) + \left(\frac{8}{n^3}\right) \left(\frac{n(n+1)(2n+1)}{6}\right) - \left(\frac{4}{n^2}\right) \left(\frac{n(n+1)}{2}\right)\end{aligned}$$

$$\begin{aligned}\int_1^3 (x^3 - 2x^2 + 1)dx &= \lim_{n \rightarrow \infty} \left\{ \frac{4(n+1)^2}{n^2} + \frac{4(n+1)(2n+1)}{3n^2} - \frac{2(n+1)}{n} \right\} \\&= 4 + \frac{8}{3} - 2 = \frac{14}{3}\end{aligned}$$



> `int(x^3-2*x^2+1,x=1..3);`



$$\frac{14}{3}$$

# Riemann Sum

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i)\Delta x$$

where

$$\Delta x = \frac{b-a}{n}$$





This method of integral approximation is known as a **Riemann sum**. There are **3 basic types** of Riemann sums:

*1) The left or lower sum*


$$\int_a^b f(x)dx \approx \sum_{i=0}^{n-1} f(x_i)(x_{i+1} - x_i)$$

*2) The right or upper sum*


$$\int_a^b f(x)dx \approx \sum_{i=1}^n f(x_i)(x_i - x_{i-1})$$

*3) The middle or midpoint sum*


$$\int_a^b f(x)dx \approx \sum_{i=1}^n f\left(\frac{x_i + x_{i-1}}{2}\right)(x_i - x_{i-1})$$



**LOWER**




**UPPER**



**MIDPOINT**

```
> with(Student[Calculus1]):
> RiemannSum(x^2, x=0.0..2.0, method = lower, output=plot, partition=4);
```




$$\int_0^2 x^2 dx \approx \sum_{i=0}^{n-1} f(x_i)(x_{i+1} - x_i)$$

$$= 0^2(0.5 - 0) + 0.5^2(1 - 0.5) + 1^2(1.5 - 1) + 1.5^2(2 - 1.5)$$

$$= 1.75$$

```
> with(Student[Calculus1]):
> RiemannSum(x^2, x=0.0..2.0, method = upper,output=plot,partition=4);
```



$$\int_0^2 x^2 dx \approx \sum_{i=1}^n f(x_i)(x_i - x_{i-1})$$

$$= 0.5^2(0.5 - 0) + 1^2(1 - 0.5) + 1.5^2(1.5 - 1) + 2^2(2 - 1.5)$$

$$= 3.75$$


```
> with(Student[Calculus1]):
> RiemannSum(x^2, x=0.0..2.0, method = midpoint, output=plot, partition=4);
```

An Approximation of the Integral of

$$f(x) = x^2$$


on the Interval  $[0, 2.0]$

Using a Midpoint Riemann Sum




$$\begin{aligned} \int_0^2 x^2 dx &\approx \sum_{i=1}^n f\left(\frac{x_i + x_{i-1}}{2}\right)(x_i - x_{i-1}) \\ &= 0.25^2(0.5 - 0) + 0.75^2(1 - 0.5) \\ &\quad + 1.25^2(1.5 - 1) + 1.75^2(2 - 1.5) \\ &= 0.5(0.25^2 + 0.75^2 + 1.25^2 + 1.75^2) = 2.625 \end{aligned}$$

# Lower Riemann Sum (Partitions=40)



**Area=2.56750000**

# Upper Riemann Sum (Partitions=40)



**Area=2.76750000**


# Midpoint Riemann Sum (Partitions=40)

An Approximation of the Integral of

$$f(x) = x^2$$

on the Interval  $[0, 2.0]$

Using a Midpoint Riemann Sum



**Area=2.66625000**




# Trapezoidal Rule

One of the simplest ways to approximate the area under a curve is to approximate the curve by a straight line. The trapezoidal rule approximates the curve by the straight line that passes through the points  $(a, f(a))$  and  $(b, f(b))$ , the two ends of the interval of interest.

We have  $x_0 = a$ ,  $x_1 = b$  and  $h = b - a$

$$\int_a^b f(x)dx \approx \frac{h}{2}[f(x_0) + f(x_1)]$$

$$\int_a^b f(x)dx \approx \frac{b-a}{2n}[f(a) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(b)]$$



The area of the  $i_{th}$  trapezoid is given by

$$\Delta x \left( \frac{f(x_i) + f(x_{i-1})}{2} \right)$$


The sum of the areas of the trapezoids reduces to a simple formula,

$$\int_a^b f(x) dx \approx \frac{b-a}{2n} [f(a) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(b)]$$

**Example:** Evaluate  $\int_0^2 x^2 dx$  using Trapezoidal Rule with 4 and 10 partitions.

**Partitions:4 (n=4)**


> `ApproximateInt(x^2, x=0..2, method = trapezoid,output=plot,partition=4);`



$$\int_a^b f(x) dx \approx \frac{b-a}{2n} [f(a) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(b)]$$

$$\begin{aligned} \int_0^2 x^2 dx &\approx \frac{2-0}{2(4)} [0 + 2(0.5^2) + 2(1^2) + 2(1.5^2) + (2^2)] \\ &= 0.25 [0 + 2(0.5^2) + 2(1^2) + 2(1.5^2) + (2^2)] = 2.75 \end{aligned}$$

## Partitions:10



$$\begin{aligned}\int_0^2 x^2 dx &\approx \frac{2-0}{2(10)} \left[ 0 + 2(0.2^2 + 0.4^2 + 0.6^2 + 0.8^2 + 1^2 + \right. \\ &\quad \left. 1.2^2 + 1.4^2 + 1.6^2 + 1.8^2) + (2^2) \right] \\ &= 0.1[0 + 2(11.4) + (4)] = 0.1[26.8] = 2.68\end{aligned}$$

## Error Estimate For The Trapezoidal Rule

Let  $T_n$  be difference between  $\int_a^b f(x)dx$  and trapezoid rule estimate for  $\int_a^b f(x)dx$  with  $n$  intervals. Then  $T_n$  is the estimation error. If  $M_2$  is the maximum value of  $|f''(x)|$  on  $[a,b]$ , then


$$|T_n| \leq \frac{M_2(b-a)^3}{12n^2}$$

**This estimate is valid only when  $f''(x)$  is defined on all of  $[a,b]$ .**

**Example:** Estimate  $\int_0^1 x^2 dx$  using the trapezoidal rule with  $n=5$  partitions (intervals):

$$\Delta x = \frac{1-0}{5} = \frac{1}{5}$$

$$x_0 = 0, x_1 = 1/5, \dots, x_5 = 1$$



$$\int_0^1 x^2 dx \approx \frac{1}{2(5)} [0 + 2(1/25) + 2(4/25) + 2(9/25) + 2(16/25) + 1] = 0.34$$

**Example:**

Find a bound on the error in estimating  $\int_0^2 x^2 dx$  using the trapezoidal rule with 5 partitions.

$f''(x) = 2$  So the maximum value of  $|f''(x)|$  on  $[0, 2]$  is  $M_2 = 2$ .

Thus

$$|T_5| \leq \frac{2(2-0)^3}{12 * 5^2} = \frac{16}{300} = 0.05333$$

$$\int_0^2 x^2 dx \approx \frac{1}{2(5)} [0 + 2(0.16) + 2(0.64) + 2(1.44) + 2(2.56) + 4] = 2.72$$

True Value:  $8/3 = 2.666$       Estimate = 2.72

$$\text{Absolute\_Error} = |2.666 - 2.72| = 0.05333$$

## Matlab (Trapezoidal Rule)

```
function Q = Trap( f, a, b, n)
h = (b-a)/n;
S = feval(f, a);
if n == 1
    Q = (S + feval(f, b))*h/2;
else
    for i = 1 : n-1
        x(i) = a + h*i;
        S = S + 2*feval(f, x(i));
    end
    S = S + feval(f, b);
    Q = h*S/2;
end
```

```
f=inline('x^2')
```

```
f =
```

```
    Inline function:
```

```
    f(x) = x^2
```

```
>> Q=Trap(f,0,2,5)
```

```
Q =
```

```
    2.7200
```


```
>> Q=Trap(f,0,2,10)
```

```
Q =
```

```
    2.6800
```



# Simpson's Rule



\*Partition

**The area under the parabola may be written in the form**

$$\int_{x_{i-1}}^{x_{i+1}} (ax^2 + bx + c)dx = \frac{\Delta x}{3} [f(x_{i-1}) + 4f(x_i) + f(x_{i+1})]$$

**and this expression is free of a,b and c.**

**Consider the problem of finding the integral of**  $\int_0^2 e^{-x^2} dx$

**The required values for applying Simpson's rule are**

$$x_0 = a = 0, x_1 = (b - a)/2 = (2 - 0)/2 = 1, x_2 = b = 2$$

**which**

**gives**

$$\int_0^2 e^{-x^2} dx \approx \frac{1}{3} [\exp(-0^2) + 4 \exp(-1^2) + \exp(-2^2)] = 0.8299$$

## Discussion

Simpson's rule is found by integrating the Lagrange interpolating polynomial for  $f(x)$

$$L(x) = \frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)} f(x_0) + \frac{(x-x_0)(x-x_2)}{(x_1-x_0)(x_2-x_2)} f(x_1) + \frac{(x-x_0)(x-x_1)}{(x_2-x_0)(x_2-x_1)} f(x_2)$$

$$x_0 = a, x_1 = x_0 + h, x_2 = b, h = \frac{b-a}{2}$$

$$\int_a^b f(x)dx \approx \int_a^b L(x)dx = \frac{h}{3} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right]$$

## In general

$$\int_a^b f(x)dx \approx \frac{(b-a)}{3n} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]$$


**n must be even number for Simpson's rule.**

**Example:** Estimate  $\int_1^3 \frac{1}{x} dx$  using the Simpson's rule with  $n=4$  intervals.

$$\Delta x = \frac{3-1}{4} = \frac{1}{2}$$

$$x_0 = 1, x_1 = 3/2, \dots, x_4 = 3$$

$$\int_1^3 \frac{1}{x} dx \approx \frac{3-1}{3(4)} [1 + 4(2/3) + 2(1/2) + 4(2/5) + 1/3] = 1.1$$



$$\int_1^3 \frac{1}{x} dx = \left| \ln x \right|_1^3 = \ln(3) = 1.09861..$$

## Simpson's Rule Error Estimate

$$|S_n| \leq \frac{M_4(b-a)^5}{180n^4}$$

$M_4$  is the maximum value of  $|f^{(4)}(x)|$  on  $[a,b]$

Find a bound on the error in estimating  $\int_1^3 \frac{1}{x} dx$  using the Simpson's rule with  $n=4$  intervals.

$f^{(4)}(x) = 24x^{-5}$  So the maximum value of  $|f^{(4)}(x)|$  on  $[1,3]$  is  $M_4=24$ .

Thus

$$|S_4| \leq \frac{24(3-1)^5}{180(4)^2} = 0.01666$$

Exact error:  $1.1 - \ln(3) = 0.00139$

## Matlab (Simpson Rule)

```
%metodod de simpson
function Q = Simp( f, a, b, n)
h = (b-a)/n;
S = feval(f, a);
if rem(n,2) != 0
    error('n must be even for Simpson method')
    return
end
if n == 2
    Q = (S + 4*feval(f,a+h) + feval(f, b))*h/3;
else
    for i = 1 : 2 : n-1
        x(i) = a + h*i;
        S = S + 4*feval(f, x(i));
    end
    for i = 2 : 2 : n-2
        x(i) = a + h*i;
        S = S + 2*feval(f, x(i));
    end
    S = S + feval(f, b);
    Q = h*S/3;
end
```

$$\int_1^3 \frac{1}{x} dx \approx \frac{3-1}{3(4)} [1 + 4(2/3) + 2(1/2) + 4(2/5) + 1/3] = 1.1$$

f=inline('1/x')

f =

Inline function:

f(x) = 1/x

>> Q=Simp(f,1,3,4)

Q =

1.1000

## Length of a Curve

*The arc length of the  $f(x)$  over the interval  $a \leq x \leq b$  is,*

$$Length = \int_a^b \sqrt{1 + (f'(x))^2} dx$$

## Surface Area

*The solid of revolution obtained by rotating the region under the curve  $f(x)$ , where  $a \leq x \leq b$ , about the axis has surface area given by*

$$SurfaceArea = 2\pi \int_a^b f(x) \sqrt{1 + (f'(x))^2} dx$$

## Other Methods

Romberg Integration  
Gaussian quadrature