



# Newton Polynomials Divided Differences



There are two **disadvantages** to using the Lagrange Polynomial (or Neville's method) for interpolation.

- **First**, they involve more **arithmetic operations** than does the divided difference method.
- **Second** and more importantly, if we desire **to add or subtract a point from the set** used to construct the polynomial, we essentially to start over in the computations.
- Both the Lagrange polynomials and Neville's method also must repeat all of the arithmetic if we must interpolate at a new x-value.
- **The divided difference method avoids all of this computation.**

**Our treatment of divided-difference tables assumes that the function,  $f(x)$ , is known at several values for  $x$ :**

$x_0$	$f_0$
$x_1$	$f_1$
$x_2$	$f_2$
$\cdot$	$\cdot$
$x_n$	$f_n$

**We do not assume that the  $x$ 's are evenly spaced or even that the values are arranged in any particular order (but some ordering may be advantageous).**

## Straight Line

**Lagrange** linear interpolation polynomial

$$\begin{aligned} P_1(x) &= y_0 \frac{(x - x_1)}{(x_0 - x_1)} + y_1 \frac{(x - x_0)}{(x_1 - x_0)} \\ &= y_0 L_{1,0}(x) + y_1 L_{1,1}(x) \end{aligned}$$

$$L_{1,0}(x) = \frac{(x - x_1)}{(x_0 - x_1)} \quad \text{and} \quad L_{1,1}(x) = \frac{(x - x_0)}{(x_1 - x_0)}.$$

**The Newton form** of the equation of a straight line passing through two points  $(x_0, y_0)$  and  $(x_1, y_1)$


$$P_1(x) = a_0 + a_1(x - x_0)$$

$$a_0 = y_0, \quad a_1 = \frac{y_1 - y_0}{x_1 - x_0}$$

### Example: (Comparison)

$$(x_0, y_0) : (1, 2)$$

$$(x_1, y_1) : (2, 3)$$




## Lagrange Interpolation

$$\begin{aligned} P_1(x) &= y_0 \frac{(x - x_1)}{(x_0 - x_1)} + y_1 \frac{(x - x_0)}{(x_1 - x_0)} \\ &= 2 \frac{(x - 2)}{(1 - 2)} + 3 \frac{(x - 1)}{(2 - 1)} \\ &= -2(x - 2) + 3(x - 1) \\ &= -2x + 4 + 3x - 3 = x + 1 \end{aligned}$$

## Newton Interpolation

$$\begin{aligned} P_1(x) &= a_0 + a_1(x - x_0) \\ &= y_0 + \frac{(y_1 - y_0)}{(x_1 - x_0)}(x - x_0) \\ &= 2 + \frac{(3 - 2)}{(2 - 1)}(x - 1) \\ &= 2 + (x - 1) = x + 1 \end{aligned}$$



# Newton Interpolation Parabola

$$P_2(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1)$$

$$a_0 = y_0, a_1 = \frac{y_1 - y_0}{x_1 - x_0},$$

$$a_2 = \frac{\frac{y_2 - y_1}{x_2 - x_1} - \frac{y_1 - y_0}{x_1 - x_0}}{x_2 - x_0}$$

## Compare with the Newton Linear Interpolation

$$P_1(x) = a_0 + a_1(x - x_0)$$

$$a_0 = y_0, a_1 = \frac{y_1 - y_0}{x_1 - x_0}$$

# Lagrange Interpolation Parabola

$$\begin{aligned} P_2(x) &= y_0 \frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)} + \\ &\quad y_1 \frac{(x-x_0)(x-x_2)}{(x_1-x_0)(x_1-x_2)} + \\ &\quad y_2 \frac{(x-x_0)(x-x_1)}{(x_2-x_0)(x_2-x_1)} \\ &= y_0 L_{2,0}(x) + y_1 L_{2,1}(x) + y_2 L_{2,2}(x) \end{aligned}$$

## Compare with the Lagrange Linear Interpolation


$$\begin{aligned} P_1(x) &= y_0 \frac{(x-x_1)}{(x_0-x_1)} + y_1 \frac{(x-x_0)}{(x_1-x_0)} \\ &= y_0 L_{1,0}(x) + y_1 L_{1,1}(x) \end{aligned}$$



**Example:** Suppose that we have the following data pairs – x values and f(x) values- unknown function.

x	y
-2	4
0	2
2	8



Find a quadratic polynomial using the three given points.



## This problem solved before using Lagrange Polynomial approximation

$$\begin{aligned} p(x) &= \frac{(x-0)(x-2)}{(-2-0)(-2-2)}(4) + \frac{(x-(-2))(x-2)}{(0-(-2))(0-2)}(2) + \frac{(x-(-2))(x-0)}{(2-(-2))(2-0)}(8) \\ &= \frac{x(x-2)}{8}(4) + \frac{(x+2)(x-2)}{-4}(2) + \frac{x(x+2)}{8}(8) \\ &= x^2 + x + 2 \end{aligned}$$

>> plot(x,y,'bo');grid on



$$p(x) = x^2 + x + 2$$

## Newton Interpolation Parabola

$$P_2(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1)$$

$$a_0 = y_0, a_1 = \frac{y_1 - y_0}{x_1 - x_0},$$

$$a_2 = \frac{\frac{y_2 - y_1}{x_2 - x_1} - \frac{y_1 - y_0}{x_1 - x_0}}{x_2 - x_0}$$

$$(x_0, y_0) = (-2, 4), (x_1, y_1) = (0, 2), (x_2, y_2) = (2, 8)$$

$$a_0 = y_0 = 4,$$

$$a_1 = \frac{y_1 - y_0}{x_1 - x_0} = \frac{2 - 4}{0 - (-2)} = -1,$$

$$a_2 = \frac{\frac{y_2 - y_1}{x_2 - x_1} - \frac{y_1 - y_0}{x_1 - x_0}}{x_2 - x_0} = \frac{\frac{8 - 2}{2 - 0} - \frac{2 - 4}{0 - (-2)}}{2 - (-2)} = 1$$

**Thus**

$$p(x) = 4 - (x + 2) + x(x + 2) = x^2 + x + 2$$

**Check this result with the result of Lagrange Polynomial.**

**The calculations can be performed in a systematic manner, using a “divided-difference table”.**

$x_i$	$y_i$	$d_i$	$dd_i$
-2	4		
		$\frac{y_1 - y_0}{x_1 - x_0} = \frac{2 - 4}{0 - (-2)} = -1$	
0	2		$\frac{d_2 - d_1}{x_2 - x_0} = \frac{3 - (-1)}{2 - (-2)} = 1$
		$\frac{y_2 - y_1}{x_2 - x_1} = \frac{8 - 2}{2 - 0} = 3,$	
2	8		

**The coefficients of the Newton polynomial are the top entries in this table.**

$$P_2(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1)$$

$$\begin{aligned} p(x) &= 4 + (-1)(x + 2) + (1)x(x + 2) \\ &= x^2 + x + 2 \end{aligned}$$

## Additional Data Points


Extend the previous example adding two new points.

**Old table**

x	y
-2	4
0	2
2	8

**New Table**

x	y
-2	4
0	2
2	8
-1	-1
1	1




The divided-difference table becomes

$x_i$	$y_i$	$d_i$	$dd_i$	$ddd_i$	$dddd_i$
-2	4	$\frac{(2-4)}{0-(-2)} = -1$			
0	2	$\frac{(8-2)}{(2-0)} = 3$	$\frac{(3+1)}{(2+2)} = 1$	$\frac{(0-1)}{(-1+2)} = -1$	
2	8	$\frac{(-1-8)}{(-1-2)} = 3$	$\frac{(3-3)}{(-1-0)} = 0$	$\frac{(2-0)}{(1-0)} = 2$	$\frac{(2+1)}{(1+2)} = 1$
-1	-1	$\frac{(1+1)}{(1+1)} = 1$	$\frac{(1-3)}{(1-2)} = 2$		
1	1				

## The Newton interpolation polynomial

$$P_4(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + a_3(x - x_0)(x - x_1)(x - x_2) + a_4(x - x_0)(x - x_1)(x - x_2)(x - x_3)$$

$$P_4(x) = 4 + (-1)(x + 2) + (1)(x + 2)(x) + (-1)(x + 2)(x)(x - 2) + (1)(x + 2)(x)(x - 2)(x + 1).$$



## Quartic interpolation polynomial

$$\begin{aligned}
 P_4(x) &= 4 + (-1)(x+2) + (1)(x+2)(x) + (-1)(x+2)(x)(x-2) + \\
 &\quad (1)(x+2)(x)(x-2)(x+1). \\
 &= x^4 - 3x^2 + x + 2.
 \end{aligned}$$

**x = -2    0    2    -1    1**

**y = 4    2    8    -1    1**

**>> P=polyfit(x,y,4)**

**P = 1.0000    0.0000    -3.0000    1.0000    2.0000**



# General Form

Consider the  $n$ th-degree polynomial written in a special way:

$$P_n(x) = a_0 + (x - x_0)a_1 + (x - x_0)(x - x_1)a_2 + \dots + (x - x_0)(x - x_1)\dots(x - x_{n-1})a_n.$$

If we choose the  $a_i$  so that  $P_n(x) = f(x)$  at the  $n + 1$  known points,  $(x_i, f_i), i = 0, \dots, n$  then  $P_n(x)$  is an interpolating polynomial. We will show that the  $a_i$ 's are readily **determined by using what are called the divided difference of the tabulated values.**

## Standard notation for divided difference

A special standard notation for divided differences is

$$f[x_0, x_1] = \frac{f_1 - f_0}{x_1 - x_0},$$

called the *first divided difference between  $x_0$  and  $x_1$* .

The function

$$f[x_1, x_2] = \frac{f_2 - f_1}{x_2 - x_1},$$

is the *first divided difference between  $x_1$  and  $x_2$* .

(We use  $f[x_0] = f_0 = f(x_0)$  )

In general,

$$f[x_s, x_t] = \frac{f_t - f_s}{x_t - x_s},$$

is the *first divided difference between  $x_s$  and  $x_r$* .

Observe that

$$f[x_s, x_t] = \frac{f_t - f_s}{x_t - x_s} = \frac{f_s - f_t}{x_s - x_t} = f[x_t, x_s]$$

## Second Order Differences

Second and higher order differences are defined in terms of lower-order differences.

**Second order:**

$$f[x_0, x_1, x_2] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0}$$

**n<sup>th</sup> - order:**

$$f[x_0, x_1, \dots, x_n] = \frac{f[x_1, x_2, \dots, x_n] - f[x_0, x_1, \dots, x_{n-1}]}{x_n - x_0}$$

**Zero order difference:**

$$f[x_s] = f_s$$

$x_0$      $f[x_0]$

$$f[x_0, x_1] = \frac{f[x_1] - f[x_0]}{x_1 - x_0}$$

$x_1$      $f[x_1]$

$$f[x_1, x_2] = \frac{f[x_2] - f[x_1]}{x_2 - x_1}$$

$x_2$      $f[x_2]$

\*

$$f[x_0, x_1, x_2] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0}$$

**With this notation the polynomial**

$$P_n(x) = a_0 + (x - x_0)a_1 + (x - x_0)(x - x_1)a_2 + \dots \\ + (x - x_0)(x - x_1)\dots(x - x_{n-1})a_n.$$

**can be re-expressed**

$$P_n(x) = f[x_0] + (x - x_0)f[x_0, x_1] + (x - x_0)(x - x_1)f[x_0, x_1, x_2] + \dots \\ + (x - x_0)(x - x_1)\dots(x - x_{n-1})f[x_0, x_1, \dots, x_n].$$

$$\begin{aligned} a_0 &= f[x_0], \\ a_1 &= f[x_0, x_1], \\ a_2 &= f[x_0, x_1, x_2], \\ &\cdot \\ &\cdot \\ &\cdot \\ a_n &= f[x_0, x_1, \dots, x_n]. \end{aligned}$$

$$P_n(x) = f[x_0] + \sum_{k=1}^n f[x_0, x_1, \dots, x_k](x - x_0)\dots(x - x_{k-1}).$$

**This is known as Newton's interpolatory divided-difference Formula.**

**Example:** Suppose that we have the following data pairs –  $x$  values and  $f(x)$  values- where  $f(x)$  is some unknown function.

<b>i</b>	<b>x</b>	<b>f(x)</b>
<b>0</b>	<b>3.2</b>	<b>22.0</b>
<b>1</b>	<b>2.7</b>	<b>17.8</b>
<b>2</b>	<b>1.0</b>	<b>14.2</b>
<b>3</b>	<b>4.8</b>	<b>38.3</b>
<b>4</b>	<b>5.6</b>	<b>51.7</b>

## Divided Differences

$x_i$	$f_i$	$f[x_i, x_{i+1}]$	$f[x_i, x_{i+1}, x_{i+2}]$	$f[x_i, \dots, x_{i+3}]$	$f[x_i, \dots, x_{i+4}]$
$x_0$	$f_0$	$f[x_0, x_1]$	$f[x_0, x_1, x_2]$	$f[x_0, x_1, x_2, x_3]$	$f[x_0, x_1, x_2, x_3, x_4]$
$x_1$	$f_1$	$f[x_1, x_2]$	$f[x_1, x_2, x_3]$	$f[x_1, x_2, x_3, x_4]$	
$x_2$	$f_2$	$f[x_2, x_3]$	$f[x_2, x_3, x_4]$		
$x_3$	$f_3$	$f[x_3, x_4]$			
$x_4$	$f_4$				

## Divided Difference Table

$x_i$	$f_i$	$f[x_i, x_{i+1}]$	$f[x_i, x_{i+1}, x_{i+2}]$	$f[x_i, \dots, x_{i+3}]$	$f[x_i, \dots, x_{i+4}]$
3.2	22.0	8.400	2.856	-0.528	0.256
2.7	17.8	2.118	2.012	0.0865	
1.0	14.2	6.342	2.263		
4.8	38.3	16.750			
5.6	51.70				

$$f[x_0, x_1] = \frac{f_1 - f_0}{x_1 - x_0} = \frac{17.8 - 22.0}{2.7 - 3.2} = \frac{-4.2}{-0.5} = 8.4$$

$$f[x_0, x_1, x_2] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0} = \frac{2.118 - 8.40}{1.0 - 3.2} = 2.856$$

$$f[x_0, x_1, x_2, x_3] = \frac{f[x_1, x_2, x_3] - f[x_0, x_1, x_2]}{x_3 - x_0} = \frac{2.012 - 2.856}{4.8 - 3.2} = -0.5275$$

$$f[x_0, \dots, x_4] = \frac{f[x_1, \dots, x_4] - f[x_0, \dots, x_3]}{x_4 - x_0} = \frac{0.0865 - (-0.528)}{5.6 - 3.2} = 0.2560$$

**Interpolating polynomial of degree-3 that fits the data at all points from  $x_0=3.2$  to  $x_3=4.8$ .**

$$P_3(x) = 22.0 + 8.4(x - 3.2) + 2.856(x - 3.2)(x - 2.7) - 0.528(x - 3.2)(x - 2.7)(x - 1.0)$$

**What is the fourth-degree polynomial that fits at all five points?**

**Interpolating polynomial of degree-4 that fits the data at all points from  $x_0=3.2$  to  $x_4=5.6$ .**

$$P_4(x) = 22.0 + 8.4(x - 3.2) + 2.856(x - 3.2)(x - 2.7) - 0.528(x - 3.2)(x - 2.7)(x - 1.0) + 0.256(x - 3.2)(x - 2.7)(x - 1.0)(x - 4.8)$$

**We only have to add one more term to  $P_3(x)$**

$$P_4(x) = P_3(x) + 0.256(x - 3.2)(x - 2.7)(x - 1.0)(x - 4.8)$$



**Find the interpolated value for  $x=3.0$  using  $P_3$**

$$P_3(x) = 22.0 + 8.4(x - 3.2) + 2.856(x - 3.2)(x - 2.7) - 0.528(x - 3.2)(x - 2.7)(x - 1.0)$$

$$**P_3(3) = 20.2120**$$

**Find the interpolated value for  $x=3.0$  using  $P_4$**

$$P_4(x) = P_3(x) + 0.256(x - 3.2)(x - 2.7)(x - 1.0)(x - 4.8)$$

$$**P_4(3) = 20.2120 + 0.256(-0.2)(0.3)(2.0)(-1.8)**  
**= 20.267296**$$

# Compare Lagrange Interpolation Polynomial

```
>> x=[3.2 2.7 1.0 4.8 5.6]  
x = 3.2000 2.7000 1.0000 4.8000 5.6000
```

```
>> y=[22.0 17.8 14.2 38.3 51.70]  
y = 22.0000 17.8000 14.2000 38.3000 51.7000
```

```
>> P=polyfit(x,y,4)  
P = 0.2558 -3.5208 18.6885 -36.1836 34.9600
```

```
>> xval=polyval(P,3.0)  
xval =
```

20.2672

Now Newton Polynomial solution was

$$\begin{aligned} P_4(x) = & 22.0 + 8.4(x - 3.2) + 2.856(x - 3.2)(x - 2.7) \\ & - 0.528(x - 3.2)(x - 2.7)(x - 1.0) \\ & + 0.256(x - 3.2)(x - 2.7)(x - 1.0)(x - 4.8) \end{aligned}$$

Simplify this equation in Maple

```
> simplify(22+8.4*(x-3.2)+2.856*(x-3.2)*(x-2.7)-0.528*(x-  
3.2)*(x-2.7)*(x-1.0)+0.256*(x-3.2)*(x-2.7)*(x-1.0)*(x-4.8));
```

$$34.97459200 - 36.20611200x + 18.70016000x^2 - 3.523200000x^3 + 0.2560000000x^4$$

## An Algorithm for Interpolation from a Divided Difference Table (Newton's Divided Difference Algorithm)

Given a set of  $n+1$  distinct numbers  $x_0, x_1, \dots, x_n$  at the number  $x$  for the function  $f$  and a value  $x=u$  at which the interpolating polynomial is to be evaluated:

We first find the coefficients of the interpolating polynomial. These are stored in vector  $dd$ .

```
For i=0 to n Step 1 Do
  Set dd[i]=f[i]
End for i.
  for j=1.. to n Step 1 Do
    set temp1=dd[j-1].
    For k=j to n Step 1 DO
      Set temp2=dd[k]
      Set dd[k]=(dd[k]-temp1)/(x[k]-x[k-j]).
      Temp1=temp2
    End for k
  End for j
```

Now compute the value of the polynomial at  $u$ . We do this by nested multiplication from the highest term.

```
Set sum = 0
For i=n Down to 1 Step 1 Do
  Set sum=(sum+dd[i])(u-x[i-1])
End for I
ddvalue=sum
ddvalue is the value of the polynomial at  $u$ ,  $P_n(u)$ .
```

## MATLAB M-File (Divided Differences Newton Interpolation Polynomial)

```
function [C,D]=newpoly(X,Y)
%Input    - X is a vector that contains a list of abscissas
%          - Y is a vector that contains a list of ordinates
%Output   - C is a vector that contains the coefficients
%           of the Newton interpolatory polynomial
%          - D is the divided difference table
n=length(X);
D=zeros(n,n);
D(:,1)=Y';
%Use formula (20) to form the divided difference table
for j=2:n
    for k=j:n
        D(k,j)=(D(k,j-1)-D(k-1,j-1))/(X(k)-X(k-j+1));
    end
end
%Determine the coefficients of the Newton interpolatory
polynomial
C=D(n,n);
for k=(n-1):-1:1
    C=conv(C,poly(X(k)));
    m=length(C);
    C(m)=C(m)+D(k,k);
End
```

### Example:

Suppose that we have the following data pairs – x values and f(x) values- where f(x) is some unknown function.

i	x	f(x)
0	3.2	22.0
1	2.7	17.8
2	1.0	14.2
3	4.8	38.3
4	5.6	51.7

```
>> X=[3.2 2.7 1.0 4.8 5.6]
```

```
X = 3.2000 2.7000 1.0000 4.8000 5.6000
```

```
>> Y=[22.0 17.8 14.2 38.3 51.7]
```

```
Y = 22.0000 17.8000 14.2000 38.3000 51.7000
```

```
>> [C,D]=newpoly(X,Y)
```

```
C =
```

```
0.2558 -3.5208 18.6885 -36.1836 34.9600
```

```
D =
```

```
22.0000 0 0 0 0
```

```
17.8000 8.4000 0 0 0
```

```
14.2000 2.1176 2.8556 0 0
```

```
38.3000 6.3421 2.0116 -0.5275 0
```

```
51.7000 16.7500 2.2626 0.0865 0.2558
```

```
>>
```


$x_i$	$f_i$	$f[x_i, x_{i+1}]$	$f[x_i, x_{i+1}, x_{i+2}]$	$f[x_i, \dots, x_{i+3}]$	$f[x_i, \dots, x_{i+4}]$
3.2	22.0	8.400	2.856	-0.528	0.256
2.7	17.8	2.118	2.012	0.0865	
1.0	14.2	6.342	2.263		
4.8	38.3	16.750			
5.6	51.70				

22.0000	0	0	0	0
17.8000	8.4000	0	0	0
14.2000	2.1176	2.8556	0	0
38.3000	6.3421	2.0116	-0.5275	0
51.7000	16.7500	2.2626	0.0865	0.2558

### Example:

Suppose that we have the following data pairs – x values and f(x) values- where f(x) is  $f(x) = x^2 e^{-x/2}$ .

i	x	f(x)
0	1.1	0.6981
1	2.0	1.4715
2	3.5	2.1287
3	5.0	2.0521
4	7.10	1.4480



$$f(x) = x^2 e^{-x/2}.$$

$x_i$	$f_i$	$f[x_i, x_{i+1}]$	$f[x_i, x_{i+1}, x_{i+2}]$	$f[x_i, \dots, x_{i+3}]$	$f[x_i, \dots, x_{i+4}]$
1.1	0.6981	0.8593	-0.1755	0.0032	0.0027
2.0	1.4715	0.4381	-0.1631	0.0191	
3.5	2.1287	-0.0511	-0.0657		
5.0	2.0521	-0.2877			
7.1	1.4480				

```
>> X=[1.10 2.0 3.5 5.0 7.1]
```

```
X = 1.1000 2.0000 3.5000 5.0000 7.1000
```

```
>> Y=[0.6981 1.4715 2.1287 2.0521 1.4480]
```

```
Y = 0.6981 1.4715 2.1287 2.0521 1.4480
```

```
>> [C,D]=newpoly(X,Y)
```

```
C = 0.0026 -0.0276 -0.0745 1.2517 -0.5558
```

```
D =
```

```
0.6981 0 0 0 0
```

```
1.4715 0.8593 0 0 0
```

```
2.1287 0.4381 -0.1755 0 0
```


```
2.0521 -0.0511 -0.1631 0.0032 0
```

```
1.4480 -0.2877 -0.0657 0.0191 0.0026
```


```
>>
```



```
> plot([0.0026*x^4-0.0276*x^3-0.0745*x^2+1.2517*x-  
0.5558],x=1.1..7.1,color=[blue]);
```



```
> plot([0.0026*x^4-0.0276*x^3-0.0745*x^2+1.2517*x-  
0.5558,(x^2)*exp(-x/2)],x=1.1..7.1,color=[red ,blue]);
```



**Find the error of interpolates for  $f(1.75)$  using polynomial of degrees 1,2, and 3.**

Degree	Interpolated Value	Actual Value	Actual Error
1.	1.25668	1.276639995	0.01966
2.	1.28520	1.276639995	-0.00856
3.	1.28611	1.276639995	-0.00947

# **Divided Differences Evenly Spaced Data (Equal Spacing)**

## Newton's interpolatory divided-difference Formula

$$P_n(x) = f[x_0] + (x - x_0)f[x_0, x_1] + (x - x_0)(x - x_1)f[x_0, x_1, x_2] + \dots + (x - x_0)(x - x_1)\dots(x - x_{n-1})f[x_0, x_1, \dots, x_n].$$

$$P_n(x) = f[x_0] + \sum_{k=1}^n f[x_0, x_1, \dots, x_k](x - x_0)\dots(x - x_{k-1})$$

**Introducing the notation**

$$h = x_{i+1} - x_i, \quad i = 0, 1, \dots, n-1$$

**And**

$$x = x_0 + sh \quad \text{or} \quad s = (x - x_0)/h$$

**The difference  $x - x_i$  can be written as**

$$x - x_i = (s - i)h;$$

**So the polynomial equation**

$$P_n(x) = f[x_0] + (x - x_0)f[x_0, x_1] + (x - x_0)(x - x_1)f[x_0, x_1, x_2] + \dots \\ + (x - x_0)(x - x_1)\dots(x - x_{n-1})f[x_0, x_1, \dots, x_n].$$

**becomes**

$$P_n(x) = P_n(x_0 + sh) = f[x_0] + shf[x_0, x_1] + s(s-1)h^2f[x_0, x_1, x_2] \\ + s(s-1)\dots(s-n+1)h^nf[x_0, x_1, \dots, x_n].$$

**In general,**

$$P_n(x) = \sum_{k=0}^n s(s-1)(s-k+1)h^k f[x_0, x_1, \dots, x_k]$$

**This formula is called the **Newton forward divided-difference formula.****

**Using binomial-coefficient notation**

$$\binom{s}{k} = \frac{s(s-1)\dots(s-k+1)}{k!}$$

**We can express  $P_n(x)$  compactly as**

$$P_n(x) = P_n(x_0 + sh) = \sum_{k=0}^n \binom{s}{k} h^k f[x_0, x_1, \dots, x_k].$$

**Example:**

**The following Table lists values of a function (the Bessel function of the first kind) at various points.**

x	f(x)
1.0	0.7651977
1.3	0.6200860
1.6	0.4554022
1.9	0.2818186
2.2	0.1103623

**We first obtain divided difference table.**

x	f(x)	First divided differences	Second divided differences	Third divided differences	Fourth divided differences
1.0	<b>0.7651977</b>	<b>-0.4837057</b>			
1.3	0.6200860	-0.5489460	<b>-0.1087339</b>		
1.6	0.4554022	-0.5786120	-0.0494433	<b>0.0658784</b>	
1.9	0.2818186	-0.5715210	0.0118183	0.0680685	<b>0.0018251</b>
2.2	0.1103623				

**The coefficients of the Newton forward divided-difference form of the interpolation polynomial are along the diagonal in the table.**

## **Non-Equal Spacing Approach:**

$$P_n(x) = f[x_0] + (x - x_0)f[x_0, x_1] + (x - x_0)(x - x_1)f[x_0, x_1, x_2] + \dots + (x - x_0)(x - x_1)\dots(x - x_{n-1})f[x_0, x_1, \dots, x_n].$$

$$\begin{aligned} P_4(x) = & 0.7651977 - 0.4837057(x-1.0) - 0.1087339(x-1.0)(x-1.3) \\ & + 0.0658784(x-1.0)(x-1.3)(x-1.6) \\ & + 0.0018251(x-1.0)(x-1.3)(x-1.6)(x-1.9) \end{aligned}$$

**$P_4(1.1) = ?$**

$$\begin{aligned} P_4(1.1) = & 0.7651977 - 0.4837057(1.1-1.0) \\ & - 0.1087339(1.1-1.0)(1.1-1.3) \\ & + 0.0658784(1.1-1.0)(1.1-1.3)(1.1-1.6) \\ & + 0.0018251(1.1-1.0)(1.1-1.3)(1.1-1.6)(1.1-1.9) \\ = & \mathbf{0.7196480} \end{aligned}$$

```
X = 1.0000 1.3000 1.6000 1.9000 2.2000
```

```
Y = 0.7652 0.6201 0.4554 0.2818 0.1104
```

```
>> [C,D]=newpoly(X,Y)
```

```
C =
```

```
0.0018 0.0553 -0.3430 0.0734 0.9777
```

```
D =
```

```
0.7652 0 0 0 0
```

```
0.6201 -0.4837 0 0 0
```

```
0.4554 -0.5489 -0.1087 0 0
```

```
0.2818 -0.5786 -0.0494 0.0659 0
```

```
0.1104 -0.5715 0.0118 0.0681 0.0018
```

**OR**

```
>> P=polyfit(X,Y,4)
```

```
P =
```

```
0.0018 0.0553 -0.3430 0.0734 0.9777
```

```
>> xval=polyval(P,1.1)
```

```
xval =
```

```
0.7196
```



**But we know the data is equally spacing.**

**So we can use**

$$P_n(x) = P_n(x_0 + sh) = f[x_0] + shf[x_0, x_1] + s(s-1)h^2 f[x_0, x_1, x_2] + s(s-1)\dots(s-n+1)h^n [x_0, x_1, \dots, x_n].$$

**If an approximation to f(1.1) is required, the reasonable choice for  $x_0, x_1, \dots, x_n$  would be**

$$x_0 = 1.0, \quad x_1 = 1.3, \quad x_2 = 1.6, \quad x_3 = 1.9, \quad x_4 = 2.2,$$

$$x = x_0 + sh \quad \text{or} \quad s = (x - x_0) / h$$

**and**

$$h=0.3,$$

$$s = (1.1-1.0)/0.3=0.1/0.3=1/3$$

$$P_4(1.1) = ?$$

$$P_4(1.1) = P_4(1.0 + 1/3(0.3)) = 0.7651977$$

$$+ (1/3) (0.3) (-0.4837057)$$

$$+ (1/3) (-2/3) (0.3)^2 (-0.1087339)$$

$$+ (1/3) (-2/3) (-5/3) (0.3)^3 (0.0658784)$$

$$+ (1/3) (-2/3) (-5/3) (-8/3) (0.3)^4 (0.0018251)$$


$$= 0.7196480$$

# **Difficulties with Polynomial Interpolation**

**There are many types of problems in which polynomial interpolation through a moderate number of data points works very poorly.**

## Example:


<b>i</b>	<b>x</b>	<b>f(x)</b>
<b>0</b>	<b>-2</b>	<b>0</b>
<b>1</b>	<b>-1.5</b>	<b>0</b>
<b>2</b>	<b>-1</b>	<b>0</b>
<b>3</b>	<b>-0.5</b>	<b>0.87</b>
<b>4</b>	<b>0</b>	<b>1</b>
<b>5</b>	<b>0.5</b>	<b>0.87</b>
<b>6</b>	<b>1</b>	<b>0</b>
<b>7</b>	<b>1.5</b>	<b>0</b>
<b>8</b>	<b>2</b>	<b>0</b>




?

## Example:

<b>i</b>	<b>x</b>	<b>f(x)</b>
<b>0</b>	<b>-1</b>	<b>0.0385</b>
<b>1</b>	<b>-0.75</b>	<b>0.0664</b>
<b>2</b>	<b>-0.5</b>	<b>0.138</b>
<b>3</b>	<b>-0.25</b>	<b>0.3902</b>
<b>4</b>	<b>0.00</b>	<b>1.00</b>
<b>5</b>	<b>0.25</b>	<b>0.3902</b>
<b>6</b>	<b>0.50</b>	<b>0.138</b>
<b>7</b>	<b>0.75</b>	<b>0.0664</b>
<b>8</b>	<b>1.00</b>	<b>0.0385</b>





**The true Function is *Runge Function***

$$f(x) = \frac{1}{1 + 25x^2}$$

