

# Duality between Total Variation and Equivalence Couplings

Adam Quinn Jaffe

April 12, 2023

# I. Stochastic processes

## Theorem (Ernst-Kendall-Roberts-Rosenthal, 2019)

## **Theorem (Ernst-Kendall-Roberts-Rosenthal, 2019)**

*For any  $\theta_1, \theta_2 \in \mathbb{R}$ , one can construct a probability space on which are defined*

## Theorem (Ernst-Kendall-Roberts-Rosenthal, 2019)

For any  $\theta_1, \theta_2 \in \mathbb{R}$ , one can construct a probability space on which are defined

- ▶ a Brownian motion  $B^{\theta_1} = \{B_t^{\theta_1}\}_{t \geq 0}$  with drift  $\theta_1$ ,

## Theorem (Ernst-Kendall-Roberts-Rosenthal, 2019)

For any  $\theta_1, \theta_2 \in \mathbb{R}$ , one can construct a probability space on which are defined

- ▶ a Brownian motion  $B^{\theta_1} = \{B_t^{\theta_1}\}_{t \geq 0}$  with drift  $\theta_1$ ,
- ▶ a Brownian motion  $B^{\theta_2} = \{B_t^{\theta_2}\}_{t \geq 0}$  with drift  $\theta_2$ , and

## Theorem (Ernst-Kendall-Roberts-Rosenthal, 2019)

For any  $\theta_1, \theta_2 \in \mathbb{R}$ , one can construct a probability space on which are defined


- ▶ a Brownian motion  $B^{\theta_1} = \{B_t^{\theta_1}\}_{t \geq 0}$  with drift  $\theta_1$ ,
- ▶ a Brownian motion  $B^{\theta_2} = \{B_t^{\theta_2}\}_{t \geq 0}$  with drift  $\theta_2$ , and
- ▶ a random time  $T$  with  $T > 0$  almost surely,

## Theorem (Ernst-Kendall-Roberts-Rosenthal, 2019)

For any  $\theta_1, \theta_2 \in \mathbb{R}$ , one can construct a probability space on which are defined

- ▶ a Brownian motion  $B^{\theta_1} = \{B_t^{\theta_1}\}_{t \geq 0}$  with drift  $\theta_1$ ,
- ▶ a Brownian motion  $B^{\theta_2} = \{B_t^{\theta_2}\}_{t \geq 0}$  with drift  $\theta_2$ , and
- ▶ a random time  $T$  with  $T > 0$  almost surely,

such that  $B_t^{\theta_1} = B_t^{\theta_2}$  for all  $0 \leq t \leq T$  almost surely.



In words:

In words:

- BM with drift “starts out as” a BM without drift.

In words:

- ▶ BM with drift “starts out as” a BM without drift.
- ▶ BMs with drift are all “locally equivalent” at time zero.

In words:

- ▶ BM with drift “starts out as” a BM without drift.
- ▶ BMs with drift are all “locally equivalent” at time zero.
- ▶ The drift of a BM cannot be detected, if it is only observed up to an adversarially-chosen time.

In words:

- ▶ BM with drift “starts out as” a BM without drift.
- ▶ BMs with drift are all “locally equivalent” at time zero.
- ▶ The drift of a BM cannot be detected, if it is only observed up to an adversarially-chosen time.

Explicit construction based on Itô excursion theory.

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

- ▶ a stochastic process  $X = \{X_t\}_{t \geq 0}$  with law  $\mathbb{P}$ ,

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

- ▶ a stochastic process  $X = \{X_t\}_{t \geq 0}$  with law  $\mathbb{P}$ ,
- ▶ a stochastic process  $X' = \{X'_t\}_{t \geq 0}$  with law  $\mathbb{P}'$ , and

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

- ▶ a stochastic process  $X = \{X_t\}_{t \geq 0}$  with law  $\mathbb{P}$ ,
- ▶ a stochastic process  $X' = \{X'_t\}_{t \geq 0}$  with law  $\mathbb{P}'$ , and
- ▶ a random time  $T$  with  $T > 0$  almost surely,

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

- ▶ a stochastic process  $X = \{X_t\}_{t \geq 0}$  with law  $\mathbb{P}$ ,
- ▶ a stochastic process  $X' = \{X'_t\}_{t \geq 0}$  with law  $\mathbb{P}'$ , and
- ▶ a random time  $T$  with  $T > 0$  almost surely,

such that  $X_t = X'_t$  for all  $0 \leq t \leq T$  almost surely.

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

- ▶ a stochastic process  $X = \{X_t\}_{t \geq 0}$  with law  $\mathbb{P}$ ,
- ▶ a stochastic process  $X' = \{X'_t\}_{t \geq 0}$  with law  $\mathbb{P}'$ , and
- ▶ a random time  $T$  with  $T > 0$  almost surely,

such that  $X_t = X'_t$  for all  $0 \leq t \leq T$  almost surely.

Know that  $(\mathbb{W}^{\theta_1}, \mathbb{W}^{\theta_2})$  has the GCP for all  $\theta_1, \theta_2 \in \mathbb{R}$ , where  $\mathbb{W}^\theta$  denotes the law of BM with drift  $\theta \in \mathbb{R}$ .

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

- ▶ a stochastic process  $X = \{X_t\}_{t \geq 0}$  with law  $\mathbb{P}$ ,
- ▶ a stochastic process  $X' = \{X'_t\}_{t \geq 0}$  with law  $\mathbb{P}'$ , and
- ▶ a random time  $T$  with  $T > 0$  almost surely,

such that  $X_t = X'_t$  for all  $0 \leq t \leq T$  almost surely.

Know that  $(\mathbb{W}^{\theta_1}, \mathbb{W}^{\theta_2})$  has the GCP for all  $\theta_1, \theta_2 \in \mathbb{R}$ , where  $\mathbb{W}^\theta$  denotes the law of BM with drift  $\theta \in \mathbb{R}$ .

Say  $\mathbb{P}$  has the *Brownian GCP* if  $(\mathbb{P}, \mathbb{W}^0)$  has the GCP.

## Definition

Say that a pair of probability measures  $(\mathbb{P}, \mathbb{P}')$  on  $D([0, \infty); \mathbb{R})$  has the *germ coupling property (GCP)* if one can construct a probability space on which are defined

- ▶ a stochastic process  $X = \{X_t\}_{t \geq 0}$  with law  $\mathbb{P}$ ,
- ▶ a stochastic process  $X' = \{X'_t\}_{t \geq 0}$  with law  $\mathbb{P}'$ , and
- ▶ a random time  $T$  with  $T > 0$  almost surely,

such that  $X_t = X'_t$  for all  $0 \leq t \leq T$  almost surely.

Know that  $(\mathbb{W}^{\theta_1}, \mathbb{W}^{\theta_2})$  has the GCP for all  $\theta_1, \theta_2 \in \mathbb{R}$ , where  $\mathbb{W}^\theta$  denotes the law of BM with drift  $\theta \in \mathbb{R}$ .

Say  $\mathbb{P}$  has the *Brownian GCP* if  $(\mathbb{P}, \mathbb{W}^0)$  has the GCP.

Which other pairs have the GCP?

## II. Some vignettes



- $\Omega$  a Polish space,

- ▶  $\Omega$  a Polish space,
- ▶  $\Delta := \{(x, x) \in \Omega \times \Omega : x \in \Omega\}$  the diagonal in  $\Omega \times \Omega$ ,

- ▶  $\Omega$  a Polish space,
- ▶  $\Delta := \{(x, x) \in \Omega \times \Omega : x \in \Omega\}$  the diagonal in  $\Omega \times \Omega$ ,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega$ , and

- ▶  $\Omega$  a Polish space,
- ▶  $\Delta := \{(x, x) \in \Omega \times \Omega : x \in \Omega\}$  the diagonal in  $\Omega \times \Omega$ ,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .


- ▶  $\Omega$  a Polish space,
- ▶  $\Delta := \{(x, x) \in \Omega \times \Omega : x \in \Omega\}$  the diagonal in  $\Omega \times \Omega$ ,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .


Then (classical, Doeblin):


- ▶  $\Omega$  a Polish space,
- ▶  $\Delta := \{(x, x) \in \Omega \times \Omega : x \in \Omega\}$  the diagonal in  $\Omega \times \Omega$ ,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .


Then (classical, Doeblin):

$$\sup_{A \in \mathcal{B}(\Omega)} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(\Delta)).$$











$$\tilde{P} := Q \circ (\text{id}, \text{id})^{-1} + \gamma (P - Q) \otimes (P' - Q)$$



- ▶  $\Omega$  a Polish space,
- ▶  $\Delta := \{(x, x) \in \Omega \times \Omega : x \in \Omega\}$  the diagonal in  $\Omega \times \Omega$ ,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .

Then (classical, Doeblin):

$$\sup_{A \in \mathcal{B}(\Omega)} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(\Delta)).$$

- $\Omega_0 := \{0, 1\}^{\mathbb{N}}$  the space of binary sequences,

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{N}}$  the space of binary sequences,
- ▶  $E_0 := \bigcup_{n \in \mathbb{N}} \{(x, x') \in \Omega_0 \times \Omega_0 : (x_n, x_{n+1}, \dots) = (x'_n, x'_{n+1}, \dots)\}$   
the equivalence relation of eventual equality,

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{N}}$  the space of binary sequences,
- ▶  $E_0 := \bigcup_{n \in \mathbb{N}} \{(x, x') \in \Omega_0 \times \Omega_0 : (x_n, x_{n+1}, \dots) = (x'_n, x'_{n+1}, \dots)\}$   
the equivalence relation of eventual equality,
- ▶  $\mathcal{T} := \bigcap_{n \in \mathbb{N}} \sigma(x_n, x_{n+1}, \dots)$  the tail  $\sigma$ -algebra,

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{N}}$  the space of binary sequences,
- ▶  $E_0 := \bigcup_{n \in \mathbb{N}} \{(x, x') \in \Omega_0 \times \Omega_0 : (x_n, x_{n+1}, \dots) = (x'_n, x'_{n+1}, \dots)\}$   
the equivalence relation of eventual equality,
- ▶  $\mathcal{T} := \bigcap_{n \in \mathbb{N}} \sigma(x_n, x_{n+1}, \dots)$  the tail  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{N}}$  the space of binary sequences,
- ▶  $E_0 := \bigcup_{n \in \mathbb{N}} \{(x, x') \in \Omega_0 \times \Omega_0 : (x_n, x_{n+1}, \dots) = (x'_n, x'_{n+1}, \dots)\}$   
the equivalence relation of eventual equality,
- ▶  $\mathcal{T} := \bigcap_{n \in \mathbb{N}} \sigma(x_n, x_{n+1}, \dots)$  the tail  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{N}}$  the space of binary sequences,
- ▶  $E_0 := \bigcup_{n \in \mathbb{N}} \{(x, x') \in \Omega_0 \times \Omega_0 : (x_n, x_{n+1}, \dots) = (x'_n, x'_{n+1}, \dots)\}$   
the equivalence relation of eventual equality,
- ▶  $\mathcal{T} := \bigcap_{n \in \mathbb{N}} \sigma(x_n, x_{n+1}, \dots)$  the tail  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .

Then (Griffeath, Pitman, Goldstein):

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{N}}$  the space of binary sequences,
- ▶  $E_0 := \bigcup_{n \in \mathbb{N}} \{(x, x') \in \Omega_0 \times \Omega_0 : (x_n, x_{n+1}, \dots) = (x'_n, x'_{n+1}, \dots)\}$   
the equivalence relation of eventual equality,
- ▶  $\mathcal{T} := \bigcap_{n \in \mathbb{N}} \sigma(x_n, x_{n+1}, \dots)$  the tail  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .

Then (Griffeath, Pitman, Goldstein):

$$\sup_{A \in \mathcal{T}} |\mathbb{P}(A) - \mathbb{P}'(A)| = 0 \text{ if and only if } \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E_0)) = 0.$$

- $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,

- $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,
- $\theta : \Omega_0 \rightarrow \Omega_0$  the left-shift operation,

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,
- ▶  $\theta : \Omega_0 \rightarrow \Omega_0$  the left-shift operation,
- ▶  $E_{\mathbb{Z}} := \bigcup_{n \in \mathbb{Z}} \{(x, x') \in \Omega_0 \times \Omega_0 : \theta^n(x) = x'\}$  the equivalence relation of shift-equivalence,

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,
- ▶  $\theta : \Omega_0 \rightarrow \Omega_0$  the left-shift operation,
- ▶  $E_{\mathbb{Z}} := \bigcup_{n \in \mathbb{Z}} \{(x, x') \in \Omega_0 \times \Omega_0 : \theta^n(x) = x'\}$  the equivalence relation of shift-equivalence,
- ▶  $\mathcal{I}_{\mathbb{Z}} = \{A \in \mathcal{B}(\Omega_0) : \theta^{-1}(A) = A\}$  the shift-invariant  $\sigma$ -algebra,

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,
- ▶  $\theta : \Omega_0 \rightarrow \Omega_0$  the left-shift operation,
- ▶  $E_{\mathbb{Z}} := \bigcup_{n \in \mathbb{Z}} \{(x, x') \in \Omega_0 \times \Omega_0 : \theta^n(x) = x'\}$  the equivalence relation of shift-equivalence,
- ▶  $\mathcal{I}_{\mathbb{Z}} = \{A \in \mathcal{B}(\Omega_0) : \theta^{-1}(A) = A\}$  the shift-invariant  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,
- ▶  $\theta : \Omega_0 \rightarrow \Omega_0$  the left-shift operation,
- ▶  $E_{\mathbb{Z}} := \bigcup_{n \in \mathbb{Z}} \{(x, x') \in \Omega_0 \times \Omega_0 : \theta^n(x) = x'\}$  the equivalence relation of shift-equivalence,
- ▶  $\mathcal{I}_{\mathbb{Z}} = \{A \in \mathcal{B}(\Omega_0) : \theta^{-1}(A) = A\}$  the shift-invariant  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,
- ▶  $\theta : \Omega_0 \rightarrow \Omega_0$  the left-shift operation,
- ▶  $E_{\mathbb{Z}} := \bigcup_{n \in \mathbb{Z}} \{(x, x') \in \Omega_0 \times \Omega_0 : \theta^n(x) = x'\}$  the equivalence relation of shift-equivalence,
- ▶  $\mathcal{I}_{\mathbb{Z}} = \{A \in \mathcal{B}(\Omega_0) : \theta^{-1}(A) = A\}$  the shift-invariant  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .

Then (Aldous-Thorisson):

- ▶  $\Omega_0 := \{0, 1\}^{\mathbb{Z}}$  the space of binary sequences,
- ▶  $\theta : \Omega_0 \rightarrow \Omega_0$  the left-shift operation,
- ▶  $E_{\mathbb{Z}} := \bigcup_{n \in \mathbb{Z}} \{(x, x') \in \Omega_0 \times \Omega_0 : \theta^n(x) = x'\}$  the equivalence relation of shift-equivalence,
- ▶  $\mathcal{I}_{\mathbb{Z}} = \{A \in \mathcal{B}(\Omega_0) : \theta^{-1}(A) = A\}$  the shift-invariant  $\sigma$ -algebra,
- ▶  $\mathbb{P}, \mathbb{P}'$  two Borel probability measures on  $\Omega_0$ , and
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  the space of all couplings of  $\mathbb{P}$  and  $\mathbb{P}'$ .

Then (Aldous-Thorisson):

$$\sup_{A \in \mathcal{I}_{\mathbb{Z}}} |\mathbb{P}(A) - \mathbb{P}'(A)| = 0 \text{ if and only if } \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E_{\mathbb{Z}})) = 0.$$

$$\sup_{A \in \mathcal{B}(\Omega)} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')}(1 - \tilde{\mathbb{P}}(\Delta)).$$

$$\sup_{A \in \mathcal{T}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(\textcolor{teal}{E}_0)).$$

$$\sup_{A \in \textcolor{teal}{\mathcal{L}}_{\mathbb{Z}}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(\textcolor{teal}{E}_{\mathbb{Z}})).$$

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)| \stackrel{?}{=} \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(\textcolor{teal}{E})).$$

In many probability settings one wants to study the *E-coupling problem*

In many probability settings one wants to study the *E-coupling problem*

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

In many probability settings one wants to study the *E-coupling problem*

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

In general this problem is hard to solve and there are not many general-purpose tools available.

In many probability settings one wants to study the *E-coupling problem*

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

In general this problem is hard to solve and there are not many general-purpose tools available.

On the other hand, the *G-total variation problem*

In many probability settings one wants to study the *E-coupling problem*

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

In general this problem is hard to solve and there are not many general-purpose tools available.

On the other hand, the *G-total variation problem*

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)|$$

In many probability settings one wants to study the *E-coupling problem*

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

In general this problem is hard to solve and there are not many general-purpose tools available.

On the other hand, the *G-total variation problem*

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)|$$

is typically easy to analyze for probabilists.

In many probability settings one wants to study the *E-coupling problem*

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

In general this problem is hard to solve and there are not many general-purpose tools available.

On the other hand, the *G-total variation problem*

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)|$$

is typically easy to analyze for probabilists.

These optimization problems are closely related!

In many probability settings one wants to study the *E-coupling problem*

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

In general this problem is hard to solve and there are not many general-purpose tools available.

On the other hand, the *G-total variation problem*

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)|$$

is typically easy to analyze for probabilists.

These optimization problems are closely related! In fact, we'll see that they are often dual, in the sense of mathematical optimization.

- I. Stochastic processes
- II. Some vignettes
- III. Problem statement
- IV. Results
- V. Applications

### **III. Problem statement**

Notation:

Notation:

- $(\Omega, \mathcal{F})$  measurable space,

Notation:

- ▶  $(\Omega, \mathcal{F})$  measurable space,
- ▶  $\mathcal{P}(\Omega, \mathcal{F})$  space of probability measures on  $(\Omega, \mathcal{F})$ ,

Notation:

- ▶  $(\Omega, \mathcal{F})$  measurable space,
- ▶  $\mathcal{P}(\Omega, \mathcal{F})$  space of probability measures on  $(\Omega, \mathcal{F})$ ,
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  space of couplings of  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ ,

Notation:

- ▶  $(\Omega, \mathcal{F})$  measurable space,
- ▶  $\mathcal{P}(\Omega, \mathcal{F})$  space of probability measures on  $(\Omega, \mathcal{F})$ ,
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  space of couplings of  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ ,
- ▶  $E$  equivalence relation on  $\Omega$ ,

Notation:

- ▶  $(\Omega, \mathcal{F})$  measurable space,
- ▶  $\mathcal{P}(\Omega, \mathcal{F})$  space of probability measures on  $(\Omega, \mathcal{F})$ ,
- ▶  $\Pi(\mathbb{P}, \mathbb{P}')$  space of couplings of  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ ,
- ▶  $E$  equivalence relation on  $\Omega$ , and
- ▶  $\mathcal{G}$  sub- $\sigma$ -algebra of  $\mathcal{F}$ .

# Definition

## Definition

Say  $E$  is *measurable* if  $E \in \mathcal{F} \otimes \mathcal{F}$ .

## Definition

Say  $E$  is *measurable* if  $E \in \mathcal{F} \otimes \mathcal{F}$ . Say  $(E, \mathcal{G})$  is *strongly dual* if  $E$  is measurable and if we have

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')}(1 - \tilde{\mathbb{P}}(E)).$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ .

## Definition

Say  $E$  is *measurable* if  $E \in \mathcal{F} \otimes \mathcal{F}$ . Say  $(E, \mathcal{G})$  is *strongly dual* if  $E$  is measurable and if we have

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')}(1 - \tilde{\mathbb{P}}(E)).$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ .

Roughly speaking,  $(\Delta, \mathcal{B}(\Omega))$ ,  $(E_0, \mathcal{T})$ , and  $(E_{\mathbb{Z}}, \mathcal{I}_{\mathbb{Z}})$  are strongly dual.

## Definition

Say  $E$  is *measurable* if  $E \in \mathcal{F} \otimes \mathcal{F}$ . Say  $(E, \mathcal{G})$  is *strongly dual* if  $E$  is measurable and if we have

$$\sup_{A \in \mathcal{G}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')}(1 - \tilde{\mathbb{P}}(E)).$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ .

Roughly speaking,  $(\Delta, \mathcal{B}(\Omega))$ ,  $(E_0, \mathcal{T})$ , and  $(E_{\mathbb{Z}}, \mathcal{I}_{\mathbb{Z}})$  are strongly dual.

Which pairs  $(E, \mathcal{G})$  are strongly dual?

If  $E$  is given, then there is a natural choice of  $\mathcal{G}$ :

If  $E$  is given, then there is a natural choice of  $\mathcal{G}$ :

**Lemma (AQJ)**

If  $E$  is given, then there is a natural choice of  $\mathcal{G}$ :

### **Lemma (AQJ)**

*If  $(E, \mathcal{G})$  is strongly dual for some  $\mathcal{G}$ ,*

If  $E$  is given, then there is a natural choice of  $\mathcal{G}$ :

### **Lemma (AQJ)**

*If  $(E, \mathcal{G})$  is strongly dual for some  $\mathcal{G}$ , then  $(E, E^*)$  is strongly dual,*

If  $E$  is given, then there is a natural choice of  $\mathcal{G}$ :

### Lemma (AQJ)

*If  $(E, \mathcal{G})$  is strongly dual for some  $\mathcal{G}$ , then  $(E, E^*)$  is strongly dual, where  $E^*$  is the  $E$ -invariant  $\sigma$ -algebra*

$$E^* := \{A \in \mathcal{F} : \forall (x, x') \in E(x \in A \Leftrightarrow x' \in A)\}$$

If  $E$  is given, then there is a natural choice of  $\mathcal{G}$ :

### Lemma (AQJ)

*If  $(E, \mathcal{G})$  is strongly dual for some  $\mathcal{G}$ , then  $(E, E^*)$  is strongly dual, where  $E^*$  is the  $E$ -invariant  $\sigma$ -algebra*

$$E^* := \{A \in \mathcal{F} : \forall (x, x') \in E(x \in A \Leftrightarrow x' \in A)\}$$

*Say that  $E$  is strongly dualizable if  $(E, E^*)$  is strongly dual.*

Connection to optimal transport?

Connection to optimal transport?

Note that the  $E$ -coupling problem

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)).$$

is exactly a Monge-Kantorovich problem

Connection to optimal transport?

Note that the  $E$ -coupling problem

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)).$$

is exactly a Monge-Kantorovich problem with cost function

$$c(x, x') = 1 - \mathbb{1}\{(x, x') \in E\}.$$

Connection to optimal transport?


Note that the  $E$ -coupling problem

$$\inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)).$$

is exactly a Monge-Kantorovich problem with cost function

$$c(x, x') = 1 - \mathbb{1}\{(x, x') \in E\}.$$

In words: cost 0 to move within an equivalence class, and cost 1 to move between equivalence classes.



Classical Monge-Kantorovich theory requires topological regularity:

Classical Monge-Kantorovich theory requires topological regularity:  $\Omega$  is a Polish space,  $\mathcal{F}$  is its Borel  $\sigma$ -algebra, and  $c$  is lower semi-continuous.

In our setting, this requires  $E$  to be closed in  $\Omega \times \Omega$ ;

Classical Monge-Kantorovich theory requires topological regularity:  $\Omega$  is a Polish space,  $\mathcal{F}$  is its Borel  $\sigma$ -algebra, and  $c$  is lower semi-continuous.

In our setting, this requires  $E$  to be closed in  $\Omega \times \Omega$ ; in this case, Kantorovich duality and some standard tricks can show that  $E$  is strongly dualizable.

Classical Monge-Kantorovich theory requires topological regularity:  $\Omega$  is a Polish space,  $\mathcal{F}$  is its Borel  $\sigma$ -algebra, and  $c$  is lower semi-continuous.

In our setting, this requires  $E$  to be closed in  $\Omega \times \Omega$ ; in this case, Kantorovich duality and some standard tricks can show that  $E$  is strongly dualizable.

However, most interesting equivalence relations, from the point of view of probability, are  $F_\sigma$  (countable union of closed) in  $\Omega \times \Omega$ .

Classical Monge-Kantorovich theory requires topological regularity:  $\Omega$  is a Polish space,  $\mathcal{F}$  is its Borel  $\sigma$ -algebra, and  $c$  is lower semi-continuous.

In our setting, this requires  $E$  to be closed in  $\Omega \times \Omega$ ; in this case, Kantorovich duality and some standard tricks can show that  $E$  is strongly dualizable.

However, most interesting equivalence relations, from the point of view of probability, are  $F_\sigma$  (countable union of closed) in  $\Omega \times \Omega$ .

**Idea:** One can exchange topological regularity (lower semi-continuity of the cost function) with algebraic regularity (the cost function is supported on an equivalence relation) while maintaining a form of duality.

## IV. Results

Some useful reductions:

Some useful reductions:

It is easy to show that we always have *weak dualizability*,

Some useful reductions:

It is easy to show that we always have *weak dualizability*, that

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)| \leq \inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ .

Some useful reductions:

It is easy to show that we always have *weak dualizability*, that

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)| \leq \inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . The difficult part is showing the reverse inequality and that the inf is attained.

Some useful reductions:

It is easy to show that we always have *weak dualizability*, that

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)| \leq \inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . The difficult part is showing the reverse inequality and that the inf is attained.

We say that  $E$  is *quasi-strongly dualizable* if

Some useful reductions:

It is easy to show that we always have *weak dualizability*, that

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)| \leq \inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . The difficult part is showing the reverse inequality and that the inf is attained.

We say that  $E$  is *quasi-strongly dualizable* if for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$  the following are equivalent:

Some useful reductions:

It is easy to show that we always have *weak dualizability*, that

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)| \leq \inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . The difficult part is showing the reverse inequality and that the inf is attained.

We say that  $E$  is *quasi-strongly dualizable* if for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$  the following are equivalent:

- $\mathbb{P}(A) = \mathbb{P}'(A)$  for all  $A \in E^*$

Some useful reductions:

It is easy to show that we always have *weak dualizability*, that

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)| \leq \inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . The difficult part is showing the reverse inequality and that the inf is attained.

We say that  $E$  is *quasi-strongly dualizable* if for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$  the following are equivalent:

- ▶  $\mathbb{P}(A) = \mathbb{P}'(A)$  for all  $A \in E^*$
- ▶ There exists  $\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')$  and  $N \in \mathcal{F} \otimes \mathcal{F}$  with  $\tilde{\mathbb{P}}(N) = 0$  and  $(\Omega \times \Omega) \setminus E \subseteq N$ .

Some useful reductions:

It is easy to show that we always have *weak dualizability*, that

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)| \leq \inf_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)),$$

for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . The difficult part is showing the reverse inequality and that the inf is attained.

We say that  $E$  is *quasi-strongly dualizable* if for all  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$  the following are equivalent:

- ▶  $\mathbb{P}(A) = \mathbb{P}'(A)$  for all  $A \in E^*$
- ▶ There exists  $\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')$  and  $N \in \mathcal{F} \otimes \mathcal{F}$  with  $\tilde{\mathbb{P}}(N) = 0$  and  $(\Omega \times \Omega) \setminus E \subseteq N$ .

Then  $E$  is strongly dualizable if and only if it is measurable and quasi-strongly dualizable.

Some basic descriptive set theory:

Some basic descriptive set theory:

A measurable space  $(S, \mathcal{S})$  is called a *standard Borel space* if there exists a Polish topology  $\tau$  on  $\Omega$  such that  $\mathcal{S} = \mathcal{B}(\tau)$ .

Some basic descriptive set theory:

A measurable space  $(S, \mathcal{S})$  is called a *standard Borel space* if there exists a Polish topology  $\tau$  on  $\Omega$  such that  $\mathcal{S} = \mathcal{B}(\tau)$ .

An equivalence relation  $E$  on an abstract measurable space  $(\Omega, \mathcal{F})$  is called *smooth* if

Some basic descriptive set theory:

A measurable space  $(S, \mathcal{S})$  is called a *standard Borel space* if there exists a Polish topology  $\tau$  on  $\Omega$  such that  $\mathcal{S} = \mathcal{B}(\tau)$ .

An equivalence relation  $E$  on an abstract measurable space  $(\Omega, \mathcal{F})$  is called *smooth* if there exists a standard Borel space  $(S, \mathcal{S})$  and a measurable function  $\phi : (\Omega, \mathcal{F}) \rightarrow (S, \mathcal{S})$

Some basic descriptive set theory:

A measurable space  $(S, \mathcal{S})$  is called a *standard Borel space* if there exists a Polish topology  $\tau$  on  $\Omega$  such that  $\mathcal{S} = \mathcal{B}(\tau)$ .

An equivalence relation  $E$  on an abstract measurable space  $(\Omega, \mathcal{F})$  is called *smooth* if there exists a standard Borel space  $(S, \mathcal{S})$  and a measurable function  $\phi : (\Omega, \mathcal{F}) \rightarrow (S, \mathcal{S})$  such that  $(x, x') \in E$  is equivalent to  $\phi(x) = \phi(x')$ .

Some basic descriptive set theory:

A measurable space  $(S, \mathcal{S})$  is called a *standard Borel space* if there exists a Polish topology  $\tau$  on  $\Omega$  such that  $\mathcal{S} = \mathcal{B}(\tau)$ .

An equivalence relation  $E$  on an abstract measurable space  $(\Omega, \mathcal{F})$  is called *smooth* if there exists a standard Borel space  $(S, \mathcal{S})$  and a measurable function  $\phi : (\Omega, \mathcal{F}) \rightarrow (S, \mathcal{S})$  such that  $(x, x') \in E$  is equivalent to  $\phi(x) = \phi(x')$ .

Roughly speaking,  $E$  is smooth if and only if the quotient  $\Omega/E$  can be given a natural standard Borel structure.

## **Lemma (AQJ)**

*The following are equivalent:*

## **Lemma (AQJ)**

*The following are equivalent:*

- (i) *E is smooth.*

## **Lemma (AQJ)**

*The following are equivalent:*

- (i) *E is smooth.*
- (ii)  *$E^*$  is countably generated.*

## **Lemma (AQJ)**

*The following are equivalent:*

- (i)  $E$  is smooth.
- (ii)  $E^*$  is countably generated.
- (iii)  $E \in E^* \otimes E^*$ .

## **Lemma (AQJ)**

*The following are equivalent:*

- (i)  $E$  is smooth.
- (ii)  $E^*$  is countably generated.
- (iii)  $E \in E^* \otimes E^*$ .

The equivalence between (i) and (ii) is classical, but the equivalence with (iii) appears to be novel.

## Theorem (AQJ)

*Every smooth equivalence relation is strongly dualizable.*

## Theorem (AQJ)

*Every smooth equivalence relation is strongly dualizable.*


Consequence: All equivalence relations with  $G_\delta$  (countable intersection of open) equivalence classes are strongly dualizable.


## Theorem (AQJ)


*Every smooth equivalence relation is strongly dualizable.*


Consequence: All equivalence relations with  $G_\delta$  (countable intersection of open) equivalence classes are strongly dualizable.


Idea of proof: Do the Doeblin coupling in  $(\Omega, E^*)$  then pull back.












$\tilde{\mu} := \nu \circ (\text{id}, \text{id})^{-1} + \gamma(\mu - \nu) \otimes (\mu' - \nu)$

$(\Omega, E^*)$

$(\Omega, E^*)$

$\mu$

$\mu'$



However, many equivalence relations of interest are not smooth.

However, many equivalence relations of interest are not smooth.  
Instead, we have the following closure result:

However, many equivalence relations of interest are not smooth.  
Instead, we have the following closure result:

### **Theorem (AQJ)**




*A countable increasing union of strongly dualizable equivalence relations is strongly dualizable.*


However, many equivalence relations of interest are not smooth.  
Instead, we have the following closure result:

### **Theorem (AQJ)**


*A countable increasing union of strongly dualizable equivalence relations is strongly dualizable.*

Idea of proof: Iterate greedily.










P







P'






$\tilde{P}_1(\cdot \cap E_1)$  $\tilde{P}_1$  $P - \tilde{P}_1(\cdot \cap E_1) \circ \pi^{-1}$  $P' - \tilde{P}_1(\cdot \cap E_1) \circ \pi'^{-1}$ 

$\tilde{\mathbb{P}}_1(\cdot \cap E_1)$  $\mathbb{P} - \tilde{\mathbb{P}}_1(\cdot \cap E_1) \circ \pi^{-1}$  $\mathbb{P}' - \tilde{\mathbb{P}}_1(\cdot \cap E_1) \circ \pi^{-1}$ 

$\widetilde{\mathbb{P}}_1(\cdot \cap E_1)$  $\widetilde{\mathbb{P}}_2$  $\mathbb{P} - \widetilde{\mathbb{P}}_1(\cdot \cap E_1) \circ \pi^{-1}$  $\mathbb{P}' - \widetilde{\mathbb{P}}_1(\cdot \cap E_1) \circ \pi'^{-1}$

$$\sum_{i=1}^2 \tilde{P}_1(\cdot \cap E_i)$$




$$\tilde{P}_2$$



$$P - \sum_{i=1}^2 \tilde{P}_1(\cdot \cap E_i) \circ \pi^{-1}$$

$$P' - \sum_{i=1}^2 \tilde{P}_1(\cdot \cap E_i) \circ \pi^{-1}$$


$$\sum_{i=1}^2 \tilde{P}_i(\cdot \cap E_i)$$



$$P - \sum_{i=1}^2 \tilde{P}_i(\cdot \cap E_i) \circ \pi^{-1}$$

$$P' - \sum_{i=1}^2 \tilde{P}_i(\cdot \cap E_i) \circ \pi'^{-1}$$


$$\sum_{i=1}^2 \tilde{\mathbb{P}}_i(\cdot \cap E_i)$$



$$\tilde{\mathbb{P}}_3$$




$$\mathbb{P} - \sum_{i=1}^2 \tilde{\mathbb{P}}_i(\cdot \cap E_i) \circ \pi^{-1}$$



$$\mathbb{P}' - \sum_{i=1}^2 \tilde{\mathbb{P}}_i(\cdot \cap E_i) \circ \pi^{-1}$$

$$\sum_{i=1}^3 \tilde{P}_i(\cdot \cap E_i)$$

$$\tilde{P}_3$$



$$P - \sum_{i=1}^3 \tilde{P}_i(\cdot \cap E_i) \circ \pi^{-1}$$

$$P' - \sum_{i=1}^3 \tilde{P}_i(\cdot \cap E_i) \circ \pi'^{-1}$$



This shows strong dualizability for all *hypersmooth* (countable union of smooth) equivalence relations,

This shows strong dualizability for all *hypersmooth* (countable union of smooth) equivalence relations, and this covers most “reasonable” equivalence relations occurring in probability.

## V. Applications

If  $E$  is strongly dualizable, then the  $E$ -coupling problem

$$\min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E))$$

If  $E$  is strongly dualizable, then the  $E$ -coupling problem

$$\min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E))$$

has a minimizer and is equivalent to the  $E^*$ -total variation problem:

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)|.$$

If  $E$  is strongly dualizable, then the  $E$ -coupling problem

$$\min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E))$$

has a minimizer and is equivalent to the  $E^*$ -total variation problem:

$$\sup_{A \in E^*} |\mathbb{P}(A) - \mathbb{P}'(A)|.$$

Thus, the  $E$ -coupling problem can be studied with the help of very many soft arguments.

Example from algorithms:

Suppose  $(X_1, X_2, \dots)$  is an i.i.d. real-valued sequence with common distribution  $\mu$ .

Example from algorithms:

Suppose  $(X_1, X_2, \dots)$  is an i.i.d. real-valued sequence with common distribution  $\mu$ . If  $A$  is a (possibly randomized) sorting algorithm terminating in a finite number of swaps, one can consider the distribution of the random sequence  $A(X_1, X_2, \dots)$ .

Example from algorithms:

Suppose  $(X_1, X_2, \dots)$  is an i.i.d. real-valued sequence with common distribution  $\mu$ . If  $A$  is a (possibly randomized) sorting algorithm terminating in a finite number of swaps, one can consider the distribution of the random sequence  $A(X_1, X_2, \dots)$ .

Which random sequences can be simulated in the way, by applying a finite, randomized sorting algorithm to an i.i.d. sequence?

$$\begin{array}{l} X : \quad 1 \ -1 \ 2 \ 2 \ 0 \ 3 \ -2 \ 1 \ 3 \ \cdots \cdots \\ X' : \quad 2 \ -1 \ 0 \ 3 \ 2 \ 1 \ -2 \ 1 \ 3 \ \cdots \cdots \end{array}$$

A pink curved arrow points from the second row of  $X$  to the first row of  $X'$ . Several pink arrows point from specific entries in the second row of  $X$  to corresponding entries in the first row of  $X'$ , specifically from 1 to 2, -1 to 0, 2 to 3, and 0 to 2.

- $\Omega := \mathbb{R}^{\mathbb{N}}$  the space of real sequences,

- ▶  $\Omega := \mathbb{R}^{\mathbb{N}}$  the space of real sequences,
- ▶  $E_{\infty}$  the equivalence relation of agreeing up to a permutation of finitely many elements.

- ▶  $\Omega := \mathbb{R}^{\mathbb{N}}$  the space of real sequences,
- ▶  $E_{\infty}$  the equivalence relation of agreeing up to a permutation of finitely many elements.
- ▶  $\mathcal{E}$  the exchangeable  $\sigma$ -algebra.

- ▶  $\Omega := \mathbb{R}^{\mathbb{N}}$  the space of real sequences,
- ▶  $E_{\infty}$  the equivalence relation of agreeing up to a permutation of finitely many elements.
- ▶  $\mathcal{E}$  the exchangeable  $\sigma$ -algebra.

Then  $E_{\infty}$  is strongly dualizable, and we have  $E_{\infty}^* = \mathcal{E}$ , and

$$\sup_{A \in \mathcal{E}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')}(1 - \tilde{\mathbb{P}}(E_{\infty})).$$

- ▶  $\Omega := \mathbb{R}^{\mathbb{N}}$  the space of real sequences,
- ▶  $E_{\infty}$  the equivalence relation of agreeing up to a permutation of finitely many elements.
- ▶  $\mathcal{E}$  the exchangeable  $\sigma$ -algebra.

Then  $E_{\infty}$  is strongly dualizable, and we have  $E_{\infty}^* = \mathcal{E}$ , and

$$\sup_{A \in \mathcal{E}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E_{\infty})).$$

Consequence: If  $\mathbb{P} \ll \mu^{\otimes \mathbb{N}}$ , then there exists a finite, randomized sorting algorithm which, when given an input consisting of an i.i.d. sequence from  $\mu$ , returns a sequence with law  $\mathbb{P}$ .

- ▶  $\Omega := \mathbb{R}^{\mathbb{N}}$  the space of real sequences,
- ▶  $E_{\infty}$  the equivalence relation of agreeing up to a permutation of finitely many elements.
- ▶  $\mathcal{E}$  the exchangeable  $\sigma$ -algebra.

Then  $E_{\infty}$  is strongly dualizable, and we have  $E_{\infty}^* = \mathcal{E}$ , and

$$\sup_{A \in \mathcal{E}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')}(1 - \tilde{\mathbb{P}}(E_{\infty})).$$

Consequence: If  $\mathbb{P} \ll \mu^{\otimes \mathbb{N}}$ , then there exists a finite, randomized sorting algorithm which, when given an input consisting of an i.i.d. sequence from  $\mu$ , returns a sequence with law  $\mathbb{P}$ .

How to construct such an algorithm explicitly?

Example from point processes:

Example from point processes:

Let  $\mathbb{P}$  be the law of a determinantal point process on  $\mathbb{R}^k$ .


Example from point processes:

Let  $\mathbb{P}$  be the law of a determinantal point process on  $\mathbb{R}^k$ . For  $\mathbf{a} = (a_1, \dots, a_\ell) \in (\mathbb{R}^k)^\ell$ , writing  $|\mathbf{a}| = \ell$ , consider reduced Palm measure  $\mathbb{P}_{\mathbf{a}}$ .

Example from point processes:

Let  $\mathbb{P}$  be the law of a determinantal point process on  $\mathbb{R}^k$ . For  $\mathbf{a} = (a_1, \dots, a_\ell) \in (\mathbb{R}^k)^\ell$ , writing  $|\mathbf{a}| = \ell$ , consider reduced Palm measure  $\mathbb{P}_{\mathbf{a}}$ .

We say that  $\mathbf{a}$  and  $\mathbf{a}'$  have *similar potential in  $\mathbb{P}$*  if there exists a coupling of  $(\omega, \omega')$  of  $\mathbb{P}_{\mathbf{a}}$  and  $\mathbb{P}_{\mathbf{a}'}$  such that  $\omega$  and  $\omega'$  have finitely many disagreements.



- $\Omega$  the space of non-negative, locally-finite, integer-valued, simple Borel measures on  $\mathbb{R}^k$ ,

- ▶  $\Omega$  the space of non-negative, locally-finite, integer-valued, simple Borel measures on  $\mathbb{R}^k$ ,
- ▶  $E$  the equivalence relation of having finitely many disagreements, and

- ▶  $\Omega$  the space of non-negative, locally-finite, integer-valued, simple Borel measures on  $\mathbb{R}^k$ ,
- ▶  $E$  the equivalence relation of having finitely many disagreements, and
- ▶  $\mathcal{F}_K := \sigma(x(U) : U \cap K = \emptyset)$  the  $\sigma$ -algebra of all information occurring outside of the compact set  $K \subseteq \mathbb{R}^k$ , and  
 $\mathcal{T} := \bigcap_{K \text{compact}} \mathcal{F}_K$  the *tail  $\sigma$ -algebra*.

- ▶  $\Omega$  the space of non-negative, locally-finite, integer-valued, simple Borel measures on  $\mathbb{R}^k$ ,
- ▶  $E$  the equivalence relation of having finitely many disagreements, and
- ▶  $\mathcal{F}_K := \sigma(x(U) : U \cap K = \emptyset)$  the  $\sigma$ -algebra of all information occurring outside of the compact set  $K \subseteq \mathbb{R}^k$ , and  
 $\mathcal{T} := \bigcap_{K \text{compact}} \mathcal{F}_K$  the *tail  $\sigma$ -algebra*.

Then  $E$  is strongly dualizable, and we have  $E^* = \mathcal{T}$ , and

$$\sup_{A \in \mathcal{T}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)).$$

- ▶  $\Omega$  the space of non-negative, locally-finite, integer-valued, simple Borel measures on  $\mathbb{R}^k$ ,
- ▶  $E$  the equivalence relation of having finitely many disagreements, and
- ▶  $\mathcal{F}_K := \sigma(x(U) : U \cap K = \emptyset)$  the  $\sigma$ -algebra of all information occurring outside of the compact set  $K \subseteq \mathbb{R}^k$ , and  
 $\mathcal{T} := \bigcap_{K \text{ compact}} \mathcal{F}_K$  the *tail  $\sigma$ -algebra*.

Then  $E$  is strongly dualizable, and we have  $E^* = \mathcal{T}$ , and

$$\sup_{A \in \mathcal{T}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E)).$$

Consequence: If  $\mathbb{P}$  is the law of a DPP with integrable kernel (Ginibre, Gamma, Bessel, Airy) and  $|\mathbf{a}| = |\mathbf{a}'|$ , then  $\mathbf{a}$  and  $\mathbf{a}'$  have similar potential in  $\mathbb{P}$ .

Example from stochastic processes:

Example from stochastic processes:

- $\Omega := D([0, \infty); \mathbb{R})$  the space of càdlàg paths,

Example from stochastic processes:

- ▶  $\Omega := D([0, \infty); \mathbb{R})$  the space of càdlàg paths,
- ▶  $E_{0+}$  the equivalence relation of agreeing for some positive time,

Example from stochastic processes:

- ▶  $\Omega := D([0, \infty); \mathbb{R})$  the space of càdlàg paths,
- ▶  $E_{0+}$  the equivalence relation of agreeing for some positive time,
- ▶  $\mathcal{F}_{0+} := \bigcap_{t>0} \mathcal{F}_t$  the *germ*  $\sigma$ -algebra.

Example from stochastic processes:

- ▶  $\Omega := D([0, \infty); \mathbb{R})$  the space of càdlàg paths,
- ▶  $E_{0+}$  the equivalence relation of agreeing for some positive time,
- ▶  $\mathcal{F}_{0+} := \bigcap_{t>0} \mathcal{F}_t$  the *germ*  $\sigma$ -algebra.

Then  $E_{0+}$  is strongly dualizable, and we have  $E_{0+}^* = \mathcal{F}_{0+}$ , and

$$\sup_{A \in \mathcal{F}_{0+}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E_{0+})).$$


Example from stochastic processes:

- ▶  $\Omega := D([0, \infty); \mathbb{R})$  the space of càdlàg paths,
- ▶  $E_{0+}$  the equivalence relation of agreeing for some positive time,
- ▶  $\mathcal{F}_{0+} := \bigcap_{t>0} \mathcal{F}_t$  the *germ*  $\sigma$ -algebra.

Then  $E_{0+}$  is strongly dualizable, and we have  $E_{0+}^* = \mathcal{F}_{0+}$ , and

$$\sup_{A \in \mathcal{F}_{0+}} |\mathbb{P}(A) - \mathbb{P}'(A)| = \min_{\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')} (1 - \tilde{\mathbb{P}}(E_{0+})).$$

Consequence:  $(\mathbb{P}, \mathbb{P}')$  has the germ coupling property (GCP) if and only if  $\mathbb{P}(A) = \mathbb{P}'(A)$  for all  $A \in \mathcal{F}_{0+}$ .



This recovers the result of Ernst-Kendall-Roberts-Rosenthal for BMs with drift.

This recovers the result of Ernst-Kendall-Roberts-Rosenthal for BMs with drift. But we also get some other interesting consequences:

This recovers the result of Ernst-Kendall-Roberts-Rosenthal for BMs with drift. But we also get some other interesting consequences:

### **Theorem (AQJ)**

*If  $\mu : \mathbb{R} \rightarrow \mathbb{R}$  and  $\sigma : \mathbb{R} \rightarrow [0, \infty)$  are Lipschitz continuous and  $x_0 \in \mathbb{R}$  is arbitrary,*

This recovers the result of Ernst-Kendall-Roberts-Rosenthal for BMs with drift. But we also get some other interesting consequences:

### Theorem (AQJ)

*If  $\mu : \mathbb{R} \rightarrow \mathbb{R}$  and  $\sigma : \mathbb{R} \rightarrow [0, \infty)$  are Lipschitz continuous and  $x_0 \in \mathbb{R}$  is arbitrary, then the strong solution  $X$  of*

$$\begin{cases} dX_t = \mu(X_t)dt + \sigma(X_t)dB_t \text{ for } 0 \leq t \leq 1 \\ X_0 = x_0, \end{cases} \quad (1)$$

This recovers the result of Ernst-Kendall-Roberts-Rosenthal for BMs with drift. But we also get some other interesting consequences:

### Theorem (AQJ)

*If  $\mu : \mathbb{R} \rightarrow \mathbb{R}$  and  $\sigma : \mathbb{R} \rightarrow [0, \infty)$  are Lipschitz continuous and  $x_0 \in \mathbb{R}$  is arbitrary, then the strong solution  $X$  of*

$$\begin{cases} dX_t = \mu(X_t)dt + \sigma(X_t)dB_t \text{ for } 0 \leq t \leq 1 \\ X_0 = x_0, \end{cases} \quad (1)$$

*has the Brownian GCP if and only if*

This recovers the result of Ernst-Kendall-Roberts-Rosenthal for BMs with drift. But we also get some other interesting consequences:

### Theorem (AQJ)

*If  $\mu : \mathbb{R} \rightarrow \mathbb{R}$  and  $\sigma : \mathbb{R} \rightarrow [0, \infty)$  are Lipschitz continuous and  $x_0 \in \mathbb{R}$  is arbitrary, then the strong solution  $X$  of*

$$\begin{cases} dX_t = \mu(X_t)dt + \sigma(X_t)dB_t \text{ for } 0 \leq t \leq 1 \\ X_0 = x_0, \end{cases} \quad (1)$$

*has the Brownian GCP if and only if  $\sigma \equiv 1$  on a neighborhood of  $x_0$ .*

This recovers the result of Ernst-Kendall-Roberts-Rosenthal for BMs with drift. But we also get some other interesting consequences:

### Theorem (AQJ)

*If  $\mu : \mathbb{R} \rightarrow \mathbb{R}$  and  $\sigma : \mathbb{R} \rightarrow [0, \infty)$  are Lipschitz continuous and  $x_0 \in \mathbb{R}$  is arbitrary, then the strong solution  $X$  of*

$$\begin{cases} dX_t = \mu(X_t)dt + \sigma(X_t)dB_t \text{ for } 0 \leq t \leq 1 \\ X_0 = x_0, \end{cases} \quad (1)$$

*has the Brownian GCP if and only if  $\sigma \equiv 1$  on a neighborhood of  $x_0$ .*

How to do this explicitly, say, for BM and OU?

For many classes of Markov processes, it turns out that the GCP can be concretely interpreted in terms of some canonical data:

For many classes of Markov processes, it turns out that the GCP can be concretely interpreted in terms of some canonical data:

### Theorem (Hummel-AQJ)

*A pair of Lévy processes in  $\mathbb{R}^k$  with characteristic triplets  $(a, \Sigma, \mu)$  and  $(a', \Sigma', \mu')$  has the GCP if and only if*

- ▶  $\Sigma = \Sigma'$ ,
- ▶  $H^2(\mu, \mu') < \infty$ , where  $H^2(\cdot, \cdot)$  denotes the Hellinger distance, and
- ▶ if  $\Sigma = \Sigma' = 0$ , then  $a = a'$ .

For many classes of Markov processes, it turns out that the GCP can be concretely interpreted in terms of some canonical data:

### Theorem (Hummel-AQJ)

*A pair of Lévy processes in  $\mathbb{R}^k$  with characteristic triplets  $(a, \Sigma, \mu)$  and  $(a', \Sigma', \mu')$  has the GCP if and only if*

- ▶  $\Sigma = \Sigma'$ ,
- ▶  $H^2(\mu, \mu') < \infty$ , where  $H^2(\cdot, \cdot)$  denotes the Hellinger distance, and
- ▶ if  $\Sigma = \Sigma' = 0$ , then  $a = a'$ .

What about particle systems?

For many classes of Markov processes, it turns out that the GCP can be concretely interpreted in terms of some canonical data:

### Theorem (Hummel-AQJ)

*A pair of Lévy processes in  $\mathbb{R}^k$  with characteristic triplets  $(a, \Sigma, \mu)$  and  $(a', \Sigma', \mu')$  has the GCP if and only if*

- ▶  $\Sigma = \Sigma'$ ,
- ▶  $H^2(\mu, \mu') < \infty$ , where  $H^2(\cdot, \cdot)$  denotes the Hellinger distance, and
- ▶ if  $\Sigma = \Sigma' = 0$ , then  $a = a'$ .

What about particle systems? Processes from population genetics?

For many classes of Markov processes, it turns out that the GCP can be concretely interpreted in terms of some canonical data:

### Theorem (Hummel-AQJ)

*A pair of Lévy processes in  $\mathbb{R}^k$  with characteristic triplets  $(a, \Sigma, \mu)$  and  $(a', \Sigma', \mu')$  has the GCP if and only if*

- ▶  $\Sigma = \Sigma'$ ,
- ▶  $H^2(\mu, \mu') < \infty$ , where  $H^2(\cdot, \cdot)$  denotes the Hellinger distance, and
- ▶ if  $\Sigma = \Sigma' = 0$ , then  $a = a'$ .

What about particle systems? Processes from population genetics? For general Feller processes, can one formulate the GCP as an analytic relation between the generators?

Thank you!

General principle of mathematical optimization is that dual representations can lead one to characterizations of primal optimality:

General principle of mathematical optimization is that dual representations can lead one to characterizations of primal optimality:

### **Theorem (AQJ)**

*Suppose  $E$  is strongly dualizable, and  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . Then  $\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')$  is optimal if and only if*

General principle of mathematical optimization is that dual representations can lead one to characterizations of primal optimality:

### Theorem (AQJ)

*Suppose  $E$  is strongly dualizable, and  $\mathbb{P}, \mathbb{P}' \in \mathcal{P}(\Omega, \mathcal{F})$ . Then  $\tilde{\mathbb{P}} \in \Pi(\mathbb{P}, \mathbb{P}')$  is optimal if and only if the measures*

$$\tilde{\mathbb{P}}(\cdot \setminus E) \circ \pi^{-1} \quad \text{and} \quad \tilde{\mathbb{P}}(\cdot \setminus E) \circ (\pi')^{-1}$$

*are mutually singular on  $(\Omega, E^*)$ , where  $\pi, \pi : \Omega \times \Omega \rightarrow \Omega$  are the coordinate projections.*

How to construct a measurable equivalence relation which is not strongly dualizable?

How to construct a measurable equivalence relation which is not strongly dualizable?

Most likely, need a space that is not standard Borel.

Is strong dualizability closed under decreasing intersections?

Is strong dualizability closed under decreasing intersections? Ongoing work says yes.

Is strong dualizability closed under decreasing intersections? Ongoing work says yes.

The greedy construction of an optimal coupling does not work, instead need to use martingale arguments.

Is strong dualizability closed under decreasing intersections? Ongoing work says yes.

The greedy construction of an optimal coupling does not work, instead need to use martingale arguments.

Let  $(\Omega, \mathcal{F})$  be a standard Borel space and  $\mathcal{E}$  a collection of measurable equivalence relations satisfying the following:

Is strong dualizability closed under decreasing intersections? Ongoing work says yes.

The greedy construction of an optimal coupling does not work, instead need to use martingale arguments.

Let  $(\Omega, \mathcal{F})$  be a standard Borel space and  $\mathcal{E}$  a collection of measurable equivalence relations satisfying the following:

- (1) If  $E$  is smooth then  $E \in \mathcal{E}$ .

Is strong dualizability closed under decreasing intersections? Ongoing work says yes.

The greedy construction of an optimal coupling does not work, instead need to use martingale arguments.

Let  $(\Omega, \mathcal{F})$  be a standard Borel space and  $\mathcal{E}$  a collection of measurable equivalence relations satisfying the following:

- (1) If  $E$  is smooth then  $E \in \mathcal{E}$ .
- (2) If  $\{E_n\}_{n=1}^{\infty}$  are in  $\mathcal{E}$  and  $E_1 \subseteq E_2 \subseteq \dots$  then  $\bigcup_{n=1}^{\infty} E_n \in \mathcal{E}$ .

Is strong dualizability closed under decreasing intersections? Ongoing work says yes.

The greedy construction of an optimal coupling does not work, instead need to use martingale arguments.

Let  $(\Omega, \mathcal{F})$  be a standard Borel space and  $\mathcal{E}$  a collection of measurable equivalence relations satisfying the following:

- (1) If  $E$  is smooth then  $E \in \mathcal{E}$ .
- (2) If  $\{E_n\}_{n=1}^{\infty}$  are in  $\mathcal{E}$  and  $E_1 \subseteq E_2 \subseteq \dots$  then  $\bigcup_{n=1}^{\infty} E_n \in \mathcal{E}$ .
- (3) If  $\{E_n\}_{n=1}^{\infty}$  are in  $\mathcal{E}$  and  $E_1 \supseteq E_2 \supseteq \dots$  then  $\bigcap_{n=1}^{\infty} E_n \in \mathcal{E}$ .

Is strong dualizability closed under decreasing intersections? Ongoing work says yes.

The greedy construction of an optimal coupling does not work, instead need to use martingale arguments.

Let  $(\Omega, \mathcal{F})$  be a standard Borel space and  $\mathcal{E}$  a collection of measurable equivalence relations satisfying the following:

- (1) If  $E$  is smooth then  $E \in \mathcal{E}$ .
- (2) If  $\{E_n\}_{n=1}^{\infty}$  are in  $\mathcal{E}$  and  $E_1 \subseteq E_2 \subseteq \dots$  then  $\bigcup_{n=1}^{\infty} E_n \in \mathcal{E}$ .
- (3) If  $\{E_n\}_{n=1}^{\infty}$  are in  $\mathcal{E}$  and  $E_1 \supseteq E_2 \supseteq \dots$  then  $\bigcap_{n=1}^{\infty} E_n \in \mathcal{E}$ .

Then, how big is  $\mathcal{E}$ ? How much bigger is (1,2,3) than (1,2)?

To probabilists: The tail  $\sigma$ -algebra  $\mathcal{T}$  gives rise to the “simplest” non-trivial zero-one law. (folkloric, informal)

To probabilists: The tail  $\sigma$ -algebra  $\mathcal{T}$  gives rise to the “simplest” non-trivial zero-one law. (folkloric, informal)

To descriptive set theorists: The eventual equality  $E_0$  is the “simplest” non-trivial equivalence relation. (breakthrough, precise)

To probabilists: The tail  $\sigma$ -algebra  $\mathcal{T}$  gives rise to the “simplest” non-trivial zero-one law. (folkloric, informal)

To descriptive set theorists: The eventual equality  $E_0$  is the “simplest” non-trivial equivalence relation. (breakthrough, precise)

Use strong duality to connect the stories?