

SAT Math Equations

December 28, 2024

Contents

1	Quadratics	2
1.1	FOIL	2
1.2	Quadratic Formula	2
1.2.1	Derivation	2
1.2.2	Alternatives	2
1.3	Discriminant	3
1.4	Vertex	3
1.5	Factoring	3
1.5.1	Aquarc Method	3
2	Sets	3
2.1	Domain and Range	3
2.2	Standard Deviation	3
3	Word Problems	3

1 Quadratics

A quadratic is a function in the form

$$f(x) = ax^2 + bx + c$$

Where $\{a, b, c\} \in \mathbb{R}$.

Or in the vertex form:

$$f(x) = a(x - h)^2 + k$$

Where $\{a, h, k\} \in \mathbb{R}$

Or in the roots form:

$$f(x) = a(x - r_1)(x - r_2)$$

Where $\{a, r_1, r_2\} \in \mathbb{R}$

1.1 FOIL

FOIL stands for First Outer Inner Last. Meaning, when you have

$$f(x) = (a_1x - r_1)(a_2x - r_2)$$

You get:

$$a_1a_2x^2 - a_1r_2x - a_2r_1x + r_1r_2 = a_1a_2x^2 - (a_1r_1 + a_2r_2)x + (r_1r_2)$$

1.2 Quadratic Formula

1.2.1 Derivation

$$ax^2 + bx + c = 0 \implies ax^2 + bx = -c \implies \frac{1}{a}ax^2 + \frac{b}{a}x = -\frac{c}{a} \implies x^2 + \frac{b}{a}x = -\frac{c}{a}$$

Complete the square:

$$\begin{aligned} x^2 + \frac{b}{a}x + \frac{b^2}{4a^2} &= -\frac{c}{a} + \frac{b^2}{4a^2} \implies \left(x + \frac{b}{2a}\right)^2 = -\frac{4ac}{4a^2} + \frac{b^2}{4a^2} \implies \left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2} \\ \implies \sqrt{\left(x + \frac{b}{2a}\right)^2} &= \frac{\pm\sqrt{b^2 - 4ac}}{2a} \implies x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \end{aligned}$$

1.2.2 Alternatives

In some scenarios it may be easier to use systems of equations.

Given a quadratic in the form:

$$ax^2 + bx + c = 0$$

We want to find r_1 and r_2 such that:

$$r_1 + r_2 = b \quad r_1r_2 = c$$

You can easily substitute around until you get the answer. If $a \neq 1$ then it is difficult to use this formula.

1.3 Discriminant

1.4 Vertex

1.5 Factoring

You can use the two equations to do it

1.5.1 Aquarc Method

The fancy X thing

2 Sets

2.1 Domain and Range

2.2 Standard Deviation

3 Word Problems

You can't really predict what these will be. For

4 Trigonometry