

INDIAN INSTITUTE OF TECHNOLOGY  
BOMBAY  
MA205 Complex Analysis Autumn 2012

Anant R. Shastri

August 24, 2012

## Lecture 10

### Radius of Convergence

### Singularities

Removable Singularities

Poles

## Recall Limsup

- ▶ Given a sequence of real numbers  $\{b_n\}$ , let us recall what we mean by  $\limsup_n \{b_n\}$ .

# Recall Limsup

- ▶ Given a sequence of real numbers  $\{b_n\}$ , let us recall what we mean by  $\limsup_n \{b_n\}$ .
- ▶ Consider  $s_n = \sup\{b_n, b_{n+1}, \dots\}$ . Then  $s_n$  is a monotonically decreasing sequence. So, the following makes sense.

# Recall Limsup

- ▶ Given a sequence of real numbers  $\{b_n\}$ , let us recall what we mean by  $\limsup_n \{b_n\}$ .
- ▶ Consider  $s_n = \sup\{b_n, b_{n+1}, \dots\}$ . Then  $s_n$  is a monotonically decreasing sequence. So, the following makes sense.
- ▶ **Definition:**  $\text{Limsup}_n \{b_n\} := \lim_{n \rightarrow \infty} s_n$ .

## Recall Limsup

- ▶ This is also the same as the least upper bound of the set of limits of all convergent subsequences of  $\{b_n\}$ .

## Recall Limsup

- ▶ This is also the same as the least upper bound of the set of limits of all convergent subsequences of  $\{b_n\}$ .
- ▶ Two important properties of the Limsup are:

## Recall Limsup

- ▶ This is also the same as the least upper bound of the set of limits of all convergent subsequences of  $\{b_n\}$ .
- ▶ Two important properties of the Limsup are:
- ▶ (**Limsup-I**) If  $\alpha > \limsup_n \{b_n\}$  then there exists  $n_0$  such that for all  $n \geq n_0$  we have,  $b_n < \alpha$ .



# Recall Limsup

- ▶ This is also the same as the least upper bound of the set of limits of all convergent subsequences of  $\{b_n\}$ .
- ▶ Two important properties of the Limsup are:
- ▶ (**Limsup-I**) If  $\alpha > \limsup_n \{b_n\}$  then there exists  $n_0$  such that for all  $n \geq n_0$  we have,  $b_n < \alpha$ .
- ▶ (**Limsup-II**) If  $\beta < \limsup_n \{b_n\}$  then there exists infinitely many  $n_j$  such that  $b_{n_j} > \beta$ .

## Recall Limsup

- Indeed,  $\limsup$  can be characterized by these two properties.

## Recall Limsup

- ▶ It is important to note that  $\limsup$  always exists and can be any value in  $[-\infty, \infty]$ . When a sequence is convergent (including  $\pm\infty$ ) the  $\limsup_n$  of the sequence will be equal to the limit.

# Cauchy-Hadamard Theorem

## Definition

A formal power series  $P(t) = \sum_n a_n t^n$  is said to be *convergent* if there exists a **non zero number**  $z$  (real or complex) such that the series of complex numbers  $\sum_n a_n z^n$  is convergent.

# Cauchy-Hadamard Theorem

## Theorem

**(Cauchy-Hadamard)** Let  $P = \sum_{n \geq 0} a_n t^n$  be a power series over  $\mathbb{C}$ . Then

(a) there exists  $0 \leq R \leq \infty$  such that for all  $0 < r < R$ , the series  $P(z)$  is absolutely and uniformly convergent in  $|z| \leq r$  and for all  $|z| > R$  the series is divergent.

(b)  $\frac{1}{R} = \limsup \sqrt[n]{|a_n|},$

# Cauchy-Hadamard Theorem

## Theorem

**(Cauchy-Hadamard)** Let  $P = \sum_{n \geq 0} a_n t^n$  be a power series over  $\mathbb{C}$ . Then

(a) there exists  $0 \leq R \leq \infty$  such that for all  $0 < r < R$ , the series  $P(z)$  is absolutely and uniformly convergent in  $|z| \leq r$  and for all  $|z| > R$  the series is divergent.

(b)  $\frac{1}{R} = \limsup \sqrt[n]{|a_n|}$ ,

Convention:

$$\frac{1}{0} = \infty; \quad \frac{1}{\infty} = 0.$$

## Radius of Convergence:

- ▶ **Proof:** Indeed, all that we have to prove is to take  $R$  as given by (b) and show that it satisfies (a).

# Radius of Convergence:

- ▶ **Proof:** Indeed, all that we have to prove is to take  $R$  as given by (b) and show that it satisfies (a).
- ▶ So, let  $0 < r < R$ . Choose  $r < s < R$ . Then  $1/s > 1/R$  and hence by property (Limsup-I), we must have  $n_0$  such that for all  $n \geq n_0$ ,  $\sqrt[n]{|a_n|} < 1/s$ .



# Radius of Convergence:

- ▶ **Proof:** Indeed, all that we have to prove is to take  $R$  as given by (b) and show that it satisfies (a).
- ▶ So, let  $0 < r < R$ . Choose  $r < s < R$ . Then  $1/s > 1/R$  and hence by property (Limsup-I), we must have  $n_0$  such that for all  $n \geq n_0$ ,  
$$\sqrt[n]{|a_n|} < 1/s.$$
- ▶ Therefore, for all  $|z| \leq r$ ,  
$$|a_n z^n| < (r/s)^n, \quad n \geq n_0.$$

# Cauchy-Hadamard

- ▶ Since  $r/s < 1$ , by Weierstrass majorant criterion, it follows that  $P(z)$  is absolutely and uniformly convergent.

# Cauchy-Hadamard

- ▶ Since  $r/s < 1$ , by Weierstrass majorant criterion, it follows that  $P(z)$  is absolutely and uniformly convergent.
- ▶ Thus we have proved that inside the disc  $|z| < R$  the series is convergent.

## Cauchy-Hadamard Theorem:

- ▶ On the other hand, suppose  $|z| > R$ . We fix  $s$  such that  $|z| > s > R$ .

## Cauchy-Hadamard Theorem:

- ▶ On the other hand, suppose  $|z| > R$ . We fix  $s$  such that  $|z| > s > R$ .
- ▶ Then  $1/s < 1/R$ , and hence by (Limsup-II) there exist infinitely many  $n_j$ , for which  $\sqrt[n_j]{|a_{n_j}|} > 1/s$ .

# Cauchy-Hadamard Theorem:

- ▶ On the other hand, suppose  $|z| > R$ . We fix  $s$  such that  $|z| > s > R$ .
- ▶ Then  $1/s < 1/R$ , and hence by (Limsup-II) there exist infinitely many  $n_j$ , for which  $\sqrt[n_j]{|a_{n_j}|} > 1/s$ .
- ▶ This means that  $|a_{n_j} z^{n_j}| > (R/s)^{n_j}$ . It follows that the sequence  $\{a_n z^n\}$  is unbounded and hence the series  $\sum_n a_n z^n$  is divergent.



# Radius of Convergence:

## Definition

The number  $R$  obtained in the above theorem is called the *radius of convergence* of  $P(t)$ .

# Radius of Convergence:

## Definition

The number  $R$  obtained in the above theorem is called the *radius of convergence* of  $P(t)$ .

- ▶ The second part of the theorem gives you the formula for it. This is called the Cauchy-Hadamard formula.



# Radius of Convergence:

## Definition

The number  $R$  obtained in the above theorem is called the *radius of convergence* of  $P(t)$ .

- ▶ The second part of the theorem gives you the formula for it. This is called the Cauchy-Hadamard formula.
- ▶ Thus the collection of all points at which a given power series converges consists of an open disc centered at the origin and perhaps some points on the boundary of the disc.

# Radius of Convergence:

- ▶ Observe that if  $P(z)$  is convergent for some  $z$ , then from the last part of (a), the radius of convergence of  $P$  is at least  $|z|$ .

## Radius of Convergence:

- ▶ Observe that if  $P(z)$  is convergent for some  $z$ , then from the last part of (a), the radius of convergence of  $P$  is at least  $|z|$ .
- ▶ Also observe that the theorem does not say anything about the convergence of the series at points on the boundary  $|z| = R$ .

# Radius of Convergence:

## Theorem

*Any given power series, its derived series and its integrated series all have the same radius of convergence.*

# Newton's Binomial Series

## Example

*Consider the principal branch  $f(z)$  of  $(1+z)^\alpha$  in the open disc  $|z| < 1$ . The MacLaurin's series  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  is called the Newton's binomial series.*

# Newton's Binomial Series

## Example

Consider the principal branch  $f(z)$  of  $(1+z)^\alpha$  in the open disc  $|z| < 1$ . The MacLaurin's series  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  is called the Newton's binomial series. We have

$$a_n = \binom{\alpha}{n} = \frac{\alpha(\alpha-1)\cdots(\alpha-n+1)}{n!}.$$

## Newton's Binomial Series continued:

Since  $f(z)$  is holomorphic in  $|z| < 1$  we know that the radius of convergence  $r$  is at least 1.

## Newton's Binomial Series continued:

Since  $f(z)$  is holomorphic in  $|z| < 1$  we know that the radius of convergence  $r$  is at least 1.

If  $r > 1$  then  $(1+z)^\alpha$  is complex differentiable as many times as you like at  $z = -1$ .



## Newton's Binomial Series continued:

Since  $f(z)$  is holomorphic in  $|z| < 1$  we know that the radius of convergence  $r$  is at least 1.

If  $r > 1$  then  $(1+z)^\alpha$  is complex differentiable as many times as you like at  $z = -1$ .

which leads to a contradiction.

# Singularity

## ► Definition

Let  $U$  be a region in  $\mathbb{C}$ . If  $f(z)$  is a function on a subset of  $U$  then the points at which  $f$  is not defined or those points at which  $f$  is not holomorphic are referred to as **singularities**.

# Singularity

## ► Definition

Let  $U$  be a region in  $\mathbb{C}$ . If  $f(z)$  is a function on a subset of  $U$  then the points at which  $f$  is not defined or those points at which  $f$  is not holomorphic are referred to as **singularities**.

## ► Definition

*A point  $z \in U$  is called an **isolated singularity** of  $f$  if  $f$  is defined and holomorphic in a neighborhood of  $z$  except perhaps at  $z$ .*

# Examples

- ▶ (i) If  $p(z)$  is a polynomial, then  $1/p(z)$  has all its singularities isolated and these are nothing but the zeros of  $p(z)$ .

# Examples

- ▶ (i) If  $p(z)$  is a polynomial, then  $1/p(z)$  has all its singularities isolated and these are nothing but the zeros of  $p(z)$ .
- ▶ (ii) Since for any holomorphic function  $f$ , the zeros of  $f$  are isolated, it follows that all the singularities of  $1/f$  are isolated.

# Examples

- ▶ (i) If  $p(z)$  is a polynomial, then  $1/p(z)$  has all its singularities isolated and these are nothing but the zeros of  $p(z)$ .
- ▶ (ii) Since for any holomorphic function  $f$ , the zeros of  $f$  are isolated, it follows that all the singularities of  $1/f$  are isolated.
- ▶ (iii) Natural examples of holomorphic functions which have non isolated singularities are branches of logarithmic function and inverse-trigonometric functions. For instance,  $\text{Ln}(z)$  has singularities along the negative real

# Removable Singularities

- ▶ In obtaining Cauchy's theorem and integral formulae, in the previous chapter, we began with the hypothesis that the function under consideration is holomorphic throughout a domain  $U$ .

# Removable Singularities

- ▶ In obtaining Cauchy's theorem and integral formulae, in the previous chapter, we began with the hypothesis that the function under consideration is holomorphic throughout a domain  $U$ .
- ▶ Later on we weakened this hypothesis to include those functions  $f$  which are holomorphic on  $U_1 = U \setminus A$ , where  $A$  is a finite subset and



# Removable Singularities

- ▶ In obtaining Cauchy's theorem and integral formulae, in the previous chapter, we began with the hypothesis that the function under consideration is holomorphic throughout a domain  $U$ .
- ▶ Later on we weakened this hypothesis to include those functions  $f$  which are holomorphic on  $U_1 = U \setminus A$ , where  $A$  is a finite subset and
- ▶ satisfying the weaker condition that  $f$  is continuous at  $a \in A$ .

# Removable Singularities

- ▶ In obtaining Cauchy's theorem and integral formulae, in the previous chapter, we began with the hypothesis that the function under consideration is holomorphic throughout a domain  $U$ .
- ▶ Later on we weakened this hypothesis to include those functions  $f$  which are holomorphic on  $U_1 = U \setminus A$ , where  $A$  is a finite subset and
- ▶ satisfying the weaker condition that  $f$  is continuous at  $a \in A$ .
- ▶ (Or just bounded in nbd of  $a$ .)

# Removable Singularities

► Thus

$$f(w) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z - w} \quad (1)$$

is valid for all  $w \in \text{int } D \setminus A$ .

# Removable Singularities

- ▶ Thus

$$f(w) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z - w} \quad (1)$$

is valid for all  $w \in \text{int } D \setminus A$ .

- ▶ On the other hand, differentiation under the integral sign tells us that the function defined by the integral on the RHS of (1) is holomorphic in the interior of the disc. All that we need is the function  $f(z)$  is continuous on  $C = \partial D$ .

# Removable Singularities

- ▶ Thus

$$f(w) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z - w} \quad (1)$$

is valid for all  $w \in \text{int } D \setminus A$ .

- ▶ On the other hand, differentiation under the integral sign tells us that the function defined by the integral on the RHS of (1) is holomorphic in the interior of the disc. All that we need is the function  $f(z)$  is continuous on  $C = \partial D$ .
- ▶ Therefore, it follows that  $f$  is holomorphic even at points of  $A$ .

# Removable Singularities

Let us consolidate this simple observation.

# Removable Singularities

Let us consolidate this simple observation.

## Definition

An isolated singularity  $z_0$  of a holomorphic function is called a **removable singularity** if  $f(z_0)$  can be defined in such a way that  $f$  becomes complex differentiable at  $z_0$ , i.e., there exists a holomorphic function  $g : U \rightarrow \mathbb{C}$  such that for all  $z \in U \setminus \{a\}$  we have  $f(z) = g(z)$ .

# Removable Singularities

## Theorem

*Let  $U$  be a domain,  $a \in U$  be any point. Suppose  $f$  is holomorphic in  $U \setminus \{a\}$ . A necessary and sufficient condition that there exists a holomorphic function  $g : U \rightarrow \mathbb{C}$  such that  $g(z) = f(z), z \in U \setminus \{a\}$  is that  $f$  is **continuous at  $a$** .*



# Removable Singularity

- **Proof:** If such a  $g$  exists as stated in the definition, then

$$\lim_{z \rightarrow a} f(z) = \lim_{z \rightarrow a} g(z) = g(a)$$

exists.

# Removable Singularity

- ▶ **Proof:** If such a  $g$  exists as stated in the definition, then

$$\lim_{z \rightarrow a} f(z) = \lim_{z \rightarrow a} g(z) = g(a)$$

exists.

- ▶ Conversely, suppose the above limit exists. Take a circular region  $D$  around  $a$  contained in  $U$  and so that  $f$  is holomorphic in  $D \setminus \{a\}$ . It is enough to find a holomorphic function  $g$  on  $D$  such that  $f(z) = g(z)$  for all  $z \in D \setminus \{a\}$ .

# Removable Singularity

- ▶ But then C.I.F. says that

$$f(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(w)}{w - z} dw$$

for all  $z \in D$  and not equal to  $a$ .

# Removable Singularity

- ▶ But then C.I.F. says that

$$f(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(w)}{w - z} dw$$

for all  $z \in D$  and not equal to  $a$ .

- ▶ On the other hand, if we take  $g(z)$  as RHS, then we know that  $g$  is holomorphic function throughout  $D$ . (by differentiating under the integral sign). This completes the proof. ♠

# Removable Singularities: Examples

- ▶ Consider the function  $f(z) = \frac{\sin z}{z}$ ,  $z \neq 0$ .

# Removable Singularities: Examples

- ▶ Consider the function  $f(z) = \frac{\sin z}{z}$ ,  $z \neq 0$ .
- ▶ Obviously  $z = 0$  is an isolated singularity. We easily see that  $\lim_{z \rightarrow 0} f(z)$  exists. Hence,  $z = 0$  is a removable singularity. Also we see that  $\lim_{z \rightarrow 0} f(z) = 1$ . So we can define  $f(0) = 1$  and make  $f$  holomorphic at  $z = 0$  also.

# Removable Singularities: Examples

- ▶ Consider the function  $f(z) = \frac{\sin z}{z}$ ,  $z \neq 0$ .
- ▶ Obviously  $z = 0$  is an isolated singularity. We easily see that  $\lim_{z \rightarrow 0} f(z)$  exists. Hence,  $z = 0$  is a removable singularity. Also we see that  $\lim_{z \rightarrow 0} f(z) = 1$ . So we can define  $f(0) = 1$  and make  $f$  holomorphic at  $z = 0$  also.
- ▶ Other similar example are  $\frac{e^z - 1}{z}$ ,  $z \cot z$  etc. for which  $z = 0$  is a removable singularity.

# Removable Singularities

## Remark

One easy way a removable singularity  $z_0$  can arise is by taking a genuine holomorphic function  $f$  around this point and then brutally redefining the value of  $f$  to be something else only at  $z_0$  or merely pretending as if  $f$  is not defined at  $z_0$ .



# Poles

## ► Definition

Let now  $z = a$  be an isolated singularity of  $f$ . We say  $a$  is a pole of  $f$  if

$$\lim_{z \rightarrow a} |f(z)| = \infty.$$

# Poles

## ► Definition

Let now  $z = a$  be an isolated singularity of  $f$ . We say  $a$  is a pole of  $f$  if

$$\lim_{z \rightarrow a} |f(z)| = \infty.$$

## ► Theorem

*Let  $z = a$  be a pole of  $f(z)$  defined and holomorphic in  $U \setminus \{a\}$ . Then there exists a positive integer  $k$  such that in a disc around  $a$ ,  $\lim_{z \rightarrow a} (z - a)^k f(z)$  exists; (equivalently  $\lim_{z \rightarrow a} (z - a)^{k+1} f(z) = 0$ ).*

# Poles

- ▶ **Proof:** Since  $\lim_{z \rightarrow a} |f(z)| = \infty$  implies that there is  $\delta > 0$  such that  $f(z) \neq 0$  in  $B_\delta(a)$ .

# Poles

- ▶ **Proof:** Since  $\lim_{z \rightarrow a} |f(z)| = \infty$  implies that there is  $\delta > 0$  such that  $f(z) \neq 0$  in  $B_\delta(a)$ .
- ▶ Consider  $g(z) = 1/f(z)$  on  $B_\delta(a) \setminus \{a\} =: U_1$ . Then  $g(z)$  is holomorphic on  $U_1$ . Moreover,  $\lim_{z \rightarrow a} g(z) = 0$ .

# Poles

- ▶ **Proof:** Since  $\lim_{z \rightarrow a} |f(z)| = \infty$  implies that there is  $\delta > 0$  such that  $f(z) \neq 0$  in  $B_\delta(a)$ .
- ▶ Consider  $g(z) = 1/f(z)$  on  $B_\delta(a) \setminus \{a\} =: U_1$ . Then  $g(z)$  is holomorphic on  $U_1$ . Moreover,  $\lim_{z \rightarrow a} g(z) = 0$ .
- ▶ Hence  $z = a$  is a removable singularity of  $g(z)$ . We are forced to define  $g(a) = 0$  in order to obtain a holomorphic function on  $B_\delta(a)$ .

# Poles

- ▶ Suppose  $a$  is a zero of  $g$  of order  $k$ . Then  $g(z) = (z - a)^k h(z)$  for a holomorphic function  $h$  on  $U_1$  and  $h(a) \neq 0$ . Therefore,  $\lim_{z \rightarrow a} (z - a)^k f(z) = \lim_{z \rightarrow a} 1/h(z)$  exists. This implies  $\lim_{z \rightarrow a} (z - a)^{k+1} f(z) = 0$ .

# Poles

- ▶ Suppose  $a$  is a zero of  $g$  of order  $k$ . Then  $g(z) = (z - a)^k h(z)$  for a holomorphic function  $h$  on  $U_1$  and  $h(a) \neq 0$ . Therefore,  $\lim_{z \rightarrow a} (z - a)^k f(z) = \lim_{z \rightarrow a} 1/h(z)$  exists. This implies  $\lim_{z \rightarrow a} (z - a)^{k+1} f(z) = 0$ .
- ▶ A partial converse is also true.

# Poles

- ▶ Suppose  $a$  is a zero of  $g$  of order  $k$ . Then  $g(z) = (z - a)^k h(z)$  for a holomorphic function  $h$  on  $U_1$  and  $h(a) \neq 0$ . Therefore,  $\lim_{z \rightarrow a} (z - a)^k f(z) = \lim_{z \rightarrow a} 1/h(z)$  exists. This implies  $\lim_{z \rightarrow a} (z - a)^{k+1} f(z) = 0$ .
- ▶ A partial converse is also true.
- ▶ For, it follows that the function  $(z - a)^{k+1} f(z)$  is holomorphic around  $a$  and vanishes at  $a$ .



# Poles

- ▶ Suppose  $a$  is a zero of  $g$  of order  $k$ . Then  $g(z) = (z - a)^k h(z)$  for a holomorphic function  $h$  on  $U_1$  and  $h(a) \neq 0$ . Therefore,  $\lim_{z \rightarrow a} (z - a)^k f(z) = \lim_{z \rightarrow a} 1/h(z)$  exists. This implies  $\lim_{z \rightarrow a} (z - a)^{k+1} f(z) = 0$ .
- ▶ A partial converse is also true.
- ▶ For, it follows that the function  $(z - a)^{k+1} f(z)$  is holomorphic around  $a$  and vanishes at  $a$ .
- ▶ If the order of zero at  $a$  is  $m$  then we have  $(z - a)^{k+1} f(z) = (z - a)^m \alpha(z)$  for a holomorphic function  $\alpha$  with  $\alpha(a) \neq 0$ .

# Poles

- ▶ If  $m$  were bigger than or equal to  $k + 1$  then it follows that  $f(z) = (z - a)^{m-k-1}\alpha(z)$  and so  $z = a$  is a removable singularity.

# Poles

- ▶ If  $m$  were bigger than or equal to  $k + 1$  then it follows that  $f(z) = (z - a)^{m-k-1}\alpha(z)$  and so  $z = a$  is a removable singularity.
- ▶ If  $m < k + 1$  then

$$|f(z)| = \left| \frac{\alpha(z)}{(z - a)^{k-m+1}} \right| \rightarrow \infty$$

as  $z \rightarrow a$ .

# Poles

## ► Definition

The least such integer  $k$  is called the **order of the pole** of  $f$  at  $z = a$ .

# Poles

## ► Definition

The least such integer  $k$  is called the **order of the pole** of  $f$  at  $z = a$ .

## ► Definition

If the order is 1 then the pole is called a **simple pole**; if the order is bigger than 1, then the pole is called a **multiple pole**.

# Poles

- ▶ Indeed we have just proved that  $f(z) = (z - a)^{-k} h(z)$ , for all  $z$  in a neighborhood of  $a$ , where  $h(z)$  is holomorphic and  $h(a) \neq 0$ , where  $k$  is the order of the pole. The number  $k$  with this property is unique.

# Poles

- ▶ Indeed we have just proved that  $f(z) = (z - a)^{-k} h(z)$ , for all  $z$  in a neighborhood of  $a$ , where  $h(z)$  is holomorphic and  $h(a) \neq 0$ , where  $k$  is the order of the pole. The number  $k$  with this property is unique.

- ▶ Definition

A function which has all its singularities, if any, as poles, is called a *meromorphic* function in  $U$ .

[Observe that the poles of a meromorphic function are required to be isolated.]

# Poles: Examples

- ▶ (i) The simplest example of a function with a pole at  $z = 0$  of order  $k$  is  $1/z^k$ .



# Poles: Examples

- ▶ (i) The simplest example of a function with a pole at  $z = 0$  of order  $k$  is  $1/z^k$ .
- ▶ (ii) More generally, functions of the form  $\frac{P(z)}{Q(z)}$  where  $P, Q$  are polynomials are meromorphic functions.

# Poles: Examples

- ▶ (i) The simplest example of a function with a pole at  $z = 0$  of order  $k$  is  $1/z^k$ .
- ▶ (ii) More generally, functions of the form  $\frac{P(z)}{Q(z)}$  where  $P, Q$  are polynomials are meromorphic functions.
- ▶ (iii) Sums, products and scalar multiples of meromorphic functions are meromorphic.

# Poles: Examples

- ▶ (iv) If  $f$  and  $g$  are non zero meromorphic functions then so is  $f/g$ . Further, the zeros of  $g$  become poles of  $f/g$ , in general. However, if  $z = a$  is a common zero of  $f$  and  $g$ , it becomes a removable singularity of  $f/g$  provided the order of the zero of  $f$  at  $a$  is bigger than or equal to that of  $g$ .

## Poles: Examples

- ▶ (iv) If  $f$  and  $g$  are non zero meromorphic functions then so is  $f/g$ . Further, the zeros of  $g$  become poles of  $f/g$ , in general. However, if  $z = a$  is a common zero of  $f$  and  $g$ , it becomes a removable singularity of  $f/g$  provided the order of the zero of  $f$  at  $a$  is bigger than or equal to that of  $g$ .
- ▶ (v) A typical example of this type is  $(\sin z)/z$ , which is indeed a holomorphic function.

## Poles: Examples

- ▶ (iv) If  $f$  and  $g$  are non zero meromorphic functions then so is  $f/g$ . Further, the zeros of  $g$  become poles of  $f/g$ , in general. However, if  $z = a$  is a common zero of  $f$  and  $g$ , it becomes a removable singularity of  $f/g$  provided the order of the zero of  $f$  at  $a$  is bigger than or equal to that of  $g$ .
- ▶ (v) A typical example of this type is  $(\sin z)/z$ , which is indeed a holomorphic function.
- ▶ (vi) Amongst trigonometric functions we have  $\tan z$  and  $\cot z$  which have infinitely many poles

# Poles: Singular part

- ▶ Let  $f$  have a pole of order  $k$  at  $z = a$  and consider  $h(z) = (z - a)^k f(z)$ , and apply the Taylor's expansion to  $h(z) = b'_0 + b'_1(z - a) + \cdots + b'_{k-1}(z - a)^{k-1} + \phi(z)(z - a)^k$  where  $\phi_k$  is holomorphic at  $z = a$ .

# Poles: Singular part

- ▶ Let  $f$  have a pole of order  $k$  at  $z = a$  and consider  $h(z) = (z - a)^k f(z)$ , and apply the Taylor's expansion to  $h(z) =$

$$b'_0 + b'_1(z - a) + \cdots + b'_{k-1}(z - a)^{k-1} + \phi(z)(z - a)^k$$

where  $\phi_k$  is holomorphic at  $z = a$ .

- ▶ For  $z \neq a$ , we can divide this expression by  $(z - a)^k$  and write

$$b_1 = b'_{k-1}, b_2 = b'_{k-2}, \dots, b_k = b'_0, \text{ to obtain}$$

$$f(z) = \frac{b_k}{(z - a)^k} + \frac{b_{k-1}}{(z - a)^{k-1}} + \cdots + \frac{b_1}{(z - a)} + \phi(z)$$

- The sum of terms which involve  $b_j$  is called the *principal part* of  $f(z)$  at  $z = a$ . Observe that  $f$  minus its principal part is a holomorphic function.

Navigation icons: back, forward, search, and other controls.



- ▶ The sum of terms which involve  $b_i$  is called the *principal part* of  $f(z)$  at  $z = a$ . Observe that  $f$  minus its principal part is a holomorphic function.
- ▶ Further, if we write Taylor's expansion for  $\phi(z)$  on the rhs above we get *Laurent<sup>1</sup> expansion* for  $f(z)$ .