


ORIGINAL ARTICLE

Multiplex polymerase chain reaction detection of Shiga toxin genes and antibiotic sensitivity of *Escherichia coli* O157:H7 isolated from beef meat in Quetta, Pakistan

Abdul Samad¹ | Ferhat Abbas¹ | Zafar Ahmad¹ | Zunera Tanveer^{2,3} | Irshad Ahmad^{4,5} | Simon G. Patching⁴  | Nighat Nawaz⁶ | Muhammad Tauseef Asmat¹ | Abdul Raziq⁷ | Asadullah¹ | Muhammad Naeem¹ | Muhammad Aleem Akhtar^{1,8,9} | Olena Pokryshko¹⁰ | Mohammad Zahid Mustafa¹

¹Center for Advanced Studies in Vaccinology and Biotechnology, University of Balochistan, Quetta, Pakistan

²Department of Physiology, University Medical and Dental College, Faisalabad, Pakistan

³Institute of Molecular Biology and Biotechnology, The University of Lahore, Lahore, Pakistan

⁴School of Biomedical Sciences and Astbury Centre for Structural Molecular Biology, University of Leeds, Leeds, United Kingdom

⁵Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, Pakistan

⁶Department of Chemistry, Islamia College Peshawar, Peshawar, Pakistan

⁷Department of Statistics, University of Balochistan, Quetta, Pakistan

⁸Department of Pharmacy, University of Balochistan, Quetta, Pakistan

⁹Department of Physiology and Endocrinology, University of Balochistan, Quetta, Pakistan

¹⁰Department of Microbiology, Virology and Immunology, Ternopil State Medical University, Ukraine

Correspondence

Mohammad Z. Mustafa, Center for Advanced Studies in Vaccinology and Biotechnology, University of Balochistan, Sariab Road, Quetta, Pakistan.

Email: zunzah@gmail.com; Simon G. Patching, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, United Kingdom. Email: s.g.patching@leeds.ac.uk, simonpatching@yahoo.co.uk

Funding information

University of Balochistan, Grant/Award Number: NA

Abstract

The objective of this study was to conduct a qualitative analysis of raw beef meat sold in the city of Quetta, Pakistan for presence and drug sensitivity of the potentially pathogenic *Escherichia coli* strain O157:H7. The study used 200 raw beef meat samples collected from retail butcher shops. Conventional and rapid biochemical tests, latex agglutination and multiplex polymerase chain reaction (PCR) using primers designed for the *rfb*_{O157} and *fliC*_{H7} genes were used to detect *E. coli* O157:H7. All O157:H7 isolates were also tested for Shiga toxin genes *stx*₁ and *stx*₂. The prevalence of *E. coli* O157:H7 in collected beef meat samples was 10%. Detection through PCR was found more sensitive than detection of O and H antigens. The quantity of *E. coli* O157:H7 isolates positive for Shiga toxins was 50% (20% for *stx*₁, 45% for *stx*₂ and 10% for both *stx*₁ and *stx*₂). Season wise variation showed highest *E. coli* O157:H7 prevalence during summer months. A further concern is that *E. coli* O157:H7 isolates were resistant to a range of common antibiotics. The results indicate an urgent need for applying proper food hygiene practices in the Quetta region to reduce incidence of foodborne diseases and they also emphasize the global problem of antimicrobial resistance.

Practical applications

E. coli O157:H7 is as a potentially threatening foodborne pathogen. A significant prevalence of *E. coli* O157:H7 detected in raw beef meat from retail outlets in the city of Quetta indicates an urgent need for applying proper food hygiene practices in the Quetta region to reduce the incidence of foodborne diseases. Furthermore, resistance of the *E. coli* O157:H7 isolates to a range of commonly used antibiotics emphasizes the global problem of antimicrobial resistance. The multiplex PCR method used here is a reliable, sensitive, and relatively rapid technique for

detecting *E. coli* O157:H7 in food and environmental samples and important for ongoing surveillance to minimize contamination of raw meat products and associated cross contamination by *E. coli* O157:H7.

1 | INTRODUCTION

Foodborne infections have large effects on public health and on the economy worldwide (Havelaar et al., 2015; Hoffmann et al., 2017; Kirk et al., 2015; Yang, Lin, Aljuffali, & Fang, 2017). Indeed, microbial contamination of meat due to improper handling and processing poses great health risks around the globe (Ahmed & Sarangi, 2013; Antwi-Agyei & Maalekuu, 2014; Bakhtyari et al., 2016; Hassan Ali, Farooqui, Khan, Khan, & Kazmi, 2010; Jayathilakan, Sultana, Radhakrishna, & Bawa, 2012; Kim & Yim, 2017). Among foodborne pathogens, *Escherichia coli* O157:H7 is broadly categorized as a threatening pathogen (Armstrong, Hollingsworth, & Morris Jr., 1996; Bidet et al., 2005; Kiranmayi, Krishnaiah, & Mallika, 2010).

E. coli is a Gram-negative bacterium from the family Enterobacteriaceae that lives in the intestines of humans and other animals and does not usually cause any health risks. Some strains, however, have modified themselves to become equipped with toxigenic or pathogenic factors. By virtue of their pathogenic factors, such strains can cause several lethal diseases. On the basis of somatic 'H' (flagellar) antigens, 'O' (lipopolysaccharide) antigens and 'K' (capsular) antigens, *E. coli* are classified as different serotypes (Fratamico et al., 2016; González Garcia, 2002; Orskov & Orskov, 1992; Yang et al., 2016). Infections caused by pathogenic *E. coli* strains are wide-ranging depending on their serotypes and include: hemolytic uremic syndrome (HUS), urinary tract infection, neonatal meningitis, respiratory illnesses, and intestinal diseases. Pathogenic or diarrheagenic *E. coli* strains are divided into different pathotypes (STEC, EPEC, ETEC, EIEC, EAEC, and DAEC) depending on the nature of the disease caused by the specific strain: enteropathogenic (EPEC), Shiga toxin-producing/enterohemorrhagic (STEC/EHEC), Shigella/enteroinvasive (EIEC), enteroaggregative (EAEC), enterotoxigenic (ETEC), and diffusely adherent (DAEC) (Ibrahim, 2015).

E. coli serotype O157:H7 was discovered in 1982 (Riley et al., 1983), it has produced outbreaks worldwide and is ranked among the most studied foodborne pathogens (Adamu, Shamsul, Desa, & Khairani-Bejo, 2014; Bell et al., 1994; Ibrahim, 2015; Kieckens et al., 2015; Torso et al., 2015). Natural reservoirs of this pathogen include cattle, sheep, and goats, and the modes of transmission of the infections are animal to person, waterborne, foodborne, and person to person (Ateba & Mbewe, 2011; Ferens & Hovde, 2011). *E. coli* O157:H7 is within the class EHEC that produces Shiga toxin 1 (Stx1) and Shiga toxin 2 (Stx2), which are regulated by the genes *stx1/stx2* along with other virulence factors (intimin and enterohemolysin) (Parsons, Zelyas, Berenger, & Chui, 2016). Symptoms of *E. coli* O157:H7 infections include bloody diarrhea (hemorrhagic colitis), non bloody diarrhea, HUS, and thrombotic thrombocytopenic purpura (Griffin et al., 1988), which can be life-threatening if not treated successfully. Conventional

methods to detect *E. coli* O157:H7 are laborious, time consuming, and suffer from poor sensitivity (Deisingh & Thompson, 2004). To ensure food safety, rapid and reliable diagnostic tools are essential for detecting *E. coli* O157:H7 in different food products. Multiplex real-time polymerase chain reaction (PCR) has emerged as the most attractive alternative to culture-based and immunological-based methods for detecting *E. coli* O157:H7 (Jinneman, Yoshitomi, & Weagant, 2003; Li, Liu, & Wang, 2017; Tabashsum, Nazneen, Ahsan, Bari, & Yasmin, 2016; Van Giau, Nguyen, Nguyen, Le, & Nguyen, 2016; Zhou et al., 2017).

The main objective of this study was to estimate the prevalence of Shiga toxin producing *E. coli* O157:H7 in raw beef sold in Quetta, Pakistan by using biochemical tests, latex agglutination, and PCR methods. Another objective was to test the sensitivity of *E. coli* O157:H7 isolates to commonly available antibiotics.

2 | MATERIALS AND METHODS

2.1 | Sample collection

This study used 200 raw beef samples (100 samples each of diced and ground/minced) that were collected randomly during January 2016–August 2016 from 100 different retail butcher shops in Quetta, Pakistan. These shops were a minimum distance of 100–300 m from each other. Samples were stored in sterilized plastic containers in an ice box and brought to the Center for Advanced Studies in Vaccinology and Biotechnology (CASVAB) Bacteriology laboratory, University of Balochistan for immediate processing.

2.2 | Culture conditions and biochemical identification

Approximately 25 g of sample was transferred aseptically to modified trypticase soy broth (mTSB) (225 mL) and this was incubated at 37 °C for 24 hr. Later, non-sorbital fermenting (NSF) colonies were selected from MacConkey agar with sorbitol, cefixime, and tellurite (CT-SMAC) plates and streaked onto eosin methylene blue (EMB) agar plates, which were further incubated at 37 °C for 24 hr. On EMB, typical *E. coli* with a metallic shine were further characterized by biochemical tests (indole, methyl red, vogesproskauer, citrate, and lysine decarboxylase tests) as reported previously (Lee & Choi, 2006; You et al., 2006). For further confirmation that the isolated colonies were *E. coli*, the Remel RapID ONE System (Thermo Fisher Scientific, U.K.) was also used, which identifies medically important Enterobacteriaceae and other oxidase-negative Gram-negative bacilli based on enzyme technology.

TABLE 1 Oligonucleotide primers for PCR amplification of the genes *rfb*_{O157}, *flic*_{H7}, *stx*₁, and *stx*₂

Target gene	Primer sequence 5'–3'	Product size	Thermal cycle program	Reference
<i>rfb</i> _{O157}	F-CGGACATCCATGTGATATGG R-TTGCTATGTACAGCTAATCC	259 bp	Initial denaturation at 94 °C for 5 min followed by 35 cycles of denaturation at 94 °C for 1 min, annealing at 56 °C for 30 sec, elongation at 72 °C for 1 min and extension at 72 °C for 10 min.	Gannon et al. (1997)
<i>flic</i> _{H7}	F-GCGCTGTCGAGTTCTATCGAG R-CAACGGTGACTTTATCGCCATTCC	625 bp		
<i>stx</i> ₁	F-ACACTGGATGATCTCAGT GG R-CTGAATCCCCCTCCATTATG	614 bp	Initial denaturation at 94 °C for 5 min followed by 35 cycles of denaturation at 94 °C for 1 min, annealing at 60 °C for 1 min, elongation at 72 °C for 2 min and extension at 72 °C for 10 min.	Manna et al. (2006)
<i>stx</i> ₂	F-CCATGACAACGGACAGCAGTT R-CCTGTCAACTGAGCACTTGG	779 bp		

2.3 | Serotyping

Serotyping was performed by latex agglutination using a Remel-Wellcolex *E. coli* O157:H7 latex agglutination kit (Thermo Fisher Scientific) to detect both O157 (somatic antigen) and H7 (flagellar antigen) (Ewing, 1986). Isolates positive for the O157 antigen were further cultivated on blood agar overnight to detect the H7 antigen. Blue color agglutination indicates a positive result for the H antigen and red color agglutination indicates the O antigen.

2.4 | Oligonucleotide primers

The primers (Macrogen, South Korea) used for PCR amplification of the *rfb*_{O157} and *flic*_{H7} genes (Gannon et al., 1997) and the Shiga toxin *stx*₁/*stx*₂ genes (Manna, Brahmane, Manna, Batabyal, & Das, 2006) are shown in Table 1.

2.5 | DNA extraction and PCR conditions

The boiling method was used to extract DNA from each isolate (Radu et al., 2000). An Applied-Biosystems (U.K., 2720) PCR thermocycler was used and programmed using standard conditions (Chen & Griffiths, 1998). Following polymerization, the sample was loaded onto an agarose gel, which was run at 100 V for 35 min, and DNA bands were visualized using a Wealtec Dolphin-view system. A positive control of *E. coli* O157:H7 (already characterized and PCR confirmed) was obtained from the CASVAB seed bank and sterile distilled water was used as the negative control.

2.6 | Antibiotic sensitivity testing

Resistance or antibiotic sensitivity of the isolated strains was tested on Mueller–Hinton agar (MHA, oxoid) by the standard single disk method as described previously (Bauer, Kirby, Sherris, & Turck, 1966) and recently (Kassim, Omuse, Premji, & Revathi, 2016). Antibigrams of the isolates were critically examined.

Briefly, the *E. coli* O157:H7 inoculum was prepared in normal saline, its turbidity was matched according to 0.5 McFarland standards (McFarland, 1907; Swenson & Thornsberry, 1984) and was spread evenly on MHA plates. Using sterilized forceps, antibiotic disks were placed on the agar surface and the plates were then incubated at 37 °C for 24 hr. After incubation the plates were observed for

appearance of a zone of inhibition. The zone of inhibition was measured according to the diameter of the clear zone (no bacterial growth) around the antibiotic disks. The sensitivity and resistivity of *E. coli* O157:H7 against each antibiotic was evaluated from the zone of inhibition and categorized into three different groups: resistant (R), intermediate (I), and sensitive (S). The standard break points for sensitivity and resistance were adopted from Clinical and Laboratory Standards Institute (CLSI) standards (CLSI. M100-S22, 2012 [http://www.facm.ucl.ac.be/intranet/CLSI/CLSI-M100S22-susceptibility%20testing-2012-original.pdf]; CLSI. M100-S23., 2013 [http://www.facm.ucl.ac.be/intranet/CLSI/CLSI-M100S23-susceptibility-testing-2013-no-protection.pdf]).

3 | RESULTS

Two hundred raw beef samples (100 each of diced and ground/minced) were collected randomly from 100 different retail butcher shops in Quetta, Pakistan to check for the prevalence of *E. coli*. Importantly, only one diced and one minced beef sample was collected from each shop and these shops were a minimum distance of 100–300 m from each other, as they were scattered throughout the city of Quetta. Multiplex PCR results (Figure 1) showed that the prevalence of *E. coli* O157:H7 was 8% ($n = 8/100$) in diced samples and 12% ($n = 12/100$) in ground/minced samples. The overall prevalence of *E. coli* O157:H7 in both diced/minced beef meat samples was therefore 10% ($n = 20/200$). Conventional culture methods using primary enrichment on mTSB and plating showed 22 (11%) NSF colonies on CT-SMAC. When tested through the RapID One biochemical system, latex agglutination and PCR, two of the isolates were confirmed not to be O157 (Table 2). Two out of 20 isolates gave negative results for the H7 antigen by latex agglutination, but when tested using PCR, the H7 gene was detected. When multiplex PCR was used to test for the presence of Shiga toxin genes in the 20 O157:H7 isolates (Figure 2), 11/20 (55%) were positive for *stx*₁ or *stx*₂, 4/20 (20%) were positive for *stx*₁, 9/20 (45%) were positive for *stx*₂, and 2/20 (10%) were positive for both *stx*₁ and *stx*₂ Shiga toxins (Table 2). Month wise detection of *E. coli* O157:H7 in the beef samples showed highest prevalence during April–August, with the highest prevalence in June (Table 3).

Antibiotic sensitivity was tested for 21 antibiotics of different classes (Table 4) and results showed that all of the isolates were sensitive to aminoglycosides, fluoroquinolones, chloramphenicol, and imipenem. All




FIGURE 1 Agarose gel (2%) electrophoresis showing amplification fragments of the *rfb*_{O157} (259 bp) and *flic*_{H7} (625 bp) genes by multiplex PCR. Ground beef: Lanes 1–7 show positive amplification of *E. coli* O157:H7. Diced beef: Lanes 1–6 show positive amplification of *E. coli* O157:H7. M1/M2 = 50 bp DNA markers, +C = positive control, –C = negative control

TABLE 2 Detection of Shiga toxin producing *E. coli* O157:H7 in beef meat samples using culture, biochemical, serological, and PCR methods

Isolates	NSF on CT-SMAC	Biochemical confirmation by RapID one system	Latex agglutination		Multiplex PCR		Multiplex PCR Shiga toxins	
			O	H	<i>rfb</i> _{O157}	<i>flic</i> _{H7}	<i>stx</i> ₁	<i>stx</i> ₂
1	+	+	+	+	+	+	–	+
2	+	+	+	+	+	+	–	+
3	+	+	+	–	+	+	–	–
4	+	+	+	+	+	+	–	–
5	+	+	+	+	+	+	+	–
6	+	+	+	+	+	+	+	+
7	+	–	–	–	–	–	–	–
8	+	+	+	+	+	+	–	+
9	+	+	+	+	+	+	–	–
10	+	+	+	–	+	+	–	+
11	+	+	+	+	+	+	–	–
12	+	+	+	+	+	+	+	–
13	+	–	–	–	–	–	–	–
14	+	+	+	–	+	+	–	+
15	+	+	+	+	+	+	–	+
16	+	+	+	+	+	+	–	–
17	+	+	+	+	+	+	–	–
18	+	+	+	+	+	+	–	–
19	+	+	+	+	+	+	–	+
20	+	+	+	+	+	+	+	+
21	+	+	+	+	+	+	–	–
22	+	+	+	+	+	+	–	–




FIGURE 2 Agarose gel (2%) electrophoresis showing amplification fragments of the *stx*₁ (614 bp) and *stx*₂ (779 bp) genes by multiplex PCR. Lanes 1, 2, 7, 9, 12, 13, and 17 are positive for *stx*₂. Lanes 5 and 11 are positive for *stx*₁. Lanes 6 and 18 are positive for both *stx*₁ and *stx*₂. M1/M2 = 100 bp DNA markers, +C = positive control, –C = negative control

TABLE 3 Seasonal (month wise) prevalence of *E. coli* O157:H7 in beef meat samples

Sample type	Month								Total positive
	January	February	March	April	May	June	July	August	
Diced beef	0/10	0/12	0/13	2/13	1/13	2/13	2/13	1/13	8/100
Ground beef	0/10	1/12	1/13	1/13	2/13	4/13	2/13	1/13	12/100
Total	0/20	1/24	1/26	3/26	3/26	6/26	4/26	2/26	20/200

of the isolates were resistant to novobiocin, cephradine, bacitracin, clindamycin, metronidazole, sulfamethoxazole + trimethoprim, penicillins, and tetracyclines. Clarithromycin and nalidixic acid showed intermediate zone of inhibition (Table 4).

4 | DISCUSSION

In this study, two of the isolates did not exhibit H7 reaction in the latex agglutination test, which is in agreement with previous studies (Kim et al., 2005). These results are also in agreement with March and Ratnam (1986), who evaluated latex agglutination as an efficient and rapid test for the O157:H7 serotype. Interestingly, PCR assay for H7 gene in these isolates showed positive results. A possible explanation is that the O157:H7 strain did not express characteristically or the antigenic protein got saturated by host antibodies. Recent advances in diagnostic techniques have made several protocols available for isolation and characterization of *E. coli* O157:H7 from food, fecal, and environmental samples (Deisingh & Thompson, 2004; Tamerat, Muktar, & Shiferaw, 2016). Further, to classify and elaborate the function of the main virulence genes, such as *stx*₁ and *stx*₂, and genes encoding accessory STEC virulence factors, such as *eaeA*, *hlyA*, and *flic*_{H7},

different PCR assays have been described (Al-Ajmi et al., 2006; Gannon, King, Kim, & Thomas, 1992; Holland, Louie, Simor, & Louie, 2000; Kumar, Grover, & Kumar Batish, 2013; Pan, Chen, & Su, 2002; Sharma, 2006; Sharma & Dean-Nystrom, 2003).

Our findings are similar to those from other studies on beef meat samples reported from several countries around the world, in most of which the *stx*₂ gene was more prevalent than *stx*₁ in *E. coli* O157 isolates. For example, studies conducted on beef cattle meat in Argentina revealed a high presence of *stx*₂ in the *E. coli* O157 serotype (Blanco et al., 2004). Our findings to some extent are similar to previous studies by Kalin and Öngör (2014) who reported 16.6% *stx*₁ and 50.0% *stx*₂ in minced beef samples in Turkey. Sallam, Mohammed, Ahdy, and Tamura (2013) revealed the presence of 46.7% *stx*₁ and 86.7% *stx*₂ genes in *E. coli* O157 isolates from retail beef samples in Egypt. However, our findings are dissimilar to some studies that reported a higher prevalence of *stx*₁ than *stx*₂ (Kiranmayi et al., 2011). The prevalence of *E. coli* O157:H7 in beef meat samples from our study is comparable with the results of Heuvelink, Zwartkruis-Nahuis, Beumer, and de-Boer (1999), who reported a prevalence rate of 10.4% in beef from the Netherlands. The occurrence of *E. coli* O157:H7 in beef samples was reported as more than 8% in South Africa (Hajian, Rahimi, & Mommtaz, 2011), 6.8% in

TABLE 4 Antibiotic sensitivity of *E. coli* O157:H7 isolates from beef meat samples

No.	Antibiotic	Class	Result
1	Novobiocin (NV-30 µg)	Aminocoumarin	Resistant
2	Amikacin (AK-30 µg)	Aminoglycoside	Sensitive
3	Gentamicin (CN-10 µg)		Sensitive
4	Kanamycin (K-5 µg)		Sensitive
5	Tobramycin (TOB-10 µg)		Sensitive
6	Chloramphenicol (C-10 µg)	Amphenicol	Sensitive
7	Imipenem (IPM-10 µg)	Carbapenem (β-lactam)	Sensitive
8	Cephadrine (CE-30 µg)	Cephalosporin (β-lactam)	Resistant
9	Bacitracin (B-10 µg)	Cyclic peptide	Resistant
10	Enrofloxacin (ENR-5 µg)	Fluoroquinolone	Sensitive
11	Ofloxacin (OFX-5 µg)		Sensitive
12	Clindamycin (DA-2 µg)	Lincosamide	Resistant
13	Clarithromycin (CLR-2 µg)	Macrolide	Intermediate zone of inhibition
14	Metronidazole (MTZ-5 µg)	Nitroimidazole	Resistant
15	Amoxycillin + clavulanic acid (AMC-3 µg)	Penicillin (β-lactam)	Resistant
16	Ampicillin (AMP-2 µg)		Resistant
17	Methicillin (MET-5 µg)		Resistant
18	Nalidixic acid (NA-30 µg)	Quinolone	Intermediate zone of inhibition
19	Sulfamethoxazole + trimethoprim (SXT-25 µg)	Sulfonamide	Resistant
20	Oxytetracycline (OT-30 µg)	Tetracycline	Resistant
21	Tetracycline (TE-10 µg)		Resistant

China, 1.5–8% in Iran (Abdul-Raouf, Ammar, & Beuchat, 1996; Maktabi, Zarei, & Mohammadpour, 2016), 9% in raw beef in Saudi Arabia, 9% in Egypt, 8.2% in Thailand, and 3.7% in the U.S. (Hessain et al., 2015; Rahimi, Kazemeini, & Salajegheh, 2012).

Summer and fall are the seasons with highest prevalence of *E. coli* O157:H7 in meat samples, which are in agreement with the findings from previous studies on beef and sheep (Byrne et al., 2003; Elder et al., 2000; Rahimi et al., 2012). Moreover, our results showed presence of the H7 antigen in all of the O157 isolates tested, which is in accordance with previous studies reporting presence of the H7 antigen in isolates of human and bovine origin (Rahimi et al., 2012).

Our antibiotic sensitivity results are in agreement with previous studies reporting 100% resistance of *E. coli* O157:H7 isolates to trimethoprim, ampicillin, and amoxicillin/clavulanic acid followed by 80% resistance to tetracycline and 40% resistance to nalidixic acid (Fard, Bokaeian, & Qureishi, 2008). Further, our findings agree with this study that revealed sensitivity of *E. coli* O157:H7 to ciprofloxacin, imipenem, gentamycin, and amikacin (Fard et al., 2008). A further important observation is that the *E. coli* O157:H7 isolates were resistant to greater than three different classes of antimicrobials, providing evidence of multidrug resistance.

5 | CONCLUSIONS

Despite frequent contamination of raw meat products by potentially virulent *E. coli* O157:H7, to date, no clinical data about the impact on human health are available in Pakistan. Our study detected *E. coli* O157:H7 in raw beef meat from retail butcher shops in Quetta, Pakistan. Therefore, it is recommended that appropriate hygiene conditions, food safety measures, and improved surveillance are used in Quetta for minimizing contamination of such raw meat products and associated cross contamination by *E. coli* O157:H7. For this purpose, multiplex PCR is a reliable, sensitive, and relatively rapid technique for detecting *E. coli* O157:H7 in food and environmental samples. Of further concern is that the O157:H7 isolates were resistant to a range of commonly used antibiotics, thus emphasizing the global problem of antimicrobial resistance.

ACKNOWLEDGEMENT

This work was funded by the University of Balochistan (Quetta, Pakistan).

CONFLICT OF INTEREST

The authors declare that they have no potential conflicts of interest.

ORCID

Simon G. Patching  <http://orcid.org/0000-0001-6132-5317>

REFERENCES

- Abdul-Raouf, U. M., Ammar, M. S., & Beuchat, L. R. (1996). Isolation of *Escherichia coli* O157:H7 from some Egyptian foods. *International Journal of Food Microbiology*, 29, 423–426.
- Adamu, M. T., Shamsul, B. M. T., Desa, M. N., & Khairani-Bejo, S. (2014). A review on *Escherichia coli* O157:H7 – The super pathogen. *Health and the Environment Journal*, 5, 118–134.
- Ahmed, S. A., & Sarangi, S. K. (2013). Analysis of bacterial contamination in fresh and finished meat products and their molecular identification. *International Journal of Pharmaceutical Science Invention*, 2, 27–32.
- Al-Ajmi, D., Padmanabham, J., Denman, S. E., Gilbert, R. A., Al Jassim, R. A., & McSweeney, C. S. (2006). Evaluation of a PCR detection method for *Escherichia coli* O157:H7/H – Bovine faecal samples. *Letters in Applied Microbiology*, 42, 386–391.
- Antwi-Agyei, P., & Maalekuu, B. K. (2014). Determination of microbial contamination in meat and fish products sold in the Kumasi metropolis (a case study of Kumasi central market and the Bantama market). *Merit Research Journal of Agricultural Science and Soil Science*, 2, 38–46.
- Armstrong, G. L., Hollingsworth, J., & Morris, J. G., Jr. (1996). Emerging foodborne pathogens: *Escherichia coli* O157:H7 as a model of entry of a new pathogen into the food supply of the developed world. *Epidemiologic Reviews*, 18, 29–51.
- Ateba, C. N., & Mbewe, M. (2011). Detection of *Escherichia coli* O157:H7 virulence genes in isolates from beef, pork, water, human and animal species in the northwest province, South Africa: Public health implications. *Research in Microbiology*, 162, 240–248.
- Bakhtyari, F., Sayevand, H. R., Remely, M., Hippe, B., Hosseini, H., & Haslberger, A. G. (2016). Evaluation of bacterial contamination sources in meat production line. *Journal of Food Quality*, 39, 750–756.
- Bauer, A. W., Kirby, W. M., Sherris, J. C., & Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology*, 45, 493–496.
- Bell, B. P., Goldoft, M., Griffin, P. M., Davis, M. A., Gordon, D. C., Tarr, P. I., ... Kobayashi, J. (1994). A multistate outbreak of *Escherichia coli* O157:H7 – Associated bloody diarrhea and hemolytic uremic syndrome from hamburgers. The Washington experience. *Journal of the American Medical Association*, 272, 1349–1353.
- Bidet, P., Mariani-Kurkdjian, P., Grimont, F., Brahimi, N., Courroux, C., Grimont, P., & Bingen, E. (2005). Characterization of *Escherichia coli* O157:H7 isolates causing haemolytic uraemic syndrome in France. *Journal of Medical Microbiology*, 54, 71–75.
- Blanco, M., Padola, N. L., Krüger, A., Sanz, M. E., Blanco, J. E., González, E. A., ... Blanco, J. (2004). Virulence genes and intimin types of Shiga-toxin-producing *Escherichia coli* isolated from cattle and beef products in Argentina. *International Microbiology*, 7, 269–276.
- Byrne, C. M., Erol, I., Call, J. E., Kaspar, C. W., Buege, D. R., Hiemke, C. J., ... Luchansky, J. B. (2003). Characterization of *E. coli* O157:H7 from downer and healthy dairy cattle in the upper Midwest region of the United States. *Applied and Environmental Microbiology*, 69, 4683–4688.
- Chen, J., & Griffiths, M. W. (1998). PCR differentiation of *Escherichia coli* from other gram-negative bacteria using primers derived from the nucleotide sequences flanking the gene encoding the universal stress protein. *Letters in Applied Microbiology*, 27, 369–371.
- Deisingh, A. K., & Thompson, M. (2004). Strategies for the detection of *Escherichia coli* O157:H7 in foods. *Journal of Applied Microbiology*, 96, 419–429.
- Elder, R. O., Keen, J. E., Siragusa, J. R., Barkocy-Gallagher, G. A., Koohmaraie, M., & Laegreid, W. W. (2000). Correlation of enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides and carcasses of beef cattle during processing. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 2999–3003.
- Ewing, E. (1986). The genus *Escherichia*. In P. R. Edwards & W. H. Ewing (Eds.), *Edwards and Ewing's identification of Enterobacteriaceae* (4th ed., pp. 96–134). New York, NY: Elsevier Science Publishing.
- Fard, A. H., Bokaeian, M., & Qureishi, M. E. (2008). Frequency of *Escherichia coli* O157:H7 in children with diarrhea in Zahedan, Islamic Republic of Iran. *Eastern Mediterranean Health Journal*, 14, 1022–1027.
- Ferens, W. A., & Hovde, C. J. (2011). *Escherichia coli* O157:H7: Animal reservoir and sources of human infection. *Foodborne Pathogens and Disease*, 8, 465–487.
- Fratamico, P. M., DebRoy, C., Liu, Y., Needleman, D. S., Baranzoni, G. M., & Feng, P. (2016). Advances in molecular serotyping and subtyping of *Escherichia coli*. *Frontiers in Microbiology*, 7, 644.
- Gannon, V. P., D'Souza, S., Graham, T., King, R. K., Rahn, K., & Read, S. (1997). Use of the flagellar H7 gene as a target in multiplex PCR assays

- and improved specificity in identification of enterohemorrhagic *Escherichia coli* strains. *Journal of Clinical Microbiology*, 35, 656–662.
- Gannon, V. P., King, R. K., Kim, J. Y., & Thomas, E. J. (1992). Rapid and sensitive method for detection of Shiga-like toxin-producing *Escherichia coli* in ground beef using the polymerase chain reaction. *Applied and Environmental Microbiology*, 58, 3809–3815.
- González García, E. A. (2002). Animal health and foodborne pathogens: Enterohaemorrhagic O157:H7 strains and other pathogenic *Escherichia coli* virotypes (EPEC, ETEC, EIEC, EHEC). *Polish Journal of Veterinary Sciences*, 5, 103–115.
- Griffin, P. M., Ostroff, S. M., Tauxe, R. V., Greene, K. D., Wells, J. G., Lewis, J. H., & Blake, P. A. (1988). Illnesses associated with *Escherichia coli* O157:H7 infections. A broad clinical spectrum. *Annals of Internal Medicine*, 109, 705–712.
- Hajian, S., Rahimi, E., & Mommtaz, H. (2011). A 3-year study of *Escherichia coli* O157:H7 in cattle, camel, sheep, goat, chicken and beef minced meat. In *International conference on food engineering and biotechnology IPCBEE* (Vol. 9). Singapore: IACSIT Press.
- Hassan Ali, N., Farooqui, A., Khan, A., Khan, A. Y., & Kazmi, S. U. (2010). Microbial contamination of raw meat and its environment in retail shops in Karachi, Pakistan. *Journal of Infection in Developing Countries*, 4, 382–388.
- Havelaar, A. H., Kirk, M. D., Torgerson, P. R., Gibb, H. J., Hald, T., Lake, R. J., ... World Health Organization Foodborne Disease Burden Epidemiology Reference Group. (2015). World Health Organization global estimates and regional comparisons of the burden of foodborne disease in 2010. *PLoS Medicine*, 12, e1001923.
- Hessain, A. M., Al-Arfaj, A. A., Zakri, A. M., El-Jakee, J. K., Al-Zogibi, O. G., Hemeg, H. A., & Ibrahim, I. M. (2015). Molecular characterization of *Escherichia coli* O157:H7 recovered from meat and meat products relevant to human health in Riyadh, Saudi Arabia. *Saudi Journal of Biological Sciences*, 22, 725–729.
- Heuvelink, A. E., Zwartkruis-Nahuis, J. T., Beumer, R. R., & de-Boer, E. (1999). Occurrence and survival of verocytotoxin-producing *Escherichia coli* O157 in meats obtained from retail outlets in the Netherlands. *Journal of Food Protection*, 62, 1115–1122.
- Hoffmann, S., Devleesschauwer, B., Aspinall, W., Cooke, R., Corrigan, T., Havelaar, A., ... Hald, T. (2017). Attribution of global foodborne disease to specific foods: Findings from a World Health Organization structured expert elicitation. *PLoS One*, 12, e0183641.
- Holland, J. L., Louie, L., Simor, A. E., & Louie, M. (2000). PCR detection of *Escherichia coli* O157:H7 directly from stools: Evaluation of commercial extraction methods for purifying fecal DNA. *Journal of Clinical Microbiology*, 38, 4108–4113.
- Ibrahim, O. O. (2015). Food borne pathogen *Escherichia coli* O157:H7 history, sources of transmission, symptoms, detection and prevention. *EC Microbiology*, 2, 214–222.
- Jayathilakan, K., Sultana, K., Radhakrishna, K., & Bawa, A. S. (2012). Utilization of byproducts and waste materials from meat, poultry and fish processing industries: A review. *Journal of Food Science and Technology*, 49, 278–293.
- Jinneman, K. C., Yoshitomi, K. J., & Weagant, S. D. (2003). Multiplex real-time PCR method to identify Shiga toxin genes *stx1* and *stx2* and *Escherichia coli* O157:H7/H-serotype. *Applied and Environmental Microbiology*, 69, 6327–6333.
- Kalin, R., & Öngör, H. (2014). Examination of *Escherichia coli* O157:H7 and some virulence genes in marketed minced meat samples. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, 20, 957–960.
- Kassim, A., Omuse, G., Premji, Z., & Revathi, G. (2016). Comparison of clinical laboratory standards institute and European committee on antimicrobial susceptibility testing guidelines for the interpretation of antibiotic susceptibility at a university teaching hospital in Nairobi, Kenya: A cross-sectional study. *Annals of Clinical Microbiology and Antimicrobials*, 15, 21.
- Kieckens, E., Rybarczyk, J., De Zutter, L., Duchateau, L., Vanrompay, D., & Cox, E. (2015). Clearance of *Escherichia coli* O157:H7 infection in calves by rectal administration of bovine lactoferrin. *Applied and Environmental Microbiology*, 81, 1644–1651.
- Kim, J. H., & Yim, D. G. (2017). Microbial levels for food contact and environmental surfaces in meat processing plants and retail shops. *Food Science and Biotechnology*, 26, 299–302.
- Kim, J. Y., Kim, S. H., Kwon, N. H., Bae, W. K., Lim, J. Y., Koo, H. C., ... Park, Y. H. (2005). Isolation and identification of *Escherichia coli* O157:H7 using different detection methods and molecular determination by multiplex PCR and RAPD. *Journal of Veterinary Science*, 6, 7–19.
- Kiranmayi, C. B., Krishnaiah, N., & Mallika, E. N. (2010). *Escherichia coli* O157:H7 – An emerging pathogen in foods of animal origin. *Veterinary World*, 3, 382–389.
- Kiranmayi, C. B., Krishnaiah, N., Subhashini, N., Amaravathi, P., Maheswari, M., & Ramya, P. (2011). PCR analysis of mutton and chicken samples for the presence of Shiga toxigenic *E. coli*. *Archives of Clinical Microbiology*, 2, 2.
- Kirk, M. D., Pires, S. M., Black, R. E., Caipo, M., Crump, J. A., Devleesschauwer, B., ... Angulo, F. J. (2015). World Health Organization estimates of the global and regional disease burden of 22 foodborne bacterial, protozoal, and viral diseases, 2010: A data synthesis. *PLoS Medicine*, 12, e1001921.
- Kumar, A., Grover, S., & Kumar Batish, V. (2013). Application of multiplex PCR assay based on *uidR* and *fliCH7* genes for detection of *Escherichia coli* O157:H7 in milk. *Journal of General and Applied Microbiology*, 59, 11–19.
- Lee, J. H., & Choi, S. J. (2006). Isolation and characteristics of sorbitol-fermenting *Escherichia coli* O157 strains from cattle. *Microbes and Infection*, 8, 2021–2026.
- Li, B., Liu, H., & Wang, W. (2017). Multiplex real-time PCR assay for detection of *Escherichia coli* O157:H7 and screening for non-O157 Shiga toxin-producing *E. coli*. *BMC Microbiology*, 17, 215.
- Maktabi, S., Zarei, M., & Mohammadpour, H. (2016). Isolation and molecular characterization of non-sorbitol fermenting *Escherichia coli* isolated from fresh ground beef. *Jundishapur Journal of Health Sciences*, 8, e31195.
- Manna, S. K., Brahmane, M. P., Manna, C., Batabyal, K., & Das, R. (2006). Occurrence, virulence characteristics and antimicrobial resistance of *Escherichia coli* O157 in slaughtered cattle and diarrhoeic calves in West Bengal, India. *Letters in Applied Microbiology*, 43, 405–409.
- March, S. B., & Ratnam, S. (1986). Sorbitol-MacConkey medium for detection of *Escherichia coli* O157:H7 associated with hemorrhagic colitis. *Journal of Clinical Microbiology*, 23, 869–872.
- McFarland, J. (1907). Nephelometer: An instrument for media used for estimating the number of bacteria in suspensions used for calculating the opsonic index and for vaccines. *Journal of the American Medical Association*, 14, 1176–1178.
- Orskov, F., & Orskov, I. (1992). *Escherichia coli* serotyping and disease in man and animals. *Canadian Journal of Microbiology*, 38, 699–704.
- Pan, T. M., Chen, L. M., & Su, Y. C. (2002). Identification of *Escherichia coli* O157:H7 by multiplex PCR with primers specific to the *hlyA*, *eaeA*, *stx1*, *stx2*, *fliC* and *rfb* genes. *Journal of the Formosan Medical Association*, 101, 661–664.
- Parsons, B. D., Zelyas, N., Berenger, B. M., & Chui, L. (2016). Detection, characterization, and typing of Shiga toxin-producing *Escherichia coli*. *Frontiers in Microbiology*, 7, 478.
- Radu, S., Rusul, G., Ling, O. W., Purwati, E., Mustakim, M., & Lihan, S. (2000). Rapid isolation and detection of *Escherichia coli* O157:H7 by use of rainbow agar O157 and PCR assay. *Southeast Asian Journal of Tropical Medicine and Public Health*, 31, 77–79.
- Rahimi, E., Kazemine, H. R., & Salajegheh, M. (2012). *Escherichia coli* O157:H7/NM prevalence in raw beef, camel, sheep, goat, and water buffalo meat in Fars and Khuzestan provinces, Iran. *Veterinary Research Forum*, 3, 15–17.
- Riley, L. W., Remis, R. S., Helgeson, S. D., McGee, H. B., Wells, J. G., Davis, B. R., ... Cohen, M. L. (1983). Hemorrhagic colitis associated with a rare *Escherichia coli* serotype. *New England Journal of Medicine*, 308, 681–685.
- Sallam, K. I., Mohammed, M. A., Ahdy, A. M., & Tamura, T. (2013). Prevalence, genetic characterization and virulence genes of sorbitol-fermenting *Escherichia coli* O157:H- and *E. coli* O157:H7 isolated from retail beef. *International Journal of Food Microbiology*, 165, 295–301.
- Sharma, V. K. (2006). Real-time reverse transcription-multiplex PCR for simultaneous and specific detection of *rfbE* and *eae* genes of *Escherichia coli* O157:H7. *Molecular and Cellular Probes*, 20, 298–306.
- Sharma, V. K., & Dean-Nystrom, E. A. (2003). Detection of enterohemorrhagic *Escherichia coli* O157:H7 by using a multiplex real-time PCR assay for genes encoding intimin and Shiga toxins. *Veterinary Microbiology*, 93, 247–260.

- Swenson, J. M., & Thornsberry, C. (1984). Preparing inoculum for susceptibility testing of anaerobes. *Journal of Clinical Microbiology*, 19, 321–325.
- Tabashum, Z., Nazneen, M., Ahsan, C. R., Bari, M. L., & Yasmin, M. (2016). Influence of detection methods in characterizing *Escherichia coli* O157:H7 in raw goat meat using conventional and molecular methods. *Biocontrol Science*, 21, 261–264.
- Tamerat, N., Muktar, Y., & Shiferaw, D. (2016). Application of molecular diagnostic techniques for the detection of *E. coli* O157:H7: A review. *Journal of Veterinary Science and Technology*, 7, 362.
- Torso, L. M., Voorhees, R. E., Forest, S. A., Gordon, A. Z., Silvestri, S. A., Kissler, B., ... Harrison, L. H. (2015). *Escherichia coli* O157:H7 outbreak associated with restaurant beef grinding. *Journal of Food Protection*, 78, 1272–1279.
- Van Giau, V., Nguyen, T. T., Nguyen, T. K. O., Le, T. T. H., & Nguyen, T. D. (2016). A novel multiplex PCR method for the detection of virulence-associated genes of *Escherichia coli* O157:H7 in food. *3 Biotech*, 6, 5.
- Yang, H., Carlson, B., Geornaras, I., Woerner, D., Sofos, J., & Belk, K. (2016). Draft genome sequence of Shiga toxin-negative *Escherichia coli* O157:H7 strain C1-057, isolated from feedlot cattle. *Genome Announcements*, 4, e00049-16.
- Yang, S. C., Lin, C. H., Aljuffali, I. A., & Fang, J. Y. (2017). Current pathogenic *Escherichia coli* foodborne outbreak cases and therapy development. *Archives in Microbiology*, 199, 811–825.
- You, J. Y., Moon, B. M., Oh, I. G., Baek, B. K., Li, L. G., Kim, B. S., ... Lee, J. H. (2006). Antimicrobial resistance of *Escherichia coli* O157 from cattle in Korea. *International Journal of Food Microbiology*, 106, 74–78.
- Zhou, B., Liang, T., Zhan, Z., Liu, R., Li, F., & Xu, H. (2017). Rapid and simultaneous quantification of viable *Escherichia coli* O157:H7 and *Salmonella* spp. in milk through multiplex real-time PCR. *Journal of Dairy Science*, 100, 8804–8813.

How to cite this article: Samad A, Abbas F, Ahmad Z, et al. Multiplex polymerase chain reaction detection of Shiga toxin genes and antibiotic sensitivity of *Escherichia coli* O157:H7 isolated from beef meat in Quetta, Pakistan. *J Food Saf.* 2018; e12540. <https://doi.org/10.1111/jfs.12540>